

Data Evaluation Report on the Acute Toxicity of XDE-638 to Common Carp (*Cyprinus carpio*)

PMRA Submission Number {.....}

EPA MRID Number 45831009

## Data Requirement:

## PMRA DATA CODE

EPA DP Barcode D288160

OECD Data Point

EPA MRID 45831009

EPA Guideline §72-1

Test material: XDE-638

Purity: 97.5%

Common name: Penoxsulam

Chemical name: IUPAC: Not reported

CAS name: 2-(2,2-Difluoroethoxy)-N-(5,8-dimethoxy[1,2,4]triazolo[1,5-C]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide

CAS No.: Not reported

Synonyms: None reported

Primary Reviewer: Rebecca Bryan  
Staff Scientist, Dynamac CorporationSignature: Rebecca Bryan  
Date: 10/17/03QC Reviewer: Christie E. Padova  
Staff Scientist, Dynamac CorporationSignature: C.E. Padova  
Date: 10/17/03Primary Reviewer: <sup>Richard F. Erickson</sup> William Erickson - Biologist  
OPP/EFED/ERB - III

Date: 2/6/04

Secondary Reviewer(s):  
{EPA/OECD/PMRA}

Date:

Reference/Submission No.:

Company Code:

Active Code:

EPA PC Code: ~~199031~~

119031

Date Evaluation Completed:

**CITATION:** Marino, T.A., *et al.* 2001. Revised Report for XDE-638: An Acute Toxicity Study with the Common Carp, *Cyprinus carpio* Linnaeus. Unpublished study performed by Toxicology & Environmental Research and Consulting, The Dow Chemical Company, Midland, MI. Laboratory Study No. 001223R. Study submitted by Dow AgroSciences, LLC, Indianapolis, IN. Study initiated October 3, 2000 and completed February 1, 2001; revised reported completed September 10, 2001.



## EXECUTIVE SUMMARY:

In a 96-hour acute toxicity study, juvenile Common Carp (*Cyprinus carpio*) were exposed to XDE-638 (penoxsulam) at nominal concentrations of 0 (negative and solvent controls) and 100 ppm (limit test). Mean-measured concentrations were <10 (LOQ, controls) and 101 ppm a.i.

After 96 hours of exposure, no mortality or sub-lethal effects were observed in any control or test group. The 96-hour LC<sub>50</sub> was >101 ppm a.i., which categorizes XDE-638 as practically nontoxic to juvenile Common carp (*Cyprinus carpio*) on an acute toxicity basis. The NOAEC and LOAEC were 101 and >101 ppm a.i., respectively.

This study is scientifically sound. However, since the Common Carp is not recognized as an acceptable species for use in an acute toxicity study with freshwater fish (§72-1), this study is classified as SUPPLEMENTAL.

### Results Synopsis

Test Organism Size/Age (mean Weight or Length): Juvenile; 0.868 ± 0.195 g, 32 ± 2 mm (mean of all surviving fish at study termination)

Test Type (Flow-through, Static, Static Renewal): Static

#### 96-Hour

LC<sub>50</sub>: >101 ppm a.i.

NOAEC: 101 ppm a.i.

LOAEC: >101 ppm a.i.

Endpoints affected: None

## I. MATERIALS AND METHODS

**GUIDELINES FOLLOWED:** The study protocol was based on procedures outlined in the OECD Guidelines for Testing of Chemicals, No. 203 (1992); and the EEC Method C.1, Acute Toxicity for Fish (1992). Deviations from U.S. EPA guideline §72-1 included:

- The species used in this study was the Common Carp (*Cyprinus carpio*), which is not recognized as an acceptable species for use in the acute toxicity test to freshwater fish.
- The initial weight and length of the fish were not specified.
- The storage conditions of test chemical was not reported.
- The size of the test vessels (12 L with a fill volume of 10 L) was less than required (fill volume of 15-30 L).
- The water hardness (66 mg CaCO<sub>3</sub>/L) was slightly greater than recommended (40-48 mg CaCO<sub>3</sub>/L).
- The pH range (6.7-7.5) slightly exceeded recommendations (7.2-7.6).
- The instantaneous biomass loading rate (0.521 g/L) slightly exceeded the maximum limit (≤0.5 g/L) for species maintained at ≥17°C.

These deviations do not affect the validity of the study; however, this study does not fulfill guideline

requirements.

**COMPLIANCE:** Signed and dated GLP, Confidentiality, and Quality Assurance statements were provided. This study was conducted in accordance with the GLP standards of the OECD, EC, and U.S. EPA.

**A. MATERIALS:**

**1. Test Material** XDE-638 (penoxsulam)

**Description:** Pink powder

**Lot No./Batch No. :** ND05167938

**Purity:** 97.5%

**Stability of Compound Under Test Conditions:** The stability of the test substance in the dilution water during the course of the study was verified by analytical determination at 0 (96.2% of nominal) and 96 hours (105% of nominal; Table 3, p. 21).

**Storage conditions of test chemical:** Not reported.

*OECD requires water solubility, stability in water and light,  $pK_a$ ,  $P_{ow}$ , and vapor pressure of the test compound. OECD requirements were not reported.*

**2. Test organism:**

**Species:** Common Carp (*Cyprinus carpio* Linnaeus)

**Age at test initiation:** Juvenile

**Weight at test initiation:** Not provided; the weight of all surviving fish measured at test termination averaged  $0.868 \pm 0.195$  g

**Length at test initiation:** Not provided; the length of all surviving fish measured at test termination averaged  $32 \pm 2$  mm

**Source:** Osage Catfisheries, Inc., Osage Beach, MO.

**B. STUDY DESIGN:**

**1. Experimental Conditions**

a) Range-finding Study: A 4-day static range-finding study was conducted with common carp and XDE-638 at nominal concentrations of 0 (negative and solvent controls) and 100 ppm. The test solution was sonicated for approximately 4 hours to complete dissolution of test material prior to addition of fish. After 96 hours, the  $LC_{50}$  was  $>100$  ppm based on biological observations. The

definitive nominal test concentration was determined based on the range-finding results.

b) Definitive Study:

**Table 1: Experimental Parameters**

Parameter	Details	Remarks
		Criteria
Acclimation period:	At least 14 days prior to testing.	
Conditions: (same as test or not)	Same as test	
Feeding:	Standard diet once per day except during the 48 hours prior to testing.	<i>EPA requires: minimum 14 days; no feeding during test OECD requires minimum of 12 days.</i>
Health: (any mortality observed)	<5% mortality during the last 48 hours of acclimation.	
Duration of the test	96 hours	
		<i>EPA/OECD requires: 96 hours</i>
Test condition		
static/flow through	Static	
Type of dilution system- for flow through method.	N/A	<i>EPA: Must provide reproducible supply of toxicant, with a consistent flow rate of 5-10 vol/24 hours, and meter systems calibrated before study and checked twice daily during test period</i>
Renewal rate for static renewal	N/A	
Aeration, if any	No aeration during testing.	
		<i>EPA requires: no aeration; OECD permits aeration</i>
<u>Test vessel</u>		
Material: (glass/stainless steel)	Glass beakers	The test vessel size and fill volume were less than required.
Size:	12 L	
Fill volume:	10 L	<i>EPA requires: Size 19 L (5 gal) or 30 x 60 x 30 cm</i> <i>Fill volume: 15-30 L of solution</i>
Source of dilution water	The dilution water was pumped to the laboratory from the upper Saginaw Bay of Lake Huron. The	

Parameter	Details	Remarks
		Criteria
		EPA 1975; Soft reconstituted water or water from a natural source, <b>not</b> dechlorinated tap water; OECD permits dechlorinated tap water.
<u>Water parameters:</u> Hardness	66 mg CaCO <sub>3</sub> /L	The hardness was slightly higher than recommended.
pH	6.7-7.5	
Dissolved oxygen	5.7-10.5 mg/L (≥67%)	
Total Organic Carbon	2,400 ng/mL	
Particulate Matter	<LOD (<1000 ng/mL, total suspended solids)	
Metals	See Table 1, p. 19	<p><b>Hardness and pH</b> EPA requires hardness of 40-48 mg/L as CaCO<sub>3</sub> and pH of 7.2-7.6; 8.0-8.3 for marine-stenohaline fishes, 7.7-8.0 for estuarine-euryhaline fishes; monthly range &lt;0.8. OECD allows hardness of 10-250 mg/L as CaCO<sub>3</sub> and pH between 6 and 8.5.</p> <p><b>Dissolved Oxygen</b> <u>Renewal</u>: ≥60% during 1<sup>st</sup> 48 hrs and ≥ 40% during 2<sup>nd</sup> 48 hrs <u>Flow-through</u>: ≥60% through out test. OECD requires at least 80% saturation value.</p> <p><b>Temperature</b> EPA requires 12°C for coldwater species and 17-22°C for warmwater species. OECD requires range of 21 - 25°C for bluegill and 13-17°C for rainbow trout.</p> <p><b>Salinity</b> 30-34 ‰ (parts per thousand) salinity, weekly range &lt; 6 ‰</p> <p><b>EPA water quality</b> measured at beginning of test and every 48 hours</p>
Pesticides	<LOD (Table 2, p. 20)	
Chlorine	22 ppb	
Temperature	21.8-22.0°C	
{Salinity for marine or estuarine species}	N/A	
Intervals of water quality measurement	DO, pH, and temperature were determined daily. Temperature was continuously recorded from one test vessel.	
<u>Concentration of test material:</u> nominal:	0 (negative and solvent controls) and 100 ppm	

Parameter	Details	Remarks
		Criteria
measured:	<10 (LOQ, negative and solvent controls) and 101 ppm a.i.	Mean-measured concentrations are provided in Table 3, p. 21.  <i>EPA/OECD requires: Control and five treatment levels. Each conc. should be 60% of the next highest conc., and should be in a geometric series</i>
Solvent (type, percentage, if used)	N,N,-dimethyl formamide (DMF), 0.1 mL/L	<i>EPA requires: Not to exceed 0.5 mL/L for static tests or 0.1 mL/L for flow-through tests; OECD requires solvent, exceed 100 mg/L.</i>
<u>Number of fish/replicates:</u> negative control:  solvent control:  treated:	30 fish, divided into 6 replicates containing 5 fish each  30 fish, divided into 6 replicates containing 5 fish each  30 fish, divided into 6 replicates containing 5 fish each	<i>EPA: <math>\geq 10</math>/concentration; OECD requires at least 7 fish/concentration</i>
Biomass loading rate	0.521 g fish/L	<i>Static: <math>\leq 0.8</math> g/L at <math>\leq 17^\circ\text{C}</math>, <math>\leq 0.5</math> g/L at <math>&gt; 17^\circ\text{C}</math>; flow-through: <math>\leq 1</math> g/L/day; OECD requires maximum of 1 g fish/L for static and semi-static with higher rates accepted for flow-through</i>
Lighting	16-hours light/8-hours dark.	<i>EPA requires: 16 hours light/8 hours dark; OECD requires 12 -16 hours photoperiod.</i>
Feeding	Animals were not fed during testing.	<i>EPA/OECD requires: No feeding during the study</i>
Recovery of chemical Level of Quantitation Level of Detection	96.2-105% of nominal 10 mg/L Not reported.	Recoveries are based on test solutions analyzed on days 0 and 4 (Table 3, p. 21).

Parameter	Details	Remarks
		Criteria
Positive control {if used, indicate the chemical and concentrations}	N/A	
Other parameters, if any	N/A	

## 2. Observations:

Table 2: Observations

Criteria	Details	Remarks/Criteria
Parameters measured including the sublethal effects/toxicity symptoms	Mortality and sub-lethal effects	
Observation intervals	24, 48, 72 and 96 hours of exposure	<i>EPA/OECD requires: minimally every 24 hours</i>
Were raw data included?	Yes, sufficient	
Other observations, if any	N/A	

## II. RESULTS AND DISCUSSION:

### A. MORTALITY:

After 96 hours of exposure, no mortality were observed in controls or 101 ppm a.i. treatment group.

Table 3: Effect of XDE-638 on mortality of Common Carp (*Cyprinus carpio*).

Treatment, ppm a.i. measured and (nominal conc.)	No. of fish at start of study						
		0-24 Hours		48-72 Hours		96 Hours	
		No Dead	% mortality	No Dead	% mortality	No Dead	% mortality
Negative control	30	0	0	0	0	0	0
Solvent control	30	0	0	0	0	0	0
101 (100)	30	0	0	0	0	0	0
NOAEC (mortality)	101 ppm a.i.						
LC <sub>50</sub> (95% C.I.)	>101 ppm a.i.						
Positive control, if used mortality: LC <sub>50</sub> :	N/A	N/A	N/A	N/A	N/A	N/A	N/A

#### B. NON-LETHAL TOXICITY ENDPOINTS:

During the 96-hour study, no sub-lethal effects were observed in controls or 101 ppm a.i. treatment group.

#### C. REPORTED STATISTICS:

The 96-hour LC<sub>50</sub>, NOAEC, and LOAEC values were visually determined, based on observed mortality data.

##### 96-Hour

LC<sub>50</sub>: >101 ppm a.i.

NOAEC: 101 ppm a.i.

LOAEC: >101 ppm a.i.

Endpoints affected: None

#### D. VERIFICATION OF STATISTICAL RESULTS:

The 96-hour LC<sub>50</sub>, NOAEC, and LOAEC for mortality and sub-lethal effects could be determined visually.

##### 96-Hour

LC<sub>50</sub>: >101 ppm a.i.

NOAEC: 101 ppm a.i.

LOAEC: >101 ppm a.i.

Endpoints affected: None



**E. STUDY DEFICIENCIES:**

This study is scientifically valid. However, since this study was conducted using the Common Carp (*Cyprinus carpio*), this study does not fulfill guideline requirements for an acute toxicity study with a freshwater fish (§72-1).

**F. REVIEWER'S COMMENTS:**

The reviewer's conclusions were identical to the study authors.

**G. CONCLUSIONS:**

This study is scientifically sound. However, since the Common Carp (*Cyprinus carpio*) were used as the test species, this study does not satisfy the guideline requirements for an acute toxicity study with freshwater fish (§72-1). This study is classified as SUPPLEMENTAL. The LC<sub>50</sub> was >101 ppm a.i., XDE-638 (penoxsulam) as practically nontoxic to juvenile Common Carp (*Cyprinus carpio*) on an acute toxicity basis. The NOAEC (for mortality and sub-lethal effects) was 101 ppm a.i.

**96-Hour**

LC<sub>50</sub>: >101 ppm a.i.

NOAEC: 101 ppm a.i.

LOAEC: >101 ppm a.i.

Endpoints affected: None

**III. REFERENCES:**

Organisation for Economic Cooperation and Development (OECD). OECD Guidelines for Testing of Chemicals. Method 203, "Fish, Acute Toxicity Test", ISBN 92-64-12221-4. Adopted July, 1992.

Official Journal of the European Communities. European Economic Communities (EEC) Method C.1. Acute Toxicity Test for Fish. ISSN 0378-6978. December 1992.

EPA-FIFRA. Environmental Protection Agency. Hazard Evaluation Division, Standard Evaluation Procedure: Acute Toxicity Test for Fish. EPA-540/9-85-006. June 1985.

U.S. Environmental Protection Agency. Office of Pesticide and Toxic Substances. Pesticide Assessment Guidelines, Subdivision E, Hazard Evaluation: Wildlife and Aquatic Organisms. Guideline 72-1, Acute Toxicity Test For Freshwater Fish. EPA-540/09-87-198. December 1986.

OECD Series on Principles on Good Laboratory Practice and Compliance Monitoring, Number 1. OECD Principles on Good Laboratory Practice (as revised in 1997) ENV/MC/CHEM(98)17.

EC Directive 99/11/EC of 8 March 1999 (OJ No. L 77/8-21, 23/3/1999).

Environmental Protection Agency-FIFRA GLP; Title 40 CFR Part 160-Federal Insecticide, Fungicide and Rodenticide Act (FIFRA); Good Laboratory Practice Standards, Final Rule.

Dow AgroSciences Test Substance Assay. TSN101773, 20 May 1999.

The Dow Chemical Company, Research Sample Safety Data Sheet, XDE-638, 2 February 1999.

Smith A.J. Dow AgroSciences Certificate of Analysis for Test/Reference/Control Substances, Report FA & PC 993090, 20 May 1999.

E.L. McClymont, *et al.* Analytical Data for XDE-638: 21 Day Chronic Toxicity Test with the Daphnia, *Daphnia magna*, Straus, Study # 001018, August 28, 2000.