

3-18-91

File: 121011
Copy

DATA EVALUATION RECORD

1. Chemical: Clethodim
2. Test Material: Select 2.0 EC; 25.6% ai
3. Study Type: Honey bee acute contact LD50

Species tested: Apis mellifera

4. Study ID: Hoxter, K., and G. Smith. 1990. Select 2.0 EC: An acute contact toxicity study with the honey bee. Wildlife International Ltd. Project No. 263-120. Submitted by Valent U.S.A. Corp., Walnut Creek, CA. EPA Acc. No. 416851-07.

5. Reviewed By:

Allen W. Vaughan
Entomologist
EEB/EFED

Signature: Allen W. Vaughan
Date: 3-18-91

6. Approved By

Norman J. Cook
Supervisory Biologist
EEB/EFED

Signature: Norman J. Cook
Date: 3-18-91

7. Conclusions:

This study is scientifically sound, and shows Select 2.0 EC to be practically nontoxic to honey bees. In an acute contact test, the LD50 was determined to be greater than 100 micrograms per bee. This study fulfills the requirement for an acute contact toxicity test on honey bees with the formulated product.

8. Recommendations: N/A

9. Background: This study was submitted in response to an EEB requirement for formulated product testing.

10. Discussion of Individual Tests: N/A

Handwritten initials/signature

11. Materials and Methods:

Four days prior to initiation of the study, one frame containing honey bee pupae was selected from Wildlife International's hives. The frame was placed in an automatic incubator to allow adult emergence. All test bees were 1 to 4 days old at initiation of the study and were apparently healthy. On the day of study initiation, all bees that had emerged were immobilized with N₂ and at least 50 bees were placed into each test chamber.

Test chambers were rolled paper containers. Each container was covered with a plastic petri dish through which a glass vial containing 50% sugar water was inserted. This food source was available to the test bees throughout the study.

The photoperiod was eight hours of light per day. Test temperatures ranged from 22 to 23° C.

Five treatment levels, 13, 22, 36, 60, and 100 micrograms per bee, were tested along with a solvent control and a negative control. Two replicates were tested at each dosage, with 25 bees per replicate. The solvent control bees received a volume of acetone equal to the largest volume used during the test.

Recently collected bees were immobilized with N₂ to facilitate handling. Each bee was individually dosed with the appropriate test solution. Solvent control bees were dosed with acetone.

Observations on mortality and signs of toxicity were made twice on the day of initiation and once on Day 1 and Day 2 after dosing.

The mortality pattern in this study was not conducive to calculating the LD50 value. An LD50 was estimated by visual inspection of the mortality data.

12. Reported Results:

The study authors found that Select 2.0 EC was practically nontoxic to honey bees, with an LD50 greater than 100 ug per bee.

13. Study Authors' Conclusions/ QA Measures

48-hr. LD50 greater than 100 ug per bee (practically nontoxic).

14. Reviewer's Discussion and Interpretation of the Study

- A. Test Procedures: Procedures were in accordance with protocols recommended in the guidelines. There were no problems in this regard.
- B. Statistical Analysis: Due to the nature of the data, no analysis was conducted.
- C. Discussion/Results: Select 2.0 EC is practically nontoxic to honey bees.
- D. Adequacy of Study:
 - 1. Classification: Core
 - 2. Rationale: Guidelines protocol
 - 3. Reparability: N/A

15. Completion of One-Liner for Study: N/A

16. CBI Appendix: N/A