



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

File copy
Acetochlor

8/1/89

Tox R 007513

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

Subject: Acetochlor - Quantitative Risk Assessment, Two Year
Chronic/Oncogenicity Sprague-Dawley Rat Study

caswell no. 3B

From: Bernice Fisher, Biostatistician *Bernice Fisher 7/21/89*
Science Support Section
Science Analysis and Coordination Branch
Health Effects Division (H7509C)

To: Stephen C. Dapson Ph.D., Pharmacologist
Review Section II
Toxicology Branch I - Insecticide/Rodenticide Support
Health Effects Division (H7509C)

Thru: John A. Quest, Ph.D., Section Head *John A. Quest 8/1/89*
Science Support Section
Science Analysis and Coordination Branch
Health Effects Division (H/509C)

Summary

The unit risk, Q_1^* , of acetochlor is $10^{-2} (\text{mg/kg/day})^{-1}$ in human equivalents. This estimate of Q_1^* is based upon papillary adenomas (nasal turbinates) in both male and female Sprague-Dawley rats fed 0, 40, 200, and 1000 ppm.

Significant differential mortality with increasing doses of acetochlor did not occur in either sex.

Both sexes had a significant increasing trend in papillary adenomas (nasal turbinates) with dose increments of acetochlor and both sexes had a significant difference in the pair-wise comparison of controls and the highest dose group. See the memorandum on Acetochlor - Qualitative Risk Assessment from a Rat 2-Year Chronic/Oncogenicity Study, C.J. Nelson 9/7/88 for details.

Background

The Peer Review Committee meeting on acetochlor on 2/8/89 concluded that the chemical compound should be classified as a [B₂] carcinogen. In addition they recommended that the unit risk, Q_1^* , should be estimated from both male and female rat nasal turbinate papillary adenoma tumor rates.

Dose-Response Analysis

As a result of the Peer Review Committee's recommendation of the use of rat papillary adenomas for the estimation of Q_1^* and since there was no significant dose related mortality in either sex the calculation of the unit risk was made by the use of the Global86, Multi-Stage Model computer program of K. Crump. The unit risk calculated from the rat data in ppm doses was converted to rat mg/kg/day by the use of Lehman's Tables and then to human equivalents by the use of interspecies surface area adjustments as recommended by EPA Cancer Guidelines (1986).

The resultant estimate of Q_1^* is as follows:

| | Rat, Q_1^* (mg/kg/day) ⁻¹ | In Human Equivalents |
|---|--|-----------------------|
| Tumor - nasal turbinate papillary adenomas | | |
| male | 2.04×10^{-3} | 1.08×10^{-2} |
| female | 1.73×10^{-3} | 9.20×10^{-3} |

It is to be noted that Q_1^* is an estimate of the upper (95%) bound on risk and that (as stated in the EPA Guidelines) the "true value of the risk is unknown and that the lower limit of the risk may be as low as zero".

References

Howe, R.B., Crump, K.S. and Van Landingham, C. (1986)
A Computer Program to Extrapolate Quantal Animal
Toxicity Data to Low Doses (unpublished report), 25 pgs.