

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460OFFICE OF PREVENTION,
PESTICIDES AND TOXIC
SUBSTANCES

March 22, 2004

MEMORANDUMSUBJECT: Drinking water assessment for abamectin
PC Code: 122804FROM: Jim Carleton, Chemist *Jim Carleton* 3/22/04
Environmental Resources Branch II
Environmental Fate and Effects Division

Nelson Thurman, Peer Reviewer

THRU: Tom Bailey, PhD, Chief *Tom A. Bailey* 3/22/04
Environmental Resources Branch II
Environmental Fate and Effects DivisionTO: Tom Harris
Registration Division

DP Barcode: D297214, D297226

CONCLUSIONS

As requested, a screening assessment of estimated environmental concentrations (EECs) for abamectin in drinking water resulting from the proposed label uses for avocados, celeriac, citrus, cucurbits, fruiting vegetables, grapes, herb crops, hops, leafy vegetables, mint, pome fruits, stone fruit, strawberries, tree nuts, and tuberous roots and corm vegetables was performed. The product considered was Agri-Mek® 0.15 EC (EPA Reg.No. 100-898) containing the active ingredient abamectin, which itself is a mixture of avermectins containing $\geq 80\%$ avermectin B_{1a}



2012918

0-demethyl avermectin A_{1a}) and ≤20% avermectin B_{1b} (5-0-demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A_{1a})]. Screening models were used to determine estimated concentrations in ground water and surface water. Based on PRZM/EXAMS modeling, the acute surface water Expected Environmental Concentration (EEC) of abamectin for the use on strawberries in Florida is 0.48 µg/L. The chronic non-cancer and cancer surface water values for use in HED's drinking water assessment are 0.10 and 0.09 µg/L, respectively.

The estimated ground water concentration of abamectin after application of abamectin to strawberries is 0.00196 µg/L. This estimate was derived using the EFED model SCI-GROW, and assuming application at the the maximum annual rate of 0.125 lb a.i. per acre. Because SCI-GROW is a screening level model, we have only moderate confidence in this result.

Background

Abamectin (also known as avermectin) is the active ingredient in the miticide/insecticide Agrimek ® 0.15, which is proposed for control of a number of insect pests, specifically mites and leafminers in leafy vegetables, fruiting vegetables, and tree crops.

Groundwater and surface water monitoring data are not available to the Environmental Fate and Effects Division (EFED) for abamectin at this time. Screening models were used to determine estimated concentrations for abamectin in groundwater and surface water for the proposed uses. Of all the crops listed on the label, strawberries have the highest seasonal application rate, at four times 0.01875 lb a.i./acre. Based upon a recently reviewed plastic mulch runoff study conducted by the registrant on Florida strawberries (DP292938), the PRZM scenario for Florida strawberries was modified to include a curve number of 88, instead of the previous values of 95-97. PRZM and EXAMS were then run to generate EECs for crops with relatively high seasonal use rates, including strawberries. The strawberry EECs turned out to be the highest, and are therefore reported here for use in HED's drinking water assessment.

SURFACE WATER

This assessment replaces the one performed four years ago for abamectin on leafy and fruiting vegetables (DP Barcode 265145; dated 4/20/2000). PRZM-EXAMS simulations were conducted for abamectin use on strawberries to evaluate the cumulative probability distribution for peak and annual mean EECs. Previous modeling took the use of plastic mulch on Florida strawberries into account by using estimated high curve numbers in the PRZM scenario. The current assessment differs in that a measured curve number (value of 88) derived from a small plot runoff study conducted by the registrant was used in the scenario.

Surface Water Exposure Inputs for PRZM/EXAMS for Parent Abamectin		
MODEL INPUT VARIABLE	INPUT VALUE	COMMENTS
Application Rate (lbs ai/A)	0.01875	Current label (EPA Reg.No. 100-898)
Maximum No. of Applications	4	Current label
Application Interval (days)	7 (between applications 1 and 2, 3 and 4) 21 (between applications 2 and 3)	Current Label
K _d (mL/g)	50	MRID 40856301
Aerobic Soil Metabolic Half-life (days)	115	90% upper-bound confidence limit of mean half-life
Is the pesticide wetted-in?	No	Current label
Depth of Incorporation (in.)	0	Current label
Spray Drift	6.4%	For ground spray
Solubility (µg/L)	78	10x reported value
Aerobic Aquatic Metabolic Half-life (days)	230	No acceptable aerobic aquatic metabolism data were available. Therefore, since there were no data and the hydrolysis rate is stable, per current EFED guidance, use 2x aerobic soil metabolism half-life as input value.
Hydrolysis (pH 7) half-life (days)	0	Stable. No MRID available. Review dated 4/18/83.
Photolysis Half-life (days)	0.5	Dark-control adjusted half-life. Ku and Jacob, 1983, No MRID available, Review dated 3/28/84.

Drinking water EECs for parent abamectin on strawberries in Florida, incorporating the Index Reservoir and Percent Cropped Area adjustments, are presented below. Because strawberries are a minor use crop, as are other crops to which abamectin is applied in Florida, the regional PCA value of 0.38 was applied to these estimates.

Estimated drinking water concentrations to be used for exposure to Abamectin in drinking water derived from Surface Water			
Toxicity Endpoint	Model EEC Value ($\mu\text{g/L}$)	Use Modeled	PCA Modeled
Acute	0.48	Strawberries in Florida; 4 ground applications @ 0.01875 lb ai/A; application intervals of 7 days, 21 days, 7 days (per label instructions for strawberries)	The regional PCA factor of 0.38 was used to reflect strawberries and other minor use crops grown in the southeast.
Chronic non-cancer	0.10		
Cancer	0.09		

Ground Water

The SCI-GROW model is based on scaled ground water concentration from ground water monitoring studies, environmental fate properties (aerobic soil half-lives and organic carbon partitioning coefficients- K_{oc} 's) and application rates. The model is based on permeable soils that are vulnerable to leaching and on shallow ground water (10-30 feet). SCI-GROW version 2.3 (executable file dated 08/05/2003) was used to estimate concentrations of abamectin that could be found in drinking water derived from ground water, using the input values listed in the table below. EECs were not calculated for degradates of abamectin.

Ground Water Exposure Inputs for SCI-GROW for Parent Abamectin		
MODEL INPUT VARIABLE	INPUT VALUE	COMMENTS
Application Rate (lbs. ai/A)	0.01875 (strawberries)	Current label ((EPA Reg.No. 100-898)
Maximum No. of Applications	4 (Strawberries)	Current label.
K_{oc}	2,531	Lowest non-sand K_{oc} , of 2,531 in Three Bridges silt loam (1.22 % OC). Lowest K_{oc} was used since the K_{oc} 's differed by more than a factor of 3. MRID 40856301
Aerobic Soil Metabolic Half-life (days)	70	Mean of 70 days from individual half-lives of 34, 41, 72, and 131 days. Ku and Jacob, 1983, No MRID available, Review dated 3/28/84.

Results from the SCI-GROW screening model predict that the maximum chronic concentration of parent abamectin in shallow ground water is not expected to exceed 1.96 ng/L for the current maximum seasonal use rate on strawberries.