EEB REVIEW

| DATE: IN | 9-30-85 | OUT | 0 2 DEC 1985 | |
|---------------------|--|----------------------|---|--|
| FILE OR REG. NO | 279-GNLE | , 279-GNLG, | 279-GNLU | and the second seco |
| PETITION OR EXP. NO |). | | and the state of the | |
| DATE OF SUBMISSION | | 9-16-85 | | |
| DATE RECEIVED BY HE | D | 9-26-85 | | <u></u> |
| RD REQUESTED COMPLI | ETION DATE _ | 12-2-85 | | · |
| EEB ESTIMATED COMPI | | | | |
| RD ACTION CODE/TYPE | | | | |
| | | | | |
| TYPE PRODUCT(S) : 3 | I, D, H, F, | | | |
| DATA ACCESSION NO(S | 5) | | | |
| PRODUCT MANAGER NO | | | | |
| PRODUCT NAME(S) | | | | |
| | | | | |
| COMPANY NAME | FMC Corp | oration | | e - (|
| SUBMISSION PURPOSE | Submissi | on of data | in response to | o EEB's |
| | review | of fish ear | ly life stage | toxicity |
| | test_usi | ng rainbow | trout | · |
| SHAUGHNESSEY NO. | СНЕМІ | CAL, & FORM | ULATION | % A.I. |
| 125401 | 2-(2-chloro | phenyl)-met | hyl-4-4- | |
| | dimethyl-3- | isoxazol id i | none | |
| | | | | |
| | | | | |
| | the state of the s | | | |

XX 641

EEB REVIEW

Command Herbicide

100 Submission Purpose and Label Information

100.1 Submission Purpose and Pesticide Use

The registrant (FMC Corp.) submitted data from a freshwater fish early life-stage test in support of registration of Command for use on soybeans.

101 Hazard Assessment

101.2 Likelihood of Adverse Effects on Nontarget Organisms

Freshwater fish

On the basis of data reviewed previously (LC50 = 19 mg/l for rainbow trout; LC50 = 34 mg/l for bluegill sunfish), EEB did not anticipate a hazard to freshwater fish from the proposed use of Command on soybeans. For this reason, a freshwater fish early life-stage test was not required. FMC Corp. did conduct the test, however, and data from that test indicate the MATC for Command technical on rainbow trout is between 2.29 mg/l and 4.35 mg/l. The MATC would be approximately 50 to 90 times the value for the aquatic estimated environmental concentration (0.05 ppm) calculated by the Exposure Assessment Branch for use of Command on soybeans. These figures support EEB's original conclusion, that use of Command on soybeans at proposed rates will not result in hazard to freshwater fish.

101.4 Adequacy of Toxicity Data

The registrant submitted a fish early life-stage test on rainbow trout. The study was determined to be supplemental, due to significant deviations from the recommended protocol.

103 Conclusions

EEB has reviewed the submitted fish early life-stage test on rainbow trout. As noted above (Sec. 101.4), the study was determined to be supplemental. The data will be placed in the EEB file for future reference.

Allen W. Vaughan, Entomologist Ecological Effects Branch Hazard Evaluation Division (TS-769)

Norman Cook, Section Head Ecological Effects Branch Hazard Evaluation Division (TS-769)

Michael Slimak, Chief

Ecological Effects Branch

Hazard Evaluation Division (TS-769)

DATA EVALUATION RECORD

- 1. Chemical: Command (FMC 57020)
- 2. Test Material: Technical, 95.6% a.i.
- 3. Study Type: Chronic fish early life-stage toxicity test (freshwater)

Species Tested: Rainbow trout (Salmo gairdneri)

- 4. Study ID: Anon. 1985. The toxicity of FMC 57020 to rainbow trout (Salmo gairdneri) embryos and larvae. (Submission of requested raw data.) Prepared by Springborn Bionomics, Inc., Wareham, MA. Submitted by FMC Corp., Princeton, NJ. Reg. Nos. 279-GNLE, 279-GNLG, 279-GNLU. Acc. No. 073830.
- 5. Reviewed By:

Allen W. Vaughan Entomologist EEB/HED Signature: <u>Allen W. Vaughan</u>

Date: 11/26/85

6. Approved By:

Norman J. Cook Supervisory Biologist EEB/HED Signature:

ate: $//.\dot{u}.$

7. Conclusions:

This study is scientifically sound, and determines that the maximum acceptable toxicant concentration (MATC) for Command technical to rainbow trout is between 2.29 mg/l and 4.35 mg/l.

The study does not fulfill the guideline requirement for a fish early life-stage toxicity test, for the following reasons:

- 1. test was initiated with "eyed" eggs which were exposed to the toxicant for only 6-8 days prior to hatch. 10 days would be the minimum acceptable period of exposure in this test.
- 2. test duration was only 50 days post-hatch. Sixty days would be the appropriate minimum duration.
- 8. Recommendations: N/A

- 9. Background: This submission, containing raw data from a fish early life-stage toxicity test with the technical pesticide, was submitted by the registrant in response to a request from EEB. EEB had reviewed the study previously and had determined the need for raw data to validate the study.
- 10. Discussion of Individual Test: N/A

11. Materials and Methods

- A. <u>Test Animals</u> were rainbow trout embryos obtained from Mount Lassen Trout Farm, Red Bluffs, CA.
- B. <u>Test System</u>: glass aquaria with constant test volume of 11 1 well water; flow-through exposure using proportional diluter; test duration = 57 days.
- C. <u>Dose</u>: Flow-through bioassay using measured concentrations; dimethyl formamide solvent.
- D. Design: Five concentrations (nominal = 1.27, 2.66, 5.31, 10.62, 21.24 mg/l; measured = 1.12, 2.29, 4.35, 8.45, 18.18 mg/l) plus control and solvent control (29 ul/1 DMF); 2 reps. of each, 40 larvae/rep.

E. Statistics:

Percentage hatchability of embryos and survival, length, and weight of larvae after 49 days post-hatch exposure, were subjected to analysis of variance (Steel and Torrie, 1960). Data for percentage hatchability and survival were transformed to arc sin percentage prior to analysis. If treatment effects were indicated, the means of these parameters were compared to those from the control and solvent control using Dunnett's procedure (Steel and Torrie, 1960). When a treatment mean was significantly different from the control means (P=0.05), that treatment was considered to be a toxicant effect level.

12. Reported Results:

All rainbow trout exposed to 18.18 mg/l died as embryos (prior to hatching) or as larvae within 24 hours of hatching. Hatchability of embryos exposed to 8.45 mg/l or less was unaffected. No rainbow trout larvae survived 49 days of post-hatch exposure to 8.45 mg/l FMC 57020. Exposure to 4.35 mg/l of FMC 57020 for 57 days resulted in significantly decreased survival, mean total length and average wet weight in comparison to the controls.

Rainbow trout larvae exposed to measured FMC 57020 concentrations of 2.29 mg/l or less suffered no apparent adverse effects. The maximum acceptable toxicant concentration (MATC) was, therefore, determined to be between 2.29 mg/l and 4.35 mg/l. Data on embryo hatchability, survival, mean total length, and average wet weight are attached.

13. Study Author's Conclusions/Q.A. Measures:

Maximum acceptable toxicant concentration (MATC): 2.29 mg/l < MATC < 4.35 mg/l.

Test was conducted according to the Springborn Bionomics protocol entitled "Methods for conducting early life-stage toxicity tests with rainbow trout (Salmo gairdneri)," 1983. The data and report were conducted in accordance with all pertinent EPA Good Laboratory Practice regulations. Also, the report was reviewed by FMC Corp., Toxicology Department's Quality Assurance Unit.

14. Reviewer's Discussion and Interpretation of the Study

- A. <u>Test Procedures</u>: Following were the major deviations from recommended procedure:
 - test was initiated with "eyed" eggs (18 days postfertilization). Standard procedure indicates that fertilized eggs may be used within 48 hours of spawning;
 - range of time-to-hatch was only 6-8 days, as opposed to 10 days minimum;
 - 3. test duration was only 50 days post-hatch, instead of 60 days.

B. Statistical Analysis:

EEB's independent validation of the analyses performed by the authors indicates that the procedures used were appropriate. Results of EEB's validation support the reported results of the study. Copies of EEB's validation are attached to this review.

C. Discussion/Results:

This study is scientifically sound, and determines that the MATC for Command technical to rainbow trout is between 2.29 and 4.35 mg/l. These figures would probably be slightly lower if the methodology had adhered more closely to the recommended protocol (i.e., longer period of exposure prior to hatch, and longer period of exposure overall). It is because of these deviations that the study is considered supplemental.

D. Adequacy of Study:

- 1. Classification: Supplemental
- 2. Rationale: Exposure period prior to hatch is too brief; overall test duration is too short.
- 3. Reparability: None
- 15. Completion of One-Liner: One-liner completed 11/22/85.
- 16. CBI Appendix: N/A

Matchability of embryos and survival, total length and wet weight of rainbow trout (Salmo gairdneri) larvae exposed to FMC 57020 for 57 days. Table 4.

| Mean measured concentrations (mg/L) | Embryo hatchability (%) | Survival (%) | Larvae (30 days post-swim-up) Mean total length (S.D) (mm) | Average wet weight (mg) |
|---|-------------------------------|-----------------|--|-------------------------------|
| 18.18 A | . bg | | 1 | 1 |
| æ | 0 ه | 1 | } | ı |
| 8.45 A | 72 | в 0 | I | 1 |
| Ø | . 72 | 0 8 | 1 | |
| 4.35 A | 81 | .82a | 38(2) ^a | 474ª |
| æ | 91 | 80a | 37(3) ^a | 452 ^a |
| 2.29 A | 78 | . 86 | 39 (2) | 571 |
| B | 72 | 86 | 38(2) | 532 |
| 1.12 A | 68 | 86 | 40 (2) | 532 |
| £ | 80 | 95 | 39 (2) | 563 |
| control A | 83 | 92 | 39 (2) | 586 |
| В | 7.7 | 9.5 | 40(1) | 529 |
| solvent A | 82 | 88 | 40 (3) | 575 |
| control B (DMF) | . 88 | 8.5 | 41(2) | 625 |

aSignificantly (P=0.05) different'from control.

| 10:03 WEDNESDAY, NOVEMBER 20, 1985 1 | | | | | 10:03 WEDNESDAY, NOVEMBER 20, 1985 2 | | | R-SQUARE C.V. | 0.969523 8.4520 | RESPONSE MEAN | 53,78058833 | S F VALUE PR > F | 13803.27073720 111.34 0.0001 10:03 WEDNESDAY, NOVEMBER 20, 1985 3 | | | | | | | | | | | | | 10:03 WEDNESDAY, NOVEMBER 20, 1985 4 | VARI ANCE C.V. |
|--------------------------------------|---------------------------------|-------------------------|----------|-------|---|---------------------------------|------------------------------|----------------|-----------------|---------------|----------------------|------------------|--|---------------------------------|--|-------------------|--|--|-----------|--------|-----------------|---------------|---------------|--------|--------|--------------------------------------|----------------|
| 10:03 WEDNESD | | | | | 10:03 WEDNESD | | | PR > F | 0.0001 | ROOT MSE | 4.54553496 | TYPE III SS | 13803.2707372 10:03 WEDNESD | | or rate, | | 6 7.49028 7.5756 | SRENT. | | | | | | | | 10:03 WEDNESD | SUM |
| | URE | | | ហ | ET = 28 | URE | | F VALUE | 111.34 | | | DE | v | URE | RESPONSE SONWISE ERR | 619 | 5 7.38102 7.49 | CANTLY DIFF | N | 4 3 | 4 A | 4 8 | 4 U | 4 C | 4 F | 4 0 ` | STD FRROR |
| SAS | GENERAL LINEAR MODELS PROCEDURE | CLASS LEVEL INFORMATION | VALUES | ABCDE | NUMBER OF OBSERVATIONS IN DATA SET SAS | GENERAL LINEAR MODELS PROCEDURE | | | 12287 | 38804 | | PR > F | 0.0001 SAS | GENERAL LINEAR MODELS PROCEDURE | OR VARIABLE: YPE I COMPARI RROR RAIE | DF=21 MSE=20.6619 | 4 7.24379 7. | E NOT SIGNIFI | MEAN | 68.369 | 66.034 | 63,498 | 60.390 | 59.825 | 58,349 | 0.000 SAS | MAXTMITM |
| ผ | ERAL LINEAR | CLASS LEVEL | S LEVELS | 7 | OF OBSERVATIONS. | ERAL LINEAR | | MEAN SQUARE | 2300,54512287 | 20,66188804 | | F VALUE | 111.34 | ERAL LINEAR | RANGE TEST FY NITROLS THE TRIMENIMISE E | ALPHA=0.05 DP | 3,01172 | ME LETTER AR | GRÒUPING | ∢. | < < < | | יטט | ၁ ပ (| טיַט | Ω Ω | MTNTMIM |
| | GEN | | CLASS | TRI | NUMBER | GEN | | SUM OF SQUARES | 13803.27073720 | 433,89964876 | 14237.17038595 | TYPE I SS | 13803.27073720 | Nab | DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESPONSE NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE, NOT THE EXPERIMENTWISE ERROR RATE | AL | NUMBER OF MEANS CRITICAL RANGE 6.6759 | MEANS WITH THE SAWE LEITER ARE NOT SIGNIFICAVILY DIFFERENT | DUNCAN GR | | en e | n m a | n m | na ma | , | | CTANDARA |
| | | | | | | | ESPONSE | ក្ន | v | 21 | 27 | Ŋ | φ | | | | NUMBE | | | | desert de | 4 | hatch | | | | MFAN |
| | | | | | rd | | DEPENDENT VARIABLE: RESPONSE | SOURCE | MODEL | ERROR | CORRECTED TOTAL | SOURCE | TRT 1 | | | | | | | | Course house to | Rainban Front | of a whole of | | | 1 | WADT ABI E |
| 216. | 217. | 220. | 221. | 223. | 225. 227. 228. | 229. 230. | 231. | 234. | 235. | 237. | 239. 240. 241. | 242. | 244. 245. 246. | 247. | 249. 250. 251. 252. | 253. | 255. 256. 257. | 258. | 260. | 262. | 265. | 267. | 268. 269. | 270. | 272. | 2/4. 275. 276. | 277. 277. |

| | 9:44 WEDNESDAY, NOVEMBER 20, 1985 1 | | | | | 9:44 WEDNESDAY, NOVEMBER 20, 1985 2 | | | R-SQUARE C.V. | 0.997711 3.1019 | RESPONSE MEAN | 61,52642561 | F VALUE PR > F | 523.10 0.0001 NOVEMBER 20, 1985 3 | - | | | | | | | | | | | 9:44 WEDNESDAY, NOVEMBER 20, 1985 4 |
|---|-------------------------------------|---------------------------------|-------------------------|--------|--------------|-------------------------------------|---------------------------------|------------------------------|----------------|-----------------|---------------|-----------------|----------------|--|---------------------------------|--|-----------------|---|---|-----------------|-------------|---------------|------------|--------|--------|--|
| a velicla velica () C | 9:44 WEDNESDAY, | | | | | 9:44 WEDNESDAY, | | | PR > F | 0.0001 | ROOT MSE | 1,90848589 | TYPE III SS | 9526.51157352 523.10 9:44 WEDNESDAY, NOVEMBER 20, | | r rate, | | 6 4.97756 | RENT. | | | | | | | 9:44 WEDNESDAY, |
| | | URE | | | Ēų ELI | SET = 12 | URE | | F VALUE | 523.10 | | | | ហ | JRE | VESPONSE SONWISE ERRO | 32 | 5 4,95925 | ZANTLY DIFFE | N TRT | 2 D | 2 0 . | 7 BB | 2 A | 2 E | K 64 |
| | | GENERAL LINEAR MODELS PROCEDURE | CLASS LEVEL INFORMATION | VALUES | ABCDI | S IN DATA SI | GENERAL LINEAR MODELS PROCEDURE | | | 470 | 841 | | PR > F | 0.0001 | GENERAL LINEAR MODELS PROCEDURE | VARIABLE: F E I COMPARIS OR RATE | MSE=3.64232 | 3 4.91923 | NOT SIGNIFIC | MEAN | 81.837 | 79.442 | 75.294 | 68.445 | 64.140 | 0000 |
| i c | SAS | L LINEAR MO | LASS LEVEL | LEVELS | ٥ | NUMBER OF OBSERVATIONS IN DATA | L LINEAR MO | | MEAN SQUARE | 1905,30231470 | 3,64231841 | | F VALUE | 523.10 SAS | L LINEAR MOI | SE TEST FOR SIS THE TYPE SNIWISE ERR | ALPHA=0.05 DF=6 | 2 36 4.84023 | LETTER ARE N | SVI | ≪ « | 4 | | O C | υ | D SAS |
| | | GENERA | ד | CLASS | TRI | NUMBER OF (| GENERA | | SUM OF SQUARES | 9526.51157352 | 21.85391044 | 9548,36548397 | TYPE I SS | 9526.51157352 | GENERAL | DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESPONSE NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONMISE ERROR RATE, NOT THE EXPERIMENTWISE ERROR RATE | ALPHA= | NUMBER OF MEANS 2 CRITICAL RANGE 4.66996 | MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT. | DUNCAN GROUPING | | m a | a ma | | | e de terretario e e e e e de Mario e e e e e e e e e e e e e e e e e e e |
| | | | | | | | | RESPONSE | Ç | ហ | 9 | 11 | DF | īÙ | | | | | | | herbicide | 4. | | | | The semestrate yes a graduate |
| fetch 2013 clr 1 w216/1 16 : INVALID. 1216/1 | | | | | | ч | | DEPENDENT VARIABLE: RESPONSE | SOURCE | MODEL | ERROR | CORRECTED TOTAL | SOURCE | TRT | | | | | | | Command her | Rainbow trout | % survival | | | 1 |
| fetal ? 1 w2 w216 : ? 1 21 | 217. | 218. | 220. | 222. | 224. 224. | 226. 227. 228. | 230. | 231. | 233. | 236. | 238. | 240. | 242. | 245. 246. | 248. | 250. 251. 252. | 254. | 256. 257. | 258. 259. | 261. | 263. | 265. | 267. | 269. | | 273. |

650

10.

| PROCEDURE |
|-----------|
| MODELS |
| LINEAR |
| GENERAL |

CLASS LEVEL INFORMATION

CLASS LEVELS VALUES

TRT 5 ABCDE

NUMBER OF OBSERVATIONS IN DATA SET = 10 SAS

SAS

9:55 WEDNESDAY, NOVEMBER 20, 1985

GENERAL LINEAR MODELS PROCEDURE

| DEPENDENT VARIABLE: RESPONS | RESPONSE | | | | | | | |
|-----------------------------|----------|----------------|-------------|-------------|---------|---|----------------------|------------------|
| SOURCE | Д. | SUM OF SQUARES | MEAN SQUARE | UARE | F VALUE | PR > F | R-SQUARE | C.V. |
| MODEL | 4 | 10.4000000 | 2.60000000 | 0000 | 5.20 | 0.0499 | 0.806202 | 1.8085 |
| ERROR | ιΩ | 2,50000000 | 0.5000000 | 0000 | | ROOT MSE | RESI | RESPONSE MEAN |
| CORRECTED TOTAL | Ø | 12,9000000 | | | | 0.70710678 | Ř | 39.10000000 |
| SOURCE | DF | TYPE I SS | F VALUE | PR > F | DF | TYPE III SS | F VALUE | PR > F |
| IRT 1 | 4 | 10,4000000 | 5.20 SAS | 0.0499 S | 4 | 10.4000000 5.20 0.0499 9:55 WEDNESDAY, NOVEMBER 20, 1985 3 | 5.20 NOVEMBER 20, | 0.0499 1985 3 |

GENERAL LINEAR MODELS PROCEDURE

2216. 2217. 2218. 2219. 22200. 22200. 22200. 22200. 22200. 22200. 22200. 22200. 2

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESPONSE NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE, NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=5 MSE=0.5

NUMBER OF MEANS 2 3 4 5 CRITICAL RANGE 1.8196 1.87547 1.89503 1.90553

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

MEAN

DUNCAN GROUPING

| Command herbicida Rainbow Frout Mean total length | 당성 | icide 19th (mm) | | | 44444 | 40.5000 39.5000 | « A O O | | | |
|---|----|--------------------|------------------------|------|---------|--------------------|----------------------|------------------------|-----------------------------------|--------|
| | | - | - | ń | ooo | 37.5000 SAS | E E | 9:55 WEDNESDAY | 9:55 WEDNESDAY, NOVEMBER 20, 1985 | 1985 4 |
| VARIABLE | z | MEAN | STANDARD DEVIATION | _ | MINIMUM | MAXIMUM VALUE | STD ERROR OF MEAN | SUM | VARI ANCE | C.V. |
| RESPONSE 2 | 2 | 2 40.50000000 | 0.70710678 40.00000000 | 40.0 | | 1K1=A | 0.50000000 | 0.50000000 81.00000000 | 0.50000000 | 1.746 |

0

| ILS PROCEDURE | FORMATION | VALUES | ABCDE |
|---------------------------------|-------------------------|-------------|-------|
| GENERAL LINEAR MODELS PROCEDURE | CLASS LEVEL INFORMATION | LEVELS VALU | 5 AB |
| GENERAL I | CLASS | CLASS | TRT |

. The second sec

| NUMBER OF OBSERVATIONS IN DATA SET = 10 9:57 WEDNESDAY, NOVEMBER 20, 1985 2 | GENERAL LINEAR MODELS PROCEDURE | | MEAN SQUARE F VALUE PR > F R-SQUARE C.V. | 4973.85000000 5.71 0.0417 0.820331 5.4277 | 871,50000000 ROOT MSE RESPONSE MEAN | 29,52117884 543,9000000 | JE PR > F DF TYPE III SS F VALUE PR > F | 5.71 0.0417 4 19895.40000000 5.71 0.0417 SAS 9:57 WEDNESDAY, NOVEMBER 20, 1985 3 | |
|---|---------------------------------|------------------------------|--|---|-------------------------------------|-------------------------|---|---|---|
| NUMBER OF OBSE | GENERAL LI | | SUM OF SQUARES MI | 19895,40000000 497 | 4357,50000000 87 | 24252,90000000 | TYPE I SS F VALUE | 19895,4000000 5. | |
| | | BLE: RESPONSE | පිට | 4 | ហ | 60 | J. | 4 | |
| - | | DEPENDENT VARIABLE: RESPONSE | SOURCE | MODEL | FRECR | CORRECTED TOTAL | SOURCE | IRI | • |

216. 2217. 2219. 2219. 2220. 2221. 2221. 2221. 2222. 2223. 2223. 2233. 2334. 2336. 2337. 2336. 2337. 2337. 2337. 2337. 2337. 2337. 2337. 2347. 2447. 2447. 2447. 2447. 2447. 2447. 2447.

GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESPONSE NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONMISE ERROR RATE, NOT THE EXPERIMENTMISE ERROR RATE

ALPHA=0.05 DF=5 MSE=871.5

| 79,5543 | DIFFERENT. |
|-----------------------------------|------------------------------------|
| 4 79.1161 | NIFICANTLY |
| 3 78.2996 | R ARE NOT SIGNIFICANILY DIFFERENT. |
| 2 75.9669 | LETTER A |
| NUMBER OF MEANS CRITICAL RANGE | MEANS WITH THE SAME LETTER |

| | w.** | | | | 9:57 WEDNESDAY, | SUM |
|----------|--------|-------------------|--------------|----------------------|-----------------|----------------------|
| IRT | Æ | æ | Ω | ပ | ជា | SEOR SAN |
| z | 8 | 64 | .2 | 7 | 6 | STD ERROR OF MEAN |
| MEAN | 600.00 | 557.50 | 551.50 | 547.50 | 463.00 | MAXIMUM VALUE |
| | | | | | SAS | |
| GROUPING | A. | 4 4 · | 4 A | 4 4 | æ | MINIMUM |
| DUNCAN | | | | | | STANDARD |
| | | 0,010 | | Avg. wet weight (mg) | ', | MEAN |
| | | A her | Took a | t werg | • | Z |
| | | Command herbicide | Rainbow Hout | Avg.we | | VARIABLE |

251. 252. 253. 254. 255. 255. 256. 266. 266.

C. <

VARIANCE

NOVEMBER 20, 1985

5,893

25,0000000 1200,0000000 1250,0000000

35,35533906 575,00000000 625,0000000

2 600,00000000

RESPONSE

TRI=A -

652

12.