

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

MEMORANDUM

MAY 13 1992

OFFICE OF PESTICIDES AND TOXIC SUBSTANCES

SUBJECT:

Dietary Exposure Analysis and De Minimis Risk

Assessment for Temporary Tolerance and Experimental

Use Permit for Tebuconazole on Peanuts

(PP#9F3724/2H5628)

FROM:

Stephen A. Schaible Aschaible

Dietary Exposure Section

SACB/HED (H7509C)

TO:

Susan T. Lewis/Julie Fairfax, PM 21

Fungicide-Herbicide Branch

Registration Division (H7505C)

THROUGH:

James P. Kariya, Chief Mariya Dietary Exposure Section

Health Effects Division

Action Requested

The Dietary Exposure Section (DES) was requested by Registration Division to perform a dietary exposure analysis and carcinogenic risk assessment for the chemical tebuconazole, to see whether the Agency's de minimis value for negligible risk would be exceeded by the carcinogenic risk posed by this Experimental Use Permit (EUP) and temporary tolerance on peanuts. Tebuconazole is a new chemical with no registered uses.

Since it is only recently submitted data that suggest this chemical may be a carcinogen, it has not yet gone to the HED Carcinogenicity Peer Review Committee, nor has its possible upper bound carcinogenic potency factor (Q1*) been determined. Without the appropriate Q*, a carcinogenic risk analysis cannot be performed by DES. In order to respond to the expedited review request by RD, an alternate analysis was performed instead, in which DES assumed the de minimis value of 10⁻⁶ for the carcinogenic risk and calculated the Q* that would be necessary to arrive at that de minimis value given the exposure value contributed by the EUP on peanuts.

Toxicological Endpoints

The Dietary Risk Evaluation System (DRES) chronic exposure analysis used a Reference Dose (RfD) of 0.01 mg/kg body weight/day, based on a no observed effect level (NOEL) of 1 mg/kg bwt/day and an uncertainty factor of 100. The RfD is based on a one year feeding

study in dogs which demonstrated as effects lenticular and corneal opacity and hepatic toxicity. This RfD has been approved by the HED RfD Peer Review Committee (3/5/91).

Toxicology Branch II recently received 6(a)(2) data concerning an oncogenicity study in mice which showed significant incidences of carcinomas and adenomas at the high dose in males and females (personal communication, A. Protzel, 5/7/92), but at this time tebuconazole is not considered a carcinogen.

Residue Information

The food use evaluated in this analysis was the temporary tolerance and EUP for tebuconazole on peanuts. The commodities listings in DRES which relate to peanuts are "peanuts-whole" and "peanuts- oil". For the purpose of this temporary tolerance request, tolerances reflecting secondary residues in animal commodities from the use of peanuts as a feed item are not necessary. Since the draft label dated 10/1/91 included a restriction against feeding treated peanut hay/vines to livestock (G. Otakie, 3/18/92), secondary residues are not expected in animal commodities. Tebuconazole is a new chemical and has no registered uses. A summary of the residue information used in this analysis is attached as Table 1.

Exposure Analysis

The DRES chronic exposure analysis used tolerance level residues and 100% crop treated to estimate the Theoretical Maximum Residue Contribution (TMRC) for the overall U.S. population and 22 population subgroups. A list of the TMRCs and their representations as percentages of the RfD are attached as Table 2.

The TMRC for the overall population from the proposed use on peanuts is 0.000010 mg/kg bwt/day, which represents approximately 0.1% of the RfD. The subgroup most highly exposed, children aged one through six years old, has a TMRC of 0.000027 mg/kg bwt/day, or 0.27% of the RfD. None of the subgroups has a TMRC that exceeds even one percent of the RfD, so it appears that the chronic risk from this EUP is minimal.

De Minimis Risk Assessment

As was mentioned before, no upper bound potency factor (Q*) has been determined for tebuconazole. Upper bound carcinogenic risk is calculated using the formula

Upper bound carcinogenic risk = Exposure (TMRC) x Q*

If the Q* is unknown, it is not possible to calculate the upper bound carcinogenic risk. However, by assuming the Agency's de minimis value of 10⁻⁶ as the upper bound carcinogenic risk value in the equation, we can arrive at a "reference Q*", the Q* which would be necessary to contribute a cancer risk of 10⁻⁶, given the known

					1
HISPANICS NON-HISPANIC WHITES NON-HISPANIC BLACKS NON-HISPANIC OTHERS	NORTHEAST REGION NORTH CENTRAL REGION SOUTHERN REGION WESTERN REGION	U.S. POPULATION - SPRING SEASON U.S. POPULATION - SUMMER SEASON U.S. POPULATION - FALL SEASON U.S. POPULATION - WINTER SEASON	U.S. POPULATION - 48 STATES	POPULATION SUBGROUP	999ZZZ(TEBUCONAZOLE) CBSWell #999ZZZ CAS No. A.I. CODE: CFR No. 180.
		NOS NOS NOS			NOEL= LEL=
				.	1.0000 mg/kg 0.00 ppm 0.0000 mg/kg 0.000 ppm
0.000000 0.000000 0.000000	0.000000 0.000000 0.000000	0.000000 0.000000 0.000000	0.000000	CURRENT TMRC*	
0.000006 0.000010 0.000006 0.000008	0.000010 0.000010 0.000008 0.000011	0.000010 0.000010 0.000009 0.000009	0.000010	TOTAL TMRC (MG/KG BODY WEIGHT/DAY) CURRENT TMRC* NEW TMRC**	
0.057530 0.104470 0.058870 0.080160	0.101130 0.100000 0.078640 0.111720	0.098220 0.103360 0.092520 0.088720	0.095710	Y) NEW TMRC AS PERCENT OF RFD	PADI UF>100 OPP RfD= 0.010000 EPA RfD= 0.000000
0.057530 0.104470 0.058870 0.080160	0.101130 0.100000 0.078640 0.111720	0.098220 0.103360 0.092520 0.088720	0.095710	DIFFERENCE AS PERCENT OF RFD	test run fo tebuconazol
				EFFECT OF A	test run for temp tol for tebuconazole on peanuts
				EFFECT OF ANTICIPATED RESIDUES ARC %RFD	

NURSING INFANTS (< 1 YEAR OLD)
NON-NURSING INFANTS (< 1 YEAR OLD)
FEMALES (13+ YEARS, PREGNANT)
FEMALES 13+ YEARS, NURSING
CHILDREN (1-6 YEARS OLD)
CHILDREN (7-12 YEARS OLD)
MALES (13-19 YEARS OLD)
MALES (13-19 YEARS OLD, NOT PREG. OR NURSING)
MALES (20 YEARS AND OLDER, NOT PREG. OR NURS)
FEMALES (20 YEARS AND OLDER, NOT PREG. OR NURS)

0.00002 0.00003 0.00007 0.000010 0.000027 0.000020 0.000010

0.101440 0.268600 0.200080 0.102390

0.019130 0.034070 0.072230 0.101440 0.268600 0.200080 0.102390

0.019130 0.034070 0.072230

0.000007 0.000006 0.000005

0.074680 0.064050 0.049860

0.074680 0.064050 0.049860

^{*}Current TMRC does not include new or pending tolerances.
**New TMRC includes new, pending, and published tolerances.

DATE: 05/07/92

PAGE:

STATUS

NOEL= LEL= STUDY TYPE 1.0000 mg/kg 0.00 ppm 0.0000 mg/kg 0.000 ppm **EFFECTS** PADI OPP RfD= 0.010000 EPA RfD= 0.000000 REFERENCE DOSES
DI UF -->100 DATA GAPS/COMMENTS
test run for temp tol for
tebuconazole on peanuts

CAS No. A.I. CODE: CFR No. 180. Caswell #999777 ONCO:

999ZZZ(TEBUCONAZOLE)

CHEMICAL INFORMATION

LISTING OF EXPOSURE BY RAC FOR: U.S. POPULATION - 48 STATES

15006AA 270070A GRAND TOTAL TMRC: CROP GROUP TOTALS FOR LEGUME VEGETABLES: CO 75 00 00 00 PEANUTS-WHOLE FOOD NAME 0.009571 GRAND TOTAL % OF THE RfD: 0.10000 NEW PENDING PUBLISHED 0.0957 |--EXISTING TOLERANCES--| TMRC (UG/KG/DAY) -NEW & PENDING TOLERANCES-TMRC (UG/KG/DAY) 0.006958 0.009571 XRFD 0.0695 0.0957

× exposure = upper bound concinegence rish

POPULATION TOTAL TMRC

0.009571

POPULATION TOTAL % OF THE RFD

0.0957

0.000000

0.0000

0.009571

POPULATION SUBGROUP TOTALS

Q *

reference Q, * = supper bound concinoquie non 20-5 exposure (TMRC)

= 0.1 (mg/kg/day)-1

exposure from the EUP on peanuts. In other words, we reshuffled the equation to read as follows

Reference $Q^* = Carcinogenic risk (10^{-6}) / Exposure (TMRC)$

The Q* arrived at using this analysis is the highest value that could be determined for the upper bound potency factor and still have the cancer risk posed by this EUP not exceed the de minimis value. Using this formula, the "reference Q*" for this action is approximately 0.1 (mg/kg/day)⁻¹. If the Q* that Toxicology Branch and the statisticians in SACB determine is less than this value, the resulting upper bound carcinogenic risk will be less than the de minimis value.

There are several assumptions that possibly make this "reference O*" a more sensitive value than it actually should be. Assumptions made in calculating the exposure value, such as tolerance level residues and 100 percent of crop treated, in all likelihood overestimate the exposure, especially the percent of crop treated assumption, since the acreage that this chemical will be applied to through this EUP probably only represents a small percent of the total acreage that peanuts are grown on. Also, this risk assessment was performed assuming exposure over 70 years while the duration of the temporary tolerance and experimental use permit for tebuconazole on peanuts would be only one growing season. If there were a way to incorporate refinements to these sources of possible overestimation, one would expect that the reference Q* would be a higher value, thus allowing the determined Q* to be more potent than 0.1 (mg/kg/day)⁻¹ and still not contribute to an upper bound carcinogenic risk of more than 10⁻⁶.

Attachments

cc: DES, Tox 1, CBTS, C. Frick, Caswell # 463P

TABLE

6

F000 C00E	999ZZZ(TE Casw CAS I A.I. CFR I	
FOOD NAME	799ZZZ(TEBUCONAZOLE) Caswell #999ZZZ CAS No. A.I. CODE: CFR No. 180.	CHEMICAL
m		
	NOEL=	S
	1.0000 mg/kg 0.00 ppm 0.0000 mg/kg 0.000 ppm	CHEMICA
PETITION NUMBER		CHEMICAL INFORMATION FOR CASWELL NUMBER 999ZZZ
N N N N		FOR CASWELL EFFECTS
P C		NUMBER
TOLERANCE (PPM) PENDING PUBLISHED	PAD1 UF>100 OPP RfD= 0.010000 EPA RfD= 0.000000	REFERENCE DOSES
	test run for temp tol fo tebuconazole on peanuts	DATA GAPS/COMMENTS
	7	PAGE:
		STATUS

15006AA PEANUTS-WHOLE 27007OA PEANUTS-OIL

9G3817 2H5628

0.100000 0.500000