

4. 3-1-93  
Reviewed by: Stanley B. Gross, Ph.D.  
Section 2, Toxicology Branch 1 (H7509C)  
Secondary Reviewer: Melba S. Morrow, DVM  
Section 2, Toxicology Branch 1, (H7509C)

*Stanley B. Gross*  
*2/23/93*  
*msm 3/1/93*

#### DATA EVALUATION REPORT

STUDY TYPE: Skin Irritation Study in Rabbits.

TOX. CHEM. NO.: New Chemical 129032

ACCESSION NUMBER: D179381

MRID NO: 421783-05; Amended MRID #41827-14.

TEST MATERIAL: (2-[1-methyl-2-(4-phenoxyphenoxy) ethoxy]  
pyridine.

SYNONYMS: Sumilarv; S-31183.

STUDY NUMBER(S): GLN 81-5. (Carried out by the guideline of the  
Ministry of Agriculture, Forestry and Fisheries, Japan)

SPONSOR: Sumitomo Chemical Co., Osaka, Japan.

TESTING FACILITY: Sumitomo Chemical Company, Limited. 5-33,  
Kitahama 4-Chome, Chuo-Ku, Osaka, Japan.

TITLE OF REPORT: Sumilarv -- Acute oral toxicity of S - 31183 in  
Rats. (Original report written in Japanese,  
translated by Takachi Suzuki, May 14, 1987.)

AUTHOR(S): Takashi Suzuki

REPORT ISSUED: Number NNT-70-0022, January 27, 1987

GLP Review: Masanori Takatsuka, Jan. 21, 1987.

#### CONCLUSIONS:

Five hundred mg of Sumilarv was applied to abraded and  
intact skin of the back of 6 rabbits for 4 hours. No reactions  
(erythema or edema) were observed over the next 48 hours.

CLASSIFICATION: Non-Irritant. ~~Core Guideline~~. Tox CAT IV  
ACCEPTABLE

#### A. MATERIALS:

1. Test compound: S-31183, Technical. Description: White  
solid, manufactured by Sumitomo Chemical Co. Batch # PTG-86011.  
Purity: 97.2 %, Impurities are noted in composition statement.

2. Test animals: Species: Rabbits. Strain: New Zealand White. Age: six weeks. Weight: 2.26 to 2.78 Kg. Source: Nihon Dobutsu, Co, Osaka.

Housing: Rabbits were housed in individual cages under standard laboratory conditions and offered food and water ad libitum.

B. STUDY DESIGN:

The hair on the backs of three males and three female rabbits was clipped on both sides of the median line (ca. 15 x 15 cm). One sided of the clipped skin area was abraded using a 18 G needle. Five hundred mg of the Sumilarv dry powder was spread on a 2.5 x 2.5 cm lint patch and moistened with a small amount of water. The lint patch with the Sumilarv was then applied to the abraded and intact skin areas and covered with an occlusive tape held in place for 4 hours. At the end of the four hour period, the patch was removed and the skin wiped with cotton moistened with water. The reaction of the skin was scored at 4.5, 24, 48 and 72 hrs. from the application time using the method of Draize<sup>1</sup>.

C. RESULTS:

The resulting scores of the skin reactions were summarized in Table 8, page 14 (attached). There were no signs of irritation (erythema or edema) in any of the animals at any of the evaluation times.

smlrvskn. 2/23/93

---

<sup>1</sup> Draize, J. H., et al (1944). J. Pharmacol. Exp. Therap., 82: 377-390.

Pyriproxyfen

RIN 4445-96

P.C. 129032

Page 3 is not included in this copy.

Pages \_\_\_\_\_ through \_\_\_\_\_ are not included.

---

The material not included contains the following type of information:

- ☐ Identity of product inert ingredients.
- ☐ Identity of product impurities.
- ☐ Description of the product manufacturing process.
- ☐ Description of quality control procedures.
- ☐ Identity of the source of product ingredients.
- ☐ Sales or other commercial/financial information.
- ☐ A draft product label.
- ☐ The product confidential statement of formula.
- ☐ Information about a pending registration action.
- ☒ FIFRA registration data.
- ☐ The document is a duplicate of page(s) \_\_\_\_\_.
- ☐ The document is not responsive to the request.

---

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.

---