


8-6-93

MRID No. 427702-30


DATA EVALUATION RECORD

1. **CHEMICAL:** Pirate® (AC 303,630).
Shaughnessey No. 129093.
2. **TEST MATERIAL:** AC 303,630 technical; CAS No. 122453-73-0;
Batch No. AC 7504-59A; 94.5% purity; a tan powder.
3. **STUDY TYPE:** 71-2. Avian Dietary LC₅₀ Test. Species
Tested: Bobwhite quail (*Colinus virginianus*).
4. **CITATION:** Gagne, J.A. and J.P. Sullivan. 1993. 8-Day
Acute Dietary LC₅₀ Test with AC 303,630 in the Northern
Bobwhite (*Colinus virginianus*). Laboratory Project ID No.
105-009-01. Performed by Bio-Life Associates, Ltd.,
Neillsville, WI. Submitted by American Cyanamid Company,
Princeton, NJ. EPA MRID No. 427702-30.
5. **REVIEWED BY:**



Mark A. Mossler, M.S.
Associate Scientist
KBN Engineering and
Applied Sciences, Inc.

Signature: 
Date: 7/1/93
6. **APPROVED BY:**

Michael Whitten, M.S.
Wildlife Toxicologist
KBN Engineering and
Applied Sciences, Inc.

Signature: 
Date: 7/1/93

Henry T. Craven, M.S.
Supervisor, EEB/EFED
USEPA

Signature: 
Date: 
7. **CONCLUSIONS:** This study is scientifically sound and meets
the guideline requirements for an avian dietary LC₅₀
toxicity test. The LC₅₀ of the test material for bobwhite
quail was 132 ppm, which classifies AC 303,630 technical as
highly toxic to bobwhite quail. The NOEC was 10 ppm.
8. **RECOMMENDATIONS:** N/A.
9. **BACKGROUND:**
10. **DISCUSSION OF INDIVIDUAL TESTS:** N/A.

11. MATERIALS AND METHODS:

- A. **Test Animals:** Bobwhite quail (*Colinus virginianus*) were hatched from eggs obtained from a commercial supplier in Gravette, AR. All birds were phenotypically indistinguishable from wild birds and were acclimated to the testing conditions for 4 days. No deaths occurred during the acclimation period. The birds were 12 days of age at test initiation.
- B. **Test System:** The birds were housed in brooder pens in a thermostatically controlled room. The pens measured 91 x 24 x 28 cm. During the test, the average minimum and maximum temperatures were 30 and 37°C, respectively, and the relative humidity ranged between 42 and 61%. A 16-hour fluorescent lighting photoperiod was used throughout the study.

For each treatment diet, test material was dissolved in acetone (covered and stirred) and added to Purina® Game Bird Startena. The diets were allowed to blend for 15 minutes and the acetone was allowed to evaporate from all diets before administration to the birds. Water and food were supplied *ad libitum*. Prior to and following the 5-day exposure period, all birds were placed on regular feed.

- C. **Dosage:** Eight-day dietary LC₅₀ test. Dosage levels selected for the study were 10, 20, 40, 80, 160, and 320 ppm. The amount of test material added to the diets was not adjusted for percent purity. A vehicle control diet was also prepared.
- D. **Design:** Ten chicks per treatment level and control were randomly assigned to pens. Observations were made twice daily for clinical signs indicative of test material effect. Inspections were made twice daily for mortalities and abundance of food and water.

The 17 birds that died during the test and five arbitrarily selected birds sacrificed from each of the control and 10, 20, 40, and 80 ppm treatment groups at the termination of the project were subjected to gross pathological examinations. The surviving three birds at the 160 ppm level were also inspected.

Individual bodyweights were measured at 0-hour on day 1 and on test days 5 and 8. Average feed consumption was determined daily by group for days 1-5 (the exposure period) and days 6-8 (the observation period).

Stability and homogeneity samples were collected from a trial diet. Stability samples were allowed to remain at room temperature for 5 and 10 days. Immediately after diet preparation for the definitive test, concentration verification samples were collected from the control and each treatment diet. The samples were frozen and sent to the sponsor for analysis using high performance liquid chromatography.

E. **Statistics:** The acute median lethal concentration (LC_{50}) and associated 95% confidence interval (C.I.) were calculated using the simplified method of Litchfield and Wilcoxon. The no-observed-effect concentration (NOEC) was based on Dunnett's test ($p \leq 0.05$) and clinical observations.

12. **REPORTED RESULTS:** No mortality was observed in the control or four lowest treatment groups during the study. Seventy and one-hundred percent mortality was observed in the 160 and 320 ppm groups, respectively (Table 1, attached). The 8-day dietary LC_{50} of AC 303,630 technical for bobwhite quail was determined to be 132 ppm (95% C.I. = 88-170 ppm). Based on this value, AC 303,630 technical would be classified as highly toxic to bobwhite quail.

No clinical signs of toxicity were noted for the control or 10 ppm birds. Various effects on the excreta of the birds were observed for the five highest treatment levels (Table 3, attached). Complete remission of these signs was achieved by the beginning of test day 5.

Mean percent change in bodyweight of the 80 and 160 ppm treatment birds was significantly reduced in comparison to the vehicle control birds for the exposure and entire test periods (Table 5, attached). Statistical analysis of the bodyweights of the highest concentration birds was not undertaken due to complete mortality by day 5.

Feed consumption appeared to be reduced at the two highest treatment levels on test day 1. All other values were equal or greater to the vehicle control values (Table 6, attached).

Gross changes were noted in all birds that died. The majority of these changes were noted in birds that died during the night, and therefore, the changes were believed to be due to postmortem autolysis (Table 7, attached).

The NOEC was determined to be 10 ppm, based on the lack of clinical signs of toxicity at this treatment level.

13. **STUDY AUTHOR'S CONCLUSIONS/QUALITY ASSURANCE MEASURES:**
No conclusions other than those stated above were presented by the authors.

Quality Assurance and Good Laboratory Practice (GLP) compliance statements were included in the report, indicating that the study was conducted in accordance with GLPs as set forth in 40 CFR Part 160. A Quality Assurance statement was also included in the analytical appendix.

14. **REVIEWER'S DISCUSSION AND INTERPRETATION OF STUDY RESULTS:**

- A. **Test Procedure:** The test procedures were in accordance with Subdivision E, ASTM, and SEP guidelines with the following exception:

The pen dimensions (91 x 24 cm = 2,184 cm²) were smaller than recommended (35 x 100 cm = 3,500 cm²).

- B. **Statistical Analysis:** The reviewer used EPA's Toxanal program to determine the LC₅₀ value and 95% confidence interval (see attached printout). The value was similar to that of the authors.

- C. **Discussion/Results:** Results of chemical analyses indicated that the material was homogeneously mixed, stable, and present at the desired levels. The results are summarized in Appendix II, Tables I, IIB, IIIA, and IVA (attached). Although the stability and homogeneity data were derived from trial diets, the reviewer believes that the definitive diets would have demonstrated these same qualities.

After review of the mortality, weight, and feed consumption data, the reviewer noted that there were food consumption reductions at the 40 and 80 ppm levels for day 1. There also appeared to be reductions at the 80 and 160 ppm levels for days 3, 4, and 5. However, since these effects were all noted at levels greater than those at which clinical signs of toxicity were noted, the NOEC value of 10 ppm is considered to be correct.

This study is scientifically sound and meets the guideline requirements for an avian dietary LC₅₀ toxicity test. The LC₅₀ of the test material for bobwhite quail was 132 ppm, which classifies AC 303,630 technical as highly toxic to bobwhite quail. The NOEC was 10 ppm, based on the lack of clinical signs of toxicity.

D. Adequacy of the Study:

(1) Classification: Core.

(2) Rationale: N/A.

(3) Repairability: N/A.

15. COMPLETION OF ONE-LINER: Yes, 6-28-93.

Page _____ is not included in this copy.

Pages ~~16~~ through 20 are not included.

The material not included contains the following type of information:

- _____ Identity of product inert ingredients.
- _____ Identity of product impurities.
- _____ Description of the product manufacturing process.
- _____ Description of quality control procedures.
- _____ Identity of the source of product ingredients.
- _____ Sales or other commercial/financial information.
- _____ A draft product label.
- _____ The product confidential statement of formula.
- _____ Information about a pending registration action.
- ☒ FIFRA registration data.
- _____ The document is a duplicate of page(s) _____.
- _____ The document is not responsive to the request.

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.

MOSSLER AC 303630 COLINUS VIRGINIANUS 6-28-93

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
320	10	10	100	9.765625E-02
160	10	7	70	17.1875
80	10	0	0	9.765625E-02
40	10	0	0	9.765625E-02
20	10	0	0	9.765625E-02
10	10	0	0	9.765625E-02

THE BINOMIAL TEST SHOWS THAT 80 AND 320 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS, BECAUSE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 136.5848

WHEN THERE ARE LESS THAN TWO CONCENTRATIONS AT WHICH THE PERCENT DEAD IS BETWEEN 0 AND 100, NEITHER THE MOVING AVERAGE NOR THE PROBIT METHOD CAN GIVE ANY STATISTICALLY SOUND RESULTS.
