MRID No. 429186-27

DATA EVALUATION RECORD

- 1. <u>CHEMICAL</u>: Fipronil and derivatives (M&B 46030). Shaughnessey No. 129121.
- 2. <u>TEST MATERIAL</u>: M&B 46030; Lot No. 78 GC90; 96.7% active ingredient; a white powder.
- 3. <u>STUDY TYPE</u>: 72-4. Freshwater Fish Early Life-Stage Test. Species Tested: Rainbow Trout (Oncorhynchus mykiss).
- 4. <u>CITATION</u>: Machado, M.W. 1992. (M&B 46030) The Toxicity to Rainbow Trout (Oncorhynchus mykiss) During an Early Life-Stage Exposure. SLI Report No. 92-1-4084. Performed by Springborn Laboratories, Inc., Wareham, MA. Submitted by Rhone-Poulenc Ag Company, Research Triangle Park, NC. EPA MRID No. 429186-27.
- 5. REVIEWED BY:

Rosemary Graham Mora, M.S. Associate Scientist KBN Engineering and Applied Sciences, Inc.

6. APPROVED BY:

Pim Kosalwat, Ph.D. Senior Scientist KBN Engineering and Applied Sciences, Inc.

James J. Goodyear, Ph.D. Project Officer, EEB/EFED USEPA

Signature:

Date: 2

signature: P. Kosalu ja t

Date: 2/4/94

signature:

Date:

7. <u>CONCLUSIONS</u>: This study is scientifically sound and meets the guideline requirements for a fish early life-stage test. Based on the effects of M&B 46030 on larval length, the MATC for rainbow trout was >6.6 and <15 μ g ai/l (geometric mean MATC = 9.9 μ g ai/l).

- 8. RECOMMENDATIONS: N/A.
- 9. BACKGROUND:
- 10. DISCUSSION OF INDIVIDUAL TESTS: N/A.

11. MATERIALS AND METHODS:

- Test Animals: Unfertilized rainbow trout (Oncorhynchus mykiss) eggs and sperm were obtained (separately packaged at 9°C; warmed to 11°C upon receipt) from Mount Lassen Trout Farm, Red Bluff, CA. The eggs were fertilized in a stainless steel bowl by adding the eggs to half of the sperm. The bowl was swirled to thoroughly mix the contents. The remaining sperm and a small amount of control water (11°C) were added to the bowl and swirled again. The contents were allowed to stand undisturbed for 2 minutes to complete fertilization. The excess sperm was rinsed from the eggs with control water. The eggs were allowed to harden for 1 hour. During the period of egg distribution, the temperature was allowed to rise to
- B. Test System: The test system was an intermittent-flow proportional diluter with a 50% dilution factor. The diluter delivered approximately 2 l of solution per cycle to each aquarium at an average rate of approximately 6.6 volume replacements per day.

Each glass test aquarium (39 x 20 x 25 cm) was equipped with a 19.5-cm high side drain that maintained a constant solution volume of 15 l. Embryo incubation cups were glass jars (5 cm O.D., 8 cm high) with 2000 micron-mesh Nitex® screen bottoms. A rocker arm apparatus was used to gently oscillate incubation cups in the test chambers.

The aquaria were impartially positioned in a water bath designed to maintain the solution temperature at 12 ±1°C. Sixteen hours of light at an intensity of 30-50 footcandles were provided each day. Sudden transitions from light to dark and dark to light were avoided.

The dilution water was well water which was pumped into an epoxy-coated concrete reservoir where it was supplemented with untreated Town of Wareham well water and aerated. The well water, characterized weekly, had total hardness and alkalinity ranges of 24-32 and 18-25 mg/l as $CaCO_3$, respectively, a pH range of 6.8-7.4, and a specific conductivity range of $100-150~\mu mhos/cm$.

A stock solution (15.4 mg ai/ml) was prepared by dissolving 0.796 g (0.770 g ai) of test material in acetone to a final volume of 50 ml.

- C. <u>Dosage</u>: Ninety-day, flow-through test. Based on results of preliminary testing, nominal test concentrations selected were 6.2, 12, 25, 50, and 100 μg ai/l. A dilution water control and a solvent control (6.5 μl acetone/l) were also included. The concentration of acetone in the solvent control was equivalent to the highest solvent concentration in any exposure concentration.
- D. <u>Design</u>: Fifty embryos were impartially distributed to each of 28 incubation cups. Two cups were suspended in each duplicate test chamber per treatment.
 - Embryo viability was determined on day 18, when all the eggs exhibited well-pronounced embryonic development. Due to the extreme sensitivity of the embryos to physical trauma, dead embryos were not removed from the incubation cups until test day 16. On day 18, ten live embryos were impartially selected from each of the two incubation cups (20 per replicate aquarium) and placed into one incubation cup. The remaining embryos were placed in the second cup of each replicate. On test day 30 (considered hatch day), the surviving larvae from the first incubation cup of each replicate were released into their respective aquarium. Unhatched embryos remained in the incubation cups until day 31. The surviving larvae from the second cup of each replicate were counted and discarded. Survival of organisms at hatch (number of live larvae/number of viable embryos) was determined on day 31 when hatching was considered complete.

When larvae reached swim-up stage and began actively feeding (day 13 post-hatch), they were fed live Artemia nauplii two to three times daily. The aquaria were brushed and siphoned when necessary.

Behavior and appearance of larvae were observed and recorded daily. Larval survival was determined twice weekly. At test termination (60 days post-hatch), the larvae were counted, and individually weighed and measured.

Dissolved oxygen concentration (DO), pH, and temperature for each replicate were measured daily. Temperature was continuously monitored in one replicate of the control. Conductivity, total hardness, and alkalinity were measured on day 0 and weekly thereafter in alternating replicates of the highest and lowest exposure groups and the control.

Water samples were collected from the midpoint of each aquarium on test day 0 and at least weekly thereafter for determination of M&B 46030 concentrations. Samples were analyzed using high performance liquid chromatography.

- E. Statistics: The percentage survival data were transformed (arcsine square-root percentage) before analysis. All statistical analyses were performed using the mean organism response in each replicate aquarium. All data were tested for assumptions of homogeneity of variance (Bartlett's test). A one-way, single-classification analysis of variance (ANOVA) was used to determine significant differences between the dilution water control and the solvent control. No significant differences were determined, therefore, the control data were pooled for comparison to the exposure concentrations. Williams' test was used to assess treatment effects. Treatment levels that caused significant survival effects were excluded from the analysis of growth data. All statistical conclusions were made at 95% confidence level, except for Bartlett's test which was done at 99% level.
- 12. REPORTED RESULTS: Mean measured concentrations were 2.6, 6.6, 15, 26, and 60 μ g ai/l which represent 42-60% of nominal concentrations (Tables 2 and 3, attached).

No significant difference in embryo viability or hatchability was observed between the exposure concentrations and the pooled control (Table 5, attached). Survival at 26 and 60 μg ai/l was significantly reduced when compared to that of the pooled control.

Larval length at 2.6 and 15 μ g ai/l was significantly reduced when compared to that of the pooled control. However, the reduction at both levels was considered biologically insignificant. "The absence of a concentration-response relationship, for organism length, at these lower treatment levels (i.e., 2.6-15 μ g ai/l) and no statistical difference between larval weight at the same exposure concentrations, corroborates the lack of biological significance between the growth of rainbow trout exposed to \leq 15 μ g ai/l M&B 46030 and the control solutions." Wet weight at concentration \leq 15 μ g ai/l was not significantly reduced when compared to pooled control data.

During the study, the test solutions had a pH range of 6.6-7.4, a mean DO of 9.6-10.0 mg/l, and a mean total hardness and alkalinity of 30-31 and 27-28 mg/l as $CaCO_3$,

respectively. Minimum/maximum temperature readings ranged from 10 to 19°C.

"Based on the statistically significant (p<0.05) reduction in larval survival following 60 days post-hatch exposure, the Lowest Observed Effect Concentration (LOEC) was determined to be 26 μ g ai/l and the No Observed Effect Concentration (NOEC) was determined to be 15 μ g ai/l. Based on these results, the Maximum Acceptable Toxicant Concentration (MATC) for M&B 46030 and rainbow trout embryos and larvae is estimated to be \geq 15 and \leq 26 μ g ai/l (geometric mean MATC = 20 μ g ai/l)."

Good laboratory practice (GLP) compliance and quality assurance statements were included in the report indicating that the study was performed in accordance with EPA GLP regulations (40 CFR Part 160), except in the case of stability, characterization, and verification of test substance identity.

14. REVIEWER'S DISCUSSION AND INTERPRETATION OF STUDY RESULTS:

A. <u>Test Procedure</u>: The test procedure is generally in accordance with the SEP and ASTM guidelines, except for the following deviations:

Raw water quality data were not included in the report as required.

Only two replicates per treatment level were used during the study; the SEP requires four replicates.

The total hardness (24-32 mg/l as $CaCO_3$) of the dilution water was slightly lower than the recommended hardness (40-48 mg/l as $CaCO_3$).

The light intensity employed in this study was 323-538 lux at the water surface. The SEP recommends the intensity of 400-800 lux.

The continuous test temperature ranged from 10 to 19°C; the SEP recommends 10 \pm 2°C during the exposure period.

Embryos were "impartially" distributed; the SEP requires random distribution of test organisms to the test vessels.

B. <u>Statistical Analysis</u>: Survival data were analyzed using Toxstat® (Version 3.3). The data failed the

assumptions of homogeneity of variance (Bartletts or Hartley tests) or normality (Chi-square or Shapiro Wilks tests), therefore a non-parametric test (Kruskal-Wallis) was used to evaluate treatment effects. The reviewer's analysis of survival data showed no significant difference in exposure concentrations when compared to the control (printout attached). However, upon visual examination of the data, it is obvious that there is a substantial reduction in survival (78 and 0% survival) at the two highest concentrations when compared to the control (98% survival).

Individual length and weight data were analyzed using a 2-way analysis of variance (ANOVA) coupled with Bonferroni's test and a contrast test (Systat®) for treatment comparisons (printout, attached). The results of Bonferroni's test for weight and length data showed a significant reduction at 26 μ g ai/l when compared to the controls. The results of the contrast comparison tests showed a significant reduction in length at 2.6, 15, and 26 μ g ai/l and a significant reduction in weight at 26 μ g ai/l when compared to the dilution water control.

C. <u>Discussion/Results</u>: Although the temperature of the test solutions ranged from 10 to 19°C, it is not likely that this deviation had a significant effect on the growth of the test organisms. This temperature range (10-19°C) represents two excursions from the general temperature (10-14°C) which each lasted 12 hours or less and were caused by a malfunction of the chiller of the water bath.

The author's statistical analysis of length data demonstrated a significant reduction at 2.6 and 15.0 μg ai/l from the pooled control, a difference which the author claimed to be biologically insignificant. The reviewer's results showed a significant reduction in length at 2.6, 15.0, and 26.0 μg ai/l. The reviewer agrees with the author that the difference at 2.6 μg ai/l is probably not due to the treatment since length at 6.6 μg ai/l was not statistically reduced. However, the reduction at 15 μg ai/l may be treatment related, therefore, a conservative estimate of the NOEC would be 6.6 μg ai/l.

This study is scientifically sound and meets the guideline requirements for a fish early life-stage test. Based on the effects of M&B 46030 on larval

length, the MATC for rainbow trout was >6.6 and <15 μg ai/l (geometric mean MATC = 9.9 μg ai/l).

- D. Adequacy of the Study:
 - (1) Classification: Core.
 - (2) Rationale: N/A.
 - (3) Repairability: N/A.
- 15. COMPLETION OF ONE-LINER: Yes; 14 January 1994.

M&B 46030: Survival of Exposed P.prometas Larvae

File: 42918627.sur

Transform: ARC SINE(SQUARE ROOT(Y))

Chi-square test for normality: actual and expected frequencies

INTERVAL	<-1.5	-1.5 to <-0.5	-0.5 to 0.5	>0.5 to 1.5	>1.5
EXPECTED OBSERVED	0.938 0	3.388	5.348 2	3.388	0.938 0

Calculated Chi-Square goodness of fit test statistic = 7.9994
Table Chi-Square value (alpha = 0.01) = 13.277

Data PASS normality test. Continue analysis.

M&B 46030: Survival of Exposed P.promelas Larvae

File: 42918627.sur

Transform: ARC SINE(SQUARE ROOT(Y))

Shapiro Wilks test for normality

D = 0.032

W = 0.814

Critical W (P = 0.05) (n = 14) = 0.874Critical W (P = 0.01) (n = 14) = 0.825

Data FAIL normality test. Try another transformation.

Warning - The two homogeneity tests are sensitive to non-normal data and should not be performed.

M&B 46030: Survival of Exposed P.promelas Larvae : 42918627.sur Transform: ARC SINE(SQUARE ROOT(Y))

Hartley test for homogeneity of variance ■ Bartletts test for homogeneity of variance

These two tests can not be performed because at least one group has zero variance.

Data FAIL to meet homogeneity of variance assumption. Additional transformations are useless.

TITLE:

M&B 46030: Survival of Exposed P.promelas Larvae

FILE:

42918627.sur

TRANSFORM: ARC SINE(SQUARE ROOT(Y))

NUMBER OF GROUPS: 7

GRP	IDENTIFICATION	REP	VALUE	TRANS VALUE
1	Control	1	1.0000	1.4588
1	Control	2	0.9500	1.3453
2	Solvent Control	1	1.0000	1.4588
2	Solvent Control	2	0.9500	1.3453
3	2.6 ug ai/l	1	1.0000	1.4588
3.	2.6 ug ai/l	2	0.9500	1.3453
4	6.6 ug ai/l	1	1.0000	1.4588
4	6.6 ug ai/l	2	0.9500	1.3453
5	15 ug ai/l	1	0.9500	1.3453
5	15 ug ai/l	2	0.9000	1.2490
6	26 ug ai/l	1	0.8000	1.1071
6	26 ug ai/l	2	0.7500	1.0472
7	60 ug ai/l	1	0.0000	0.1120
7	60 ug ai/l	2	0.0000	0.1120
_				

M&B 46030: Survival of Exposed P.promelas Larvae

File: 42918627.sur

Transform: ARC SINE(SQUARE ROOT(Y))

KRUSKAL-WALLIS ANOVA BY RANKS - TABLE 1 OF 2 (p=0.05)

GROUP	IDENTIFICATION	TRANSFORMED MEAN	MEAN CALCULATED IN ORIGINAL UNITS	RANK SUM
1	Control	1.402	0.975	20.500
ż	Solvent Control	1.402	0,975	20.500
3	2.6 ug ai/l	1.402	0.975	20.500
4	6.6 ug ai/l	1.402	0.975	20.500
5	15 ug ai/l	1.297	0.925	13.000
6	26 ug ai/l	1.077	0.775	7.000
	60 ug ai/l	0.112	0.000	3.000

Calculated H Value = 10.210

Critical H Value Table = 12.590

Since Calc H < Crit H FAIL TO REJECT Ho: All groups are equal.

M&B 46030: Survival of Exposed P.promelas Larvae

File: 42918627.sur

Transform: ARC SINE(SQUARE ROOT(Y))

DUNNS MULTIPLE COMPARISON - KRUSKAL-WALLIS - TABLE 2 OF 2 (p=0.05)

						GI	RO	JP		
		TRANSFORMED	ORIGINAL	0	0	0	0	0	0	0
GROUP	IDENTIFICATION	MEAN	MEAN	7	6	5	4	1	2	3
				-	-	•	-	-	-	-
7	60 ug ai/l	0.112	0.000	١						
6	26 ug ai/l	1.077	0.775	•	١					
5	15 ug ai/l	1.297	0.925		7,0	١				
4	6.6 ug ai/l	1.402	0.975	,•	-		1			
1	Control	1.402	0.975	•	-	•		١		
2	Solvent Control	1.402	0.975	•		٠.			1	
3	2.6 ug ai/l	1.402	0.975	•		,				1
								*		

^{* =} significant difference (p=0.05) Table q value (0.05,7) = 3.038

. = no significant difference

SE = 4.038

Solvent Control

2 = Dilution Water Control TRT 3 = 6.2 ug ai/l nominal 4 = 12 ug ai/lTRT 5 = 25 ug ai/l6 = 50 ug ai/lTRT REP LENGTH WEIGHT CASE 1.0000 2.0184 1.0000 59,0000 2 1.0000 CASE 1.0000 68.0000 2.9042 CASE 1.0000 1.0000 67.0000 2.9512 CASE 4 1.0000 4.0C00 62,0000 2.1924 5 1.0000 63.0000 CASE 1.0000 2.4922 67 CASE 1.0000 1.0000 60.0000 1.8659 CASE 1.0000 1.0000 59.0000 1.8044 CASE 8 1.0000 1.0000 65.0000 2.8730 CASE 9 1.0000 1.0000 68.0000 3.1342 CASE 10 1.0000 1.0000 58.0000 2.0128 CASE 11 1.0000 1.0000 61.0000 2.1073 CASE 12 1.0000 1.0000 52.0000 1.5091 CASE 13 1.0000 1.0000 67.0000 2.6213 CASE 14 1.0000 1.0000 61.0000 2.1185 CASE 15 1.0000 1.0000 66.0000 2.4689 CASE 16 1.0000 65.0000 1.0000 2.4956 17 CASE 1.0000 1.0000 58.0000 1.7375 CASE 18 1.0000 1.0000 58.0000 1.7073 CASE 19 58.0000 1.0000 1.0000 2.1178 CASE 20 1.0000 2.0000 61.0000 2.3766 CASE 21 1.0000 2.0000 58.0000 2.0160 CASE 22 1.0000 63.0000 2.0000 2.5023 23 24 CASE 1.0000 2.0000 61.0000 2.3050 CASE 1.0000 2.0000 59.0000 2.0099 61.0000 CASE 25 1.0000 2,0000 2.4929 CASE 26 1.0000 2.0000 57.0000 1.7915 CASE 27 1.0000 2.0000 63.0000 2.3305 CASE 28 1.0000 2.0000 61.0000 2.2203 29 1.0000 CASE 2.0000 58.0000 1.9859 30 CASE 1.0000 2.0000 57.0000 1.9055 0.4772 CASE 31 1.0000 2.0000 39.0000 CASE 32 1.0000 2,0000 63.0000 2.4049 33 34 1.0000 CASE 2.0000 64.0000 2.3151 CASE 1.0000 2.0000 58.0000 1.9241 CASE 35 1.0000 2.0000 64.0000 2.5178 CASE 36 1.0000 2.0000 59.0000 1.9004 37 CASE 1.0000 2.0000 72.0000 3.7679 CASE 38 1.0000 2.0000 59.0000 2.1221 39 1.0000 CASE 2.0000 58.0000 2.0005 40 CASE 2.0000 1.0000 62.0000 2.4568 CASE 41 2.0000 1.0000 63.0000 2.2853 42 CASE 2.0000 1.0000 64.0000 2.6258 43 CASE 2.0000 1.0000 56.0000 1.6325 CASE 44 2.0000 1.0000 59.0000 2.1117 CASE 45 2.0000 1.0000 59.0000 1.9496 46 47 CASE 2.0000 1.0000 57.0000 2.1573 CASE 2.0000 1.0000 64.0000 2.3289 CASE 48 2.0000 1.0000 61.0000 1.9198 CASE 49 2.0000 1.0000 59.0000 1.9873 50 CASE 2.0000 1.0000 62.0000 2.2965 CASE 51 2.0000 1.0000 59.0000 1,9357 CASE 52 2.0000 1.0000 60.0000 2.0329

Dilution Water pur

TRT 1 = Solvent Control

CASE	53	2.0000	1.0000	59.0000	2.1616
CASE	54	2.0000	1.0000	58.0000	1.8451
	55	2.0000			
CASE			1.0000	59.0000	2.1197
CASE	56	2.0000	1.0000	58.0000	1.7856
CASE	57	2.0000	1.0000	63.0000	2.4197
CASE	58	2.0000	1.0000	61.0000	1.9661
CASE	59	2.0000	2.0000	61.0000	2.5143
CASE	60	2.0000	2.0000	62.0000	2.2648
CASE	61	2.0000	2.0000	60.0000	2.1687
CASE	62	2.0000	2.0000	59.0000	2.0143
CASE	63	2.0000	2.0000	58.0000	1.9720
CASE	64	2.0000	2.0000	59.0000	1.9948
CASE	65	2.0000	2.0000	60.0000	2.3425
CASE	66	2.0000	2.0000	53.0000	1.4280
CASE	67	2.0000	2.0000	56.0000	1.6978
CASE	68	2.0000	2.0000	62.0000	2.3799
CASE	69	2.0000	2.0000	63.0000	2.4058
CASE -		2. 0000	2.0000	61.0000	2.0680
CASE	71	2.0000	2.0000	62.0000	2.2360
CASE	72	2.0000	2.0000	59.0000	2.2153
CASE	73	2.0000	2.0000	58.0000	1.9963
CASE	74	2.0000	2.0000		
				63.0000	2.4272
CASE	75	2.0000	2.0000	56.0000	1.6888
CASE	76	2.0000	2.0000	60.0000	1.9401
CASE	77	2.0000	2.0000	64.0000	2,4028
CASE	78	2.0000		56.0000	
			2.0000		1.7299
CASE	79	3.0000	1.0000	54.0000	2.1309
CASE	80	3.0000	1.0000	62.0000	2.2239
CASE	81	3.0000	1.0000	65.0000	2.8293
CASE	82	3.0000	1.0000	50.0000	
					1.5195
CASE	83	3.0000	1.0000	51.0000	2.0544
CASE	84	3.0000	1.0000	60.0000	2.1598
CASE	85	3.0000	1.0000	54.0000	2,2389
CASE	86	3.0000	1.0000		2.5677
				65.0000	
CASE	87	3.0000	1.0000	60.0000	2.1518
CASE	88	3.0000	1.0000	52.0000	1.8591
CASE	89	3.0000	1.0000	54.0000	1.4865
CASE	90	3.0000	1.0000	64.0000	2.4456
CASE	91	3.0000	1.0000	59.0000	2.1462
CASE	92	3.0000	1.0000	52.0000	1.3032
CASE	93	3.0000	1.0000	63.0000	2.7105
CASE	94	3.0000	1.0000	63.0000	2.2636
CASE	95	3.0000	1.0000	51.0000	1.7073
CASE	96	3.0000	1.0000	65.0000	2.7045
CASE	97	3.0000	1.0000	60.0000	2.2563
CASE	98	3.0000	1.0000	53.0000	1.9363
CASE	99	3.0000	2.0000	57.0000	1.7200
CASE	100	3.0000	2.0000	55.0000	2.3392
CASE	101	3.0000	2.0000	62.0000	2.3467
CASE	102	3.0000	2.0000	58.0000	2.1312
CASE	103	3.0000	2.0000	59.0000	2.0862
CASE	104	3.0000	2.0000	58.0000	1.9607
CASE	105	3.0000	2.0000	65.0000	2.5621
CASE	106	3.0000	2.0000	63.0000	2.4596
CASE	107	3.0000	2.0000	60.0000	1.9556
CASE	108	3.0000	2.0000	57.0000	2.0618
CASE	109	3.0000	2.0000	56.0000	2.2787
CASE	110	3.0000	2.0000	60.0000	2.0037
CASE	111	3.0000	2.0000	59.0000	2.1247
CASE	112	3.0000	2.0000	52.0000	1.8789
CASE	113	3.0000	2.0000	53.0000	1.7492
CASE	114	3.0000	2.0000	64.0000	2.5381
CASE	115	3.0000	2.0000	56.0000	2.1130
CASE	116	3.0000	2.0000	58.0000	2.0488
CASE	117	3.0000	2.0000	59.0000	2.2896
CASE	118	4.0000	1.0000	65.0000	2.8410
UNDE	. 10	4.0000		05.0000	E . 04 LU

CASE	119	4.0000	1.0000	63.0000	2.4224
CASE	120	4.0000	1.0000	59.0000	1.8317
CASE	121	4.0000	1.0000	60.0000	2.0447
CASE	122	4.0000	1.0000	62.0000	2.3103
CASE	123	4.0000	1.0000	62.0000	2.0585
CASE	124	4.0000	1.0000	59.0000	2.0887
	125			57.0000	
CASE	126	4.0000	1.0000		1.8493
CASE		4.0000	1.0000	59.0000	2.0620
CASE	127	4.0000	1.0000	57.0000	1.9968
CASE	128	4.0000	1.0000	58.0000	1.8490
CASE	129	4.0000	1.0000	59.0000	2.1758
CASE	130	4.0000	1.0000	58.0000	1.5330
CASE	131	4.0000	1.0000	60.0000	2.0160
CASE	132	4.0000	1.0000	60.0000	2.1176
CASE	133	4.0000	1.0000	61.0000	2.0980
CASE	134	4.0000	1.0000	63.0000	2.7727
CASE	135	4.0000	1.0000	56.0000	1.7871
CASE	136	4.0000	1 .0000	60.0000	2.2197
CASE	137	4.0000	1.0000	58.0000	1.9759
CASE	138	4.0000	2.0000	65.0000	2.8434
CASE	139	4.0000	2.0000	61.0000	2.1875
CASE	140	4.0000	2.0000	55.0000	1.9014
CASE	141	4.0000	2.0000	58.0000	2.1235
CASE	142	4.0000	2.0000	64.0000	2.4352
CASE	143	4.0000	2.0000	63.0000	2.7455
CASE	144	4.0000	2.0000	61.0000	2.1321
CASE	145	4.0000	2.0000	58.0000	1.9478
CASE	146	4.0000	2.0000	60.0000	2.0346
CASE	147	4.0000	2.0000	63.0000	2.3820
CASE	148	4.0000	2.0000	56.0000	1.7395
CASE	149	4.0000	2.0000	58.0000	1.9617
CASE	150	4.0000	2.0000	59.0000	2.0867
CASE	151	4.0000	2.0000	57.0000	1.9550
CASE	152	4.0000	2.0000	58.0000	1.9062
CASE	153	4.0000	2.0000	62.0000	2.3528
CASE	154	4.0000	2.0000	61.0000	2.2753
CASE	155	4.0000	2.0000	62.0000	2.7297
CASE	156	4.0000	2.0000	58.0000	1.9854
CASE	157	5.0000	1.0000	60.0000	1.9224
CASE	158	5.0000	1.0000	62.0000	2.4270
CASE	159	5.0000	1.0000	61.0000	2.2892
CASE	160	5.0000	1.0000	59.0000	2.3693
CASE	161	5.0000	1.0000	65.0000	3.0089
CASE	162	5.0000	1.0000	60.0000	2.3342
CASE	163	5.0000	1.0000	61.0000	2.5554
CASE	164	5.0000	1.0000	50.0000	1.3835
CASE	165	5.0000	1.0000	56.0000	2.0277
CASE	166	5.0000	1.0000	57.0000	1.9054
CASE	167	5.0000	1.0000	59.0000	1.8905
CASE	168	5.0000	1.0000	56.0000	1.6257
CASE	169	5.0000	1.0000	59,0000	1.9689
CASE	170	5.0000	1.0000	58.0000	2.1062
CASE	171	5.0000	1.0000	54.0000	1.5689
CASE	172	5.0000	1.0000	59.0000	1.9713
CASE	173	5.0000	1.0000	63.0000	2.4546
CASE	174	5.0000	1.0000	56.0000	1.9098
CASE	175	5.0000	1.0000	55.0000	1.7398
CASE	176	5.0000	2.0000	59.0000	2.2797
CASE	177	5.0000	2.0000	58.0000	2.0629
CASE	178	5.0000	2.0000	67.0000	3.1215
CASE	179	5.0000	2.0000	59.0000	2.0080
CASE	180	5.0000	2.0000	60.0000	2.0834
	181	5.0000	2.0000	57.0000	2.0005
CASE					2.3947
CASE	182 183	5.0000 5.0000	2.0000 2.0000	60.0000 57.0000	1.8599
CASE CASE	184	5.0000	2.0000	58.0000	2.1013
ONOL	.07	J.0000	2.0000	30.0000	£. 1015

	CASE	185	5.0000	2.0000	54.0000	1.8913
	CASE	186	5.0000	2.0000	50.0000	1.1821
	CASE	187	5.0000	2.0000	58.0000	1.9492
	CASE	188	5.0000	2.0000	56.0000	1.8872
	CASE	189	5.0000	2.0000	61.0000	2.3735
	CASE	190	5.0000	2.0000	54.0000	1.8578
	CASE	191	5.0000	2.0000	60.0000	2.3265
	CASE	192	5.0000	2.0000	61.0000	2.4760
	CASE	193	5.0000	2.0000	57.0000	1.8513
	CASE	194	6.0000	1.0000	27.0000	0.5724
	CASE	195	6.0000	1.0000	60.0000	2.4954
	CASE	196	6.0000	1.0000	57.0000	2.0504
	CASE	197	6.0000	1.0000	37.0000	1.1052
	CASE	198	6.0000	1.0000	38.0000	0.9652
	CASE	199	6.0000	1.0000	52.0000	1.8383
	CASE	200	6.0000	1.0000	60.0000	2.2801
	CASE	201	6.0000	1.0000	57.0000	2.4415
ķ .	CASE_	202	&_000 0	1.0000	58.0000	2:0557
	CASE	203	6.0000	1.0000	51.0000	1.5620
	CASE	204	6.0000	1.0000	56.0000	1.9092
	CASE	205	6.0000	1.0000	51.0000	1.6476
	CASE	206	6.0000	1.0000	41.0000	0.9155
	CASE	207	6.0000	1.0000	57.0000	1.8582
	CASE	208	6.0000	1.0000	36.0000	0.5597
	CASE	209	6.0000	2.0000	55.0000	1.9588
	CASE	210	6.0000	2.0000	53.0000	1.6930
	CASE	211	6.0000	2.0000	60.0000	2.6144
	CASE	212	6.0000	2.0000	41.0000	1.3414
	CASE	213	6.0000	2.0000	50.0000	1.8974
	CASE	214	6.0000	2.0000	58.0000	2.5183
	CASE	215	6.0000	2.0000	56.0000	1.7978
	CASE	216	6.0000	2.0000	58.0000	1.9550
	CASE	217	6.0000	2.0000	41.0000	1.1683
	CASE	218	6.0000	2.0000	53.0000	1.7232
	CASE	219	6.0000	2.0000	55.0000	1.8575
	CASE	220	6.0000	2.0000	47.0000	1.5572
	CASE	221	6.0000	2.0000	32.0000	0.5257
	CASE	222	6.0000	2.0000	46.0000	1.5460
	CASE	223	6.0000	2.0000	50.0000	1.4117

ANOVA on Weights

	1.0000	2.0000	3.0000	4.0000	5.0000	6.000
REP	1.0000	2.0000				
EP VA	R: WEIGHT	N: 2	23 MULTIPLE R:	0.393 SQUA	RED MULTIPLE	R: 0.15
		ANALYSI	S OF VARIANCE			
SOURCE	SUM-	OF-SQUARES	DF MEAN-SQUARE	F-RATIO	P	
TRT			1.2707	7.4604	0.0000	
REP		Annual Control of the	1 0.0124	0.0727	0.7877	
rrt*re	۲	0.2399	5 ~ 0.0480	0.2817	0.9228	
ERROR		35.9381 21		.*		
	-WATSON D S		.071			
IRST	ORDER AUTOC	ORRELATION -	.037			
OST H	OC TEST OF	WEIGHT				
-,						
MATRIX	 	E MEAN DIFFER	ENCES:			
MATRIX	 	E MEAN DIFFER	ENCES:	3	4	5
MATRIX	OF PAIRWIS			3	4	5
MATRIX	OF PAIRWIS	1 0.0000 -0.1189	2		4	5
MATRIX	OF PAIRWIS	1 0.0000 -0.1189 -0.0822	2 0.0000 0.0368	0.0000		5
MATRIX	OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699	2 0.0000 0.0368 0.0490	0.0000 0.0122	0.0000	
MATRIX	OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334	0.0000 0.0368 0.0490 -0.0145	0.0000 0.0122 -0.0513	0.0000 -0.0635	0.0000
MATRIX	OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699	2 0.0000 0.0368 0.0490	0.0000 0.0122	0.0000	
MATRIX	OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585	0.0000 0.0368 0.0490 -0.0145	0.0000 0.0122 -0.0513	0.0000 -0.0635	0.0000
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000	0.0000 0.0368 0.0490 -0.0145 -0.4395	0.0000 0.0122 -0.0513	0.0000 -0.0635	0.0000
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000	0.0000 0.0368 0.0490 -0.0145	0.0000 0.0122 -0.0513	0.0000 -0.0635	0.0000
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000	0.0000 0.0368 0.0490 -0.0145 -0.4395	0.0000 0.0122 -0.0513	0.0000 -0.0635	0.0000
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES:	0.0000 0.0122 -0.0513 -0.4763	0.0000 -0.0635 -0.4886	0.0000 -0.4251
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST OF PAIRWIS	0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES:	0.0000 0.0122 -0.0513 -0.4763	0.0000 -0.0635 -0.4886	0.0000 -0.4251
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON 1 1.0000 1.0000 1.0000	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES:	0.0000 0.0122 -0.0513 -0.4763	0.0000 -0.0635 -0.4886	0.0000 -0.4251
BONFER	OF PAIRWIS 1 2 3 4 5 6 RONI ADJUST OF PAIRWIS 1 2 3 4	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON 1 1.0000 1.0000 1.0000	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES: 2 1.0000 1.0000 1.0000	0.0000 0.0122 -0.0513 -0.4763 3	0.0000 -0.0635 -0.4886 4	0.0000 -0.4251 5
BONFER	OF PAIRWIS 1 2 3 4 5 6 RONI ADJUST OF PAIRWIS 1 2 3 4	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON 1 1.0000 1.0000 1.0000 1.0000	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES: 2 1.0000 1.0000 1.0000 1.0000	0.0000 0.0122 -0.0513 -0.4763 3 1.0000 1.0000 1.0000	0.0000 -0.0635 -0.4886 4 1.0000 1.0000	0.0000 -0.4251 5
BONFER	OF PAIRWIS 1 2 3 4 5 6 RRONI ADJUST OF PAIRWIS	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON 1 1.0000 1.0000 1.0000	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES: 2 1.0000 1.0000 1.0000	0.0000 0.0122 -0.0513 -0.4763 3	0.0000 -0.0635 -0.4886 4	0.0000 -0.4251 5
BONFER	OF PAIRWIS 1 2 3 4 5 6 RONI ADJUST OF PAIRWIS 1 2 3 4	1 0.0000 -0.1189 -0.0822 -0.0699 -0.1334 -0.5585 6 0.0000 MENT. E COMPARISON 1 1.0000 1.0000 1.0000 1.0000	2 0.0000 0.0368 0.0490 -0.0145 -0.4395 PROBABILITIES: 2 1.0000 1.0000 1.0000 1.0000	0.0000 0.0122 -0.0513 -0.4763 3 1.0000 1.0000 1.0000	0.0000 -0.0635 -0.4886 4 1.0000 1.0000	0.0000 -0.4251 5

ANOVA on Lengths

	ENCOUNTERE	D DURING PROCE	SSING ARE:			
TRT	1.0000	2.0000	3.0000	4.0000	5.0000	6.0000
REP	1.0000	2.0000				
DEP VA	R: LENGTH		3 MULTIPLE R: OF VARIANCE	0.584 SQU/	ARED MULTIPLE	R: 0.342
SOURCE	SUM-		F MEAN-SQUARE	F-RATIO) P	
TRT		2613.7275 5	522.7455	21.4541		
REP TRT*RE	:р	1.7088 1 57.3201 5		0.0701 0.4705		
	I-WATSON D ORDER AUTO	5141.1795 211 STATISTIC 2. CORRELATION	052		-	
USING POST H	LEAST SQUA		 	roni.		
		1	2	3	4	5
	1	0.0000				
	2	-0.9171	0.0000			
	3	-2.6342	-1.7171	0.0000		54
	4	-0.9224	-0.0053	1.7118	0.0000	
	5	-2.5300	-1.6129	0.1042	-1.6076	0.0000
	6	-11.0294	-10.1123	-8.3952	-10.1070	-8.4994
		6				
	6	0.0000				
	RRONI ADJUS K OF PAIRWI	TMENT. SE COMPARISON F	PROBABILITIES:			
		1	2	3	4	5
	1	1.0000				
	ż	1.0000	1.0000			
	3	0.2910	1.0000	1.0000		
		1.0000	1.0000	1.0000	1.0000	
	5	0.3994	1.0000	1.0000	1.0000	1.0000
	4 5 6	0.0000	0.0000	0.0000	0.0000	0.0000
		6				
	6	1.0000				
•	U	1.0000				

THE FOLLOWING RESULTS ARE FOR:

1.0000 1.0000

TOTAL OBSERVATIONS:

	WEIGHT	LENGTH
N OF CASES	19	19
MINIMUM	1.5091	52.0000
MAXIMUM	3.1342	68.0000
MEAN	2.2701	61.8421
STANDARD DEV	0.4718	4.3750

THE FOLLOWING RESULTS ARE FOR:

1.0000

2.0000

TOTAL OBSERVATIONS: 20

	WEIGHT	LENGTH
N OF CASES	20	20
MINIMUM	0.4772	39.0000
MAXIMUM	3.7679	72.0000
MEAN	2.1683	59.7500
STANDARD DEV	0.5795	5.9989

THE FOLLOWING RESULTS ARE FOR:
TRT = 2.0000

1.0000

TOTAL OBSERVATIONS: 19

	WEIGHT	LENGTH
N OF CASES	19	19
MINIMUM	1.6325	56.0000
MAXIMUM	2.6258	64.0000
MEAN	2.1062	60.1579
STANDARD DEV	0.2514	2.3396

THE FOLLOWING RESULTS ARE FOR:

TRT = 2.0000
2.0000

TOTAL OBSERVATIONS:

WEIGHT LENGTH N OF CASES MINIMUM 20 1.4280 20 53.0000 64.0000 59.6000 2.8359 2.5143 MAXIMUM 2.0944 MEAN STANDARD DEV

THE FOLLOWING RESULTS ARE FOR:

TRT = 3.0000

REP = 1.0000

TOTAL OBSERVATIONS: 20

 WEIGHT
 LENGTH

 N OF CASES
 20
 20

 MINIMUM
 1.3032
 50.0000

 MAXIMUM
 2.8293
 65.0000

 MEAN
 2.1348
 57.8500

 STANDARD DEV
 0.4148
 5.4703

THE FOLLOWING RESULTS ARE FUR:

TRT = 3.0000

REP = 2.0000

TOTAL OBSERVATIONS: 19

 WEIGHT
 LENGTH

 N OF CASES
 19
 19

 MINIMUM
 1.7200
 52.0000

 MAXIMUM
 2.5621
 65.0000

 MEAN
 2.1394
 58.4737

 STANDARD DEV
 0.2430
 3.4378

THE FOLLOWING RESULTS ARE FOR:

TRT = 4.0000

REP = 1.0000

TOTAL OBSERVATIONS: 20

WEIGHT LENGTH 20 N OF CASES 56.0000 MINIMUM 1.5330 2.8410 65.0000 MAXIMUM MEAN 2.1025 59.8000 STANDARD DEV 0.3101 2.3079

THE FOLLOWING RESULTS ARE FOR:

TRT = 4.0000

REP = 2.0000

TOTAL OBSERVATIONS:

19

WEIGHT LENGTH N OF CASES 19 19 55.0000 65.0000 MINIMUM 1.7395 2.8434 MAXIMUM MEAN 2.1961 59.9474 STANDARD DEV 0.3139 2.7983

THE FOLLOWING RESULTS ARE FOR:

TRT 5.0000

1.0000 REP

TOTAL OBSERVATIONS:

•	WEIGHT	LENGTH	
N OF CASES	19	19	
MINIMUM	1.3835	50.0000	
MAXIMUM	3.0089	65.0000	
MEAN	2.0768	58.4211	
STANDARD DEV	0.3914	3.4851	

THE FOLLOWING RESULTS ARE FOR:

5.0000 TRT

2.0000

TOTAL OBSERVATIONS:

	WEIGHT	LENGTH
N OF CASES	18	18
MINIMUM	1.1821	50.0000
MAXIMUM	3.1215	67.0000
MEAN	2.0948	58.1111
STANDARD DEV	0.3897	3.5627

THE FOLLOWING RESULTS ARE FOR:

TRT 6.0000

1.0000 REP

TOTAL OBSERVATIONS:

	WEIGHT	LENGTH	
N OF CASES	15	15	
MINIMUM	0.5597	27.0000	
MAXIMUM	2.4954	60.0000	
MEAN	1.6171	49.2000	
STANDARD DEV	0.6467	10.5641	

THE FOLLOWING RESULTS ARE FOR:

6.0000 TRT 2.0000 REP

TOTAL OBSERVATIONS: 15

WEIGHT LENGTH N OF CASES 0.5257 32.0000 MINIMUM 2.6144 60.0000 50.3333 MAXIMUM MEAN STANDARD DEV 0.5091 7.7521

SUMMARY STATISTICS FOR WEIGHT

BARTLETT TEST FOR HOMOGENEITY OF GROUP VARIANCES

CHI-SQUARE =

37.0695 DF= 11 PROBABILITY =

0.0001

3.5119

ANALYSIS OF VARIANCE

SOURCE

SUM OF SQUARES DF MEAN SQUARE

PROBABILITY

BETWEEN GROUPS WITHIN GROUPS

6.5797 11 35.9381 211

0.5982 0.1703 0.0002

SUMMARY STATISTICS FOR LENGTH

BARTLETT TEST FOR HOMOGENEITY OF GROUP VARIANCES

CHI-SQUARE =

91.0994 DF= 11 PROBABILITY =

0.0000

9.9523

ANALYSIS OF VARIANCE

SOURCE

SUM OF SQUARES DF MEAN SQUARE

PROBABILITY

BETWEEN GROUPS WITHIN GROUPS

2667.4483 11 5141.1795 211

242.4953 24.3658 0.0000

KOLMOGOROV-SMIRNOV ONE SAMPLE TEST USING STANDARD NORMAL DISTRIBUTION

VARIABLE

N-OF-CASES MAXDIF PROBABILITY (2-TAIL)

WEIGHT

223.0000

0.8686

0.0000

LENGTH -

223.0000

1.0000

0.0000

LEVELS ENCOUNT TRT 1.000 REP 1.000	ERED DURING F 2.000 2.000	ROCESSII 3.00		000 5.00	00 6.000
DEP VAR: LENG			LE R: 0.584 VARIANCE	SQUARED MULTI	PLE R: 0.342
SOURCE S	UM-OF-SQUARES	DF 1	MEAN-SQUARE	F-RATIO	P
TRT REP TRT*REP	2613.728 1.709 57.320		522.746 1.709 11.464	21.454 0.070 0.470	0.000 0.791 0.798
ERROR	5141.180			01410	0.17,0
Post-hoc contr TEST FOR EFFEC TEST OF HYPOTH	T CALLED:	ent 1 w	1		
SOURCE	SS	DF	MS	F	P
HYPOTHESIS ERROR		1 211	16.390 24.366	0.673	0.413
Post-hoc contra TEST FOR EFFEC	ast of treatm T CALLED:		ith control. F OF HYPOTHE		
SOURCE	SS	DF	MS	· F	·P
HYPOTHESIS ERROR	135.223 5141.180	1 211	135.223 24.366	5.550	0.019
Post-hoc contra TEST FOR EFFEC TEST OF HYPOTHI	T CALLED:	ent 3 wi	ith control.		ngang palipag di sagal na mang mang mang mang mang mang mang m
SOURCE	SS	DF	MS	F	Р
HYPOTHESIS ERROR	16.579 5141.180	1 211	16.579 24.366	0.680	0.410
Post-hoc contra TEST FOR EFFEC TEST OF HYPOTH	CALLED:	ent 4 wi TRT	ith control.		
SOURCE	ss	DF	MS	F	P
HYPOTHESIS ERROR	121.446 5141.180	1 211	121.446 24.366	4.984	0.027
Post-hoc contra TEST FOR EFFECT TEST OF HYPOTHE	CALLED:	ent 5 wi TRT	th control.	tarian para esta de la composição de la co	energetty i i i i i i i i i i i i i i i i i i
SOURCE	SS	DF	MS	F	Р
HYPOTHESIS ERROR	2062.126 5141.180	1 211	2062.126 24.366	84.632	0.000

LEVELS ENCOUNTER FRT 1.000 REP 1.000	2.000 2.000		ING ARE: 000 4.00	5.000	6.000
EP VAR: WEIGHT	N:	223	MULTIPLE R: 0.	393 SQUARED I	MULTIPLE R: 0.155
SOURCE SUM			F VARIANCE MEAN-SQUARE	F-RATIO	P
RT REP RT*REP	6.353 0.012 0.240	5 1 5	1.271 0.012 0.048	7.460 0.073 0.282	0.000 0.788 0.923
RROR	35.938	211	0.170		
Post-hoc contras EST FOR EFFECT EST OF HYPOTHES	CALLED:	ent 1	with control.	· · · · · · · · · · · · · · · · · · ·	
SOURCE	ss	DF	MS	F	Р
HYPOTHESIS ERROR	0.276 35.938		0.276 0.170	1.618	0.205
Post-hoc contras EST FOR EFFECT EST OF HYPOTHES	CALLED:	nent 2 TRT	with control.		
SOURCE	SS	DF	MS	F	P
HYPOTHESIS ERROR	0.132 35.938		0.132 0.170	0.772	0.381
Post-hoc contras TEST FOR EFFECT TEST OF HYPOTHES	CALLED:		with control.	Andreas Andreas Andreas	
SOURCE	ss	DF	MS	F	P
HYPOTHESIS ERROR	0.095 35.938		0.095 0.170	0.559	0.455
Post-hoc contras TEST FOR EFFECT TEST OF HYPOTHES	CALLED:	nent 4 TRT	with control.	in in the second se	
SOURCE	SS	DF	MS	F	P
HYPOTHESIS ERROR	0.338 35.938	1 211	0.338 0.170	1.983	0.161
Post-hoc contras TEST FOR EFFECT TEST OF HYPOTHES	CALLED:	nent 5 TRT	with control.	 	•
SOURCE	SS	DF	MS	F	P
HYPOTHESIS	5.287 35.938	1 211	5.287 0.170	31.042	0.000

Ecological Effects Branch One-Liner Data Entry Form

Pesticide Use_ Chemical Fibronil and derivatives EUP (M&B 46030) Shaughnessy No. 129121

AQUATIC VERTEBRATE TOX.	% AI	LC ₅₀ (95%CL) BLO PE	HRS/ TYPE	NOEC	STUDY/ REVIEW DATES	MRID/ CATEGORY	LAB	RC
1.				۷.				
2.						4.0		
3.								
4.								
5.						#N,		
• 9								
7.								
CHRONIC TOX.	% AI	MATC	DAYS	AFFECTED PARA.	STUDY/ REVIEW DATES	MRID/ CATEGORY	LAB	RC
1. Oncorhynchus mykiss	96.7%	>6.6 <15 µg ai/l	06	length	1992/ 1994	429186-27 Core	гтѕ	RGM
2.						ą.		
3.								

COMMENTS: Results based on mean measured concentrations. SLI=Springborn Laboratories,

Figural Margit

Page is not included in this copy.
Pages 13 hot included in this copy.
The material not included contains the following type of information:
Identity of product inert ingredients.
Identity of product impurities.
Description of the product manufacturing process.
Description of quality control procedures.
Identity of the source of product ingredients.
Sales or other commercial/financial information.
A draft product label.
The product confidential statement of formula.
Information about a pending registration action.
The document is a duplicate of page(s)
The document is not responsive to the request.
The information not included is generally considered confidentia by product registrants. If you have any questions, please contacthe individual who prepared the response to your request.