



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OPP OFFICIAL RECORD
HEALTH EFFECTS DIVISION
SCIENTIFIC DATA REVIEWS
EPA SERIES 361

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

September 13, 2001

MEMORANDUM

SUBJECT: Propargite; Chemical No. 097601. HED's Revised Human Health Risk Assessment for Propargite, Case # 0243. DP Barcode: D276544.

From: Thurston G. Morton, Risk Assessor
Reregistration Branch 4
Health Effects Division (7509C)

Thurston G. Morton
9/13/01

Thru: Susan V. Hummel, Branch Senior Scientist
Reregistration Branch 4
Health Effects Division (7509C)

Susan V. Hummel

To: Dayton Eckerson/Robert McNally
Special Review Branch
Special Review and Reregistration Division (7508W)

Attached is HED's revised human health risk assessment for the acaricide, propargite. The disciplinary science chapters and other supporting documents for Propargite are included as attachments as follows:

Propargite - Report of the Hazard Identification Assessment Review Committee. J.Rowland/P. Wagner (6/22/99)
Propargite - Report of the FQPA Safety Factor Committee. B. Tarplee/J. Rowland (8/12/99)
Revised Q₁* Memorandum (L. Brunsman, 8/2/01)
Recommendation for Selection of Cancer Dose-Response Model and Q₁* for Propargite (R. Kent, 8/14/01, D276502)
Product & Residue Chemistry Chapter. J. Stokes (1/20/00, D250257)
Revised Toxicology Chapter. S. Shallal (5/24/00, D266213)
Revised Occupational and Residential Exposure Assessment. S. Tadayon (7/31/01, D276513)
Revised Anticipated Residues and Acute, Chronic, and Cancer Dietary Exposure and Risk Analyses for the HED Human Health Risk Assessment. T. Morton (5/24/00, D266001)
Revised Cancer Dietary Exposure and Risk Analysis for the HED Human Health Risk Assessment. T. Morton(8/28/01)
Review of Propargite Incident Reports. J. Blondell/M. Spann (12/8/99, D261208)

1.0 EXECUTIVE SUMMARY

Propargite [2-(p-tert-butylphenoxy)cyclohexyl-2-propynyl sulfite] is a non-systemic acaricide currently registered for food/feed uses on a variety of field, fruit, and vegetable crops. Tolerances for residues of propargite in/on food and feed commodities are currently established under 40 CFR §180.259(a) and (b), §185.5000, and §186.5000(a) and are expressed in terms of propargite *per se*.

Hazard Identification

Propargite is an organosulfite acaricide used for the control of agricultural pests. The toxicological database for propargite is complete (See Table 1) and will support reregistration eligibility. In general, based on animal studies, propargite has low acute toxicity via the oral, dermal, and inhalation routes of exposure (Category III), but causes severe eye and skin irritation (Category I).

Propargite is considered **corrosive** and has been placed in Category I **for both eye and dermal irritation** in rabbits. There have also been documented reports of dermal and eye irritation developing in workers exposed to propargite in the field. Evidence for its dermal sensitization potential have been noted; a study that provides conclusive results has not been possible due to the irritating properties of this chemical.

Toxicity Doses And Endpoints Selected For Risk Assessment

On June 3, 1999, the Health Effects Division (HED) Hazard Identification Assessment Review Committee (HIARC) evaluated the toxicology database of propargite, established Reference Doses (RfDs), and selected the toxicological endpoints and doses for occupational exposure risk assessments. All endpoints are based solely on animal toxicity studies.

For acute dietary risk assessment, the NOAEL of 8 mg/kg/day from a developmental toxicity study in rabbits was chosen based on increased incidence of fetuses with fused sternbrae at the LOAEL of 10 mg/kg/day. The acute RfD was calculated using a 10x interspecies and 10x intraspecies uncertainty factor. The acute Population Adjusted Dose (aPAD) was 0.08 mg/kg/day (acute RfD 0.08 mg/kg/day ÷ 1x FQPA safety factor) and is applicable to Females 13-50 years only.

The HIARC reaffirmed use of an RfD of 0.04 mg/kg/day for chronic dietary risk assessments based on the results of a chronic feeding and carcinogenicity study in rats in which the NOAEL was 4 mg/kg/day. The NOAEL of 4 mg/kg/day was based on decreased body weight / body weight gain and increased mortality at the LOAEL of 19 mg/kg/day. The chronic Population Adjusted Dose (cPAD) was 0.04 mg/kg/day (chronic RfD 0.04 mg/kg/day ÷ 1X FQPA safety factor).

For estimating dermal risk, short- and intermediate-term animal studies reflecting oral administration of the pesticide were used, along with a dermal absorption factor of 14%. A 14% dermal absorption factor was selected based on the highest absorption/elimination noted in two submitted studies. This percentage is deemed valid since it corresponds to the amount of propargite which was actually detected in the excretions of animals. For short-term dermal risk assessments, a NOAEL of 6 mg/kg/day was selected based on decreased maternal body weight gain at the maternal systemic oral LOAEL of 8 mg/kg/day in a developmental toxicity study in rabbits. For intermediate-term dermal risk assessments, a NOAEL of 4 mg/kg/day was selected based on reduction in body weight at a parental oral LOAEL of 20 mg/kg/day in a reproductive toxicity study in rats. For long-term dermal risk assessments, a NOAEL of 4 mg/kg/day was selected based on decreased body weight/body weight gain and increased mortality in a chronic feeding and carcinogenicity study in rats at a LOAEL of 20 mg/kg/day. For inhalation exposure risk assessments at all durations, a LOAEL of 0.31 mg/L (50 mg/kg/day) was chosen based on increased mortality in males in an acute inhalation study in rats. The target MOE is 100 for dermal occupational risk assessments. The target MOE is 1000 for inhalation exposure risk assessments because of an additional 10x due to the lack of a NOAEL and the severity of effects at the lowest dose tested.

On January 23, 1992, the Cancer Peer Review Committee (CPRC) determined that based on the evidence presented, propargite was classified a Group B2, "likely" human carcinogen. It was concluded that administration of propargite was associated with the appearance of extremely rare jejunal tumors in male and female Sprague-Dawley rats. There was an increase in the incidence of undifferentiated sarcoma of the jejunum in males and females receiving 800 ppm propargite compared to concurrent and historical controls. A Q_1^* for propargite was calculated as $0.33 \times 10^{-1} \text{ (mg/kg/day)}^{-1}$ using the 3/4 scaling factor as documented in several documents (L. Brunsman, 8/2/01 and R. Kent, 8/14/01, D276502).

The FQPA Safety Factor Committee met on August 9, 1999 to evaluate hazard and exposure data for propargite and recommend application of the FQPA Safety Factor (as required by Food Quality Protection Act of August 3, 1996), to ensure the protection of infants and children from exposure to propargite. Based on the following: 1) lack of increased susceptibility following *in utero* exposure to rats and rabbits and pre/post natal exposure to rats; 2) adequacy of the database; 3) no currently registered residential uses; and 4) the exposure assessments will not underestimate the potential dietary (food and drinking water) exposures for infants and children from the use of propargite, the FQPA committee recommended reduction of the FQPA Safety Factor to 1X.

Dietary Exposure and Risk Assessment (General Population)

Registered propargite end-use products include emulsifiable concentrate (EC) and wettable powder (WP) formulations. Depending on the crops, these formulations may be applied as broadcast, banded or directed spray or chemigation foliar treatments pre- or postharvest using ground or aerial equipment. Single application rates range from 0.8 to 4.5 pounds per acre.

Preharvest intervals range from 7 to 60 days. The nature of the residue in plants and animals is adequately understood. The HED Metabolism Assessment Review Committee (MARC) concluded the residue of concern in plants and animals is propargite *per se*. Analytical methods are available for enforcing propargite tolerances in PAM II. For most commodities, adequate storage stability data are available. Additional storage stability data are required for an oily commodity to support residue studies on peanut and walnut, and storage stability data are required to support corn and peanut processing studies.

Adequate field trials are available pending submission of required storage stability data, sample storage information, or required label amendments. Adequate processing studies have been submitted for potatoes, citrus, field corn, grapes and peanuts. However, storage stability data are required to support the corn and peanut processing studies.

The reregistration requirements for animal feeding studies are fulfilled. Acceptable ruminant and poultry feeding studies have been submitted and evaluated.

The metabolism of propargite in rotated crops is similar to that in primary crops. Based on an adequate confined rotational crop study and limited field rotational crop studies, the Agency concluded that a six-month plantback interval (PBI) for root crops and a two-month PBI for all other crops are acceptable.

Estimated acute dietary exposure is below HED's level of concern for all female subpopulations at the 99.9th percentile. Use of USDA Pesticide Data Program (PDP) monitoring data, field trial data, and calculated livestock anticipated residues (ARs) results in a dietary risk estimate of 2 % of the aPAD for Females (13-50 years).

Estimated chronic dietary exposure is below HED's level of concern. Use of PDP monitoring data, field trial data, and calculated livestock ARs results in a maximum risk of <1 % of the chronic PAD (% cPAD) for the U.S. Population and all subpopulations.

The cancer dietary exposure and risk estimate for propargite is 1.8×10^{-7} . Results of the analyses indicate potential residues in milk are one of the contributors to the estimated exposure and risk. There were no detections of propargite in PDP data for milk and the highest residue in the 2x feeding study was at the 0.011 ppm. A sensitivity analysis was performed by inserting zeroes for the milk commodities resulting in a cancer dietary exposure and risk estimate of 1.6×10^{-7} .

Drinking Water

Acute drinking water levels of concern (DWLOCs) were calculated based on the acute dietary (food) exposure and default body weights and water consumption figures. The acute DWLOC for Females 13-50 years is 2400 ppb. The EECs for surface water (PRZM-EXAMS) and groundwater (SCI-GROW) were less than the acute DWLOC's, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The peak PRZM-EXAMS EEC was 34 ppb, while the estimated groundwater EEC was 0.006 ppb.

The EECs for surface water (PRZM-EXAMS) and groundwater (SCI-GROW) were less than the chronic DWLOCs, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The average PRZM-EXAMS EEC was 8.7 ppb, while the estimated SCI-GROW EEC was 0.006 ppb.

A cancer DWLOC was calculated based on the cancer dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were greater than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is greater than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). The PRZM-EXAMS and SCI-GROW EECs were 4.8 and 0.006 ppb, respectively. Surface water concentrations below 1.0 ppb would result in cancer risks below 1×10^{-6} for drinking water alone when back calculated. Time weighted average propargite concentration in surface water samples from the USGS NAWQA (Oristimba Creek Watershed) for the years 1992-1993 were 0.30 and 1.24 ppb, respectively. Therefore, even when monitoring data are used cancer exposure to propargite from surface water sources is greater than HED's level of concern.

Occupational Exposure and Risk Assessment

Registered propargite end-use products include emulsifiable concentrate (EC) and wettable powder (WP) formulations. Depending on the crops, these formulations may be applied as broadcast, banded or directed spray or chemigation foliar treatments pre- or postharvest using ground or aerial equipment. Single application rates range from 0.8 to 4.5 pounds per acre. Preharvest intervals range from 7 to 60 days. EPA has determined that there are potential exposures to mixers, loaders, applicators, or other handlers during usual use-patterns associated with propargite. Based on the use patterns and potential exposures described above, 14 major agricultural exposure scenarios are identified in this document to represent the extent of propargite uses.

Short-term handler exposure scenarios resulted in risk estimates expressed as MOEs, ranging from less than 1 to 2,570. A total of 68 exposure scenarios were evaluated for the various application rates assessed in each scenario. Based on the maximum level of protection (e.g.,

various levels of PPE or engineering controls) all scenarios had MOEs estimated to be greater than 100.

The results of the **intermediate-term handler** assessments indicate that all potential exposure scenarios provide at least one application rate with a total MOE(s) greater than or equal to 100 at either the **maximum PPE** (i.e., long pants, long sleeved shirts, and chemical resistant gloves while using open systems) or using **engineering controls** (i.e., closed systems). In the majority of cases, it is dermal exposure rather than the inhalation exposure contributing most to the exposure estimate (dermal and inhalation exposures were not combined). More specifically, the MOEs for all the scenarios range from 1 to 2,000. In total, 68 MOEs were calculated for the various application rates. Based on the maximum level of protection (i.e., various levels of PPE or engineering controls) all MOEs are greater than 100.

The baseline cancer risk estimates for handlers ranged from 3.2E-3 to 2.7E-6. When engineering controls were added the cancer risk was mitigated to 6.1E-5 to 6.9E-8.

For occupational **postapplication** exposure, propargite exposure estimates have MOEs equal to or exceeding 100 for all scenarios. Current propargite labels allow reentry in 48 hours. Field worker experience and reported incident data suggest that the skin irritation of propargite can be severe for several days after treatment. Longer REIs established in this document would help reduce incidents. This has been demonstrated in California when they extended the REIs in 1991 for various agricultural crops.

Residential Exposure and Risk Assessment

There are no registered residential uses of propargite.

This assessment reflects the Agency's current approaches for completing residential exposure assessments based on the guidance provided in the *Draft: Series 875-Occupational and Residential Exposure Test Guidelines, Group B-Postapplication Exposure Monitoring Test Guidelines (7/24/97 Version)*, the *Draft: Standard Operating Procedures (SOPs) for Residential Exposure Assessment (12/11/97 Version)*, and the *Overview of Issues Related to the Standard Operating Procedures for Residential Exposure Assessment* presented at the September 1999 meeting of the FIFRA Scientific Advisory Panel (SAP). The Agency is, however, currently in the process of revising its guidance for completing these types of assessments. Modifications to this assessment shall be incorporated as updated guidance becomes available and it is feasible from a regulatory perspective. This will include expanding the scope of the residential exposure assessments by developing guidance for characterizing exposures from other sources already not addressed such as from spray drift; residential residue track-in; exposures to farmworker children; and exposures to children in schools.

Aggregate Risk Assessment

There are no registered residential uses of propargite, so aggregation would contain only food and water risk estimates.

Acute aggregate risk estimates do not exceed HED's level of concern. The estimated environmental concentrations (EECs) for surface water (PRZM-EXAMS) were less than the acute DWLOCs, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The acute DWLOC for Females 13-50 years is 2400 ppb. The EECs for groundwater (SCI-GROW) were less than the acute DWLOC's, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The PRZM-EXAMS EEC was 34 ppb, while the estimated groundwater EEC was 0.006 ppb.

Chronic aggregate risk estimates do not exceed HED's level of concern. The EECs for surface water (PRZM-EXAMS) were less than the chronic DWLOCs, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the chronic DWLOC's, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The PRZM-EXAMS and SCI-GROW EECs were 8.7 and 0.006 ppb, respectively.

A cancer DWLOC was calculated based on the cancer dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were greater than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is greater than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). The PRZM-EXAMS and SCI-GROW EECs were 4.8 and 0.006 ppb, respectively. Surface water concentrations below 1.0 ppb would result in cancer risks below 1×10^{-6} for drinking water alone when back calculated. Time weighted average propargite concentration in surface water samples from the USGS NAWQA (Oristimba Creek Watershed) for the years 1992-1993 were 0.30 and 1.24 ppb, respectively. Therefore, even when monitoring data are used cancer exposure to propargite from surface water sources is greater than HED's level of concern.

cc : Chem F, Chron F. Morton


RDI:Team (1/6/00); RARC (2/1/00); SVH:9/13/01

TM, Thurston Morton, Rm. 816D CM2, 305-6691, mail code 7509C

2.0 PHYSICAL/CHEMICAL PROPERTIES CHARACTERIZATION

DESCRIPTION OF CHEMICAL

Propargite [2-(p-tert-butylphenoxy)cyclohexyl-2-propynyl sulfite] is a non-systemic organosulfite acaricide.



| | |
|--------------------|--|
| Trade Name: | Omite, Comite |
| Empirical Formula: | C ₁₉ H ₂₆ O ₄ S |
| Molecular Weight: | 350.5 g/mole |
| CAS Registry No.: | 2312-35-8 |
| PC Code: | 097601 |

IDENTIFICATION OF ACTIVE INGREDIENT

Propargite technical is a light to dark brown viscous liquid which decomposes (~200° C) before boiling, has a specific gravity of 1.10 at 20° C, octanol/water partition coefficient (log K_{ow}) of 5.8 at 25° C, and vapor pressure of 4.49 x 10⁻⁹ mm Hg at 25° C. Propargite is only slightly soluble in water (1.9 ppm at 25° C), but is soluble in most organic solvents (>200 g/L in acetone, dichloromethane, hexane, methanol, and toluene).

3.0 HAZARD CHARACTERIZATION

Table 1. Summary of Toxicology Endpoint Selection.

| EXPOSURE SCENARIO | DOSE (mg/kg/day) | ENDPOINT | STUDY TYPE/ MRID |
|--|--|--|---|
| Acute Dietary- females 13-50 | NOAEL= 8 UF = 100 FQPA = 1 | Increased incidence of fused sternebrae in fetuses at 10 mg/kg/day (LOAEL). | Developmental Toxicity in Rabbits 41336301 |
| Acute Dietary- general population | NOAEL= N/A UF = N/A FQPA = 1 | No relevant single exposure endpoint was identified. | N/A |
| | | Acute RfD (females 13-50) = 0.08 mg/kg/day Acute RfD (Gen. Pop.) = N/A aPAD = 0.08 mg/kg/day | |
| Chronic Dietary | NOAEL = 4 UF = 100 FQPA = 1 | Decreased body weight/body weight gain and increased mortality at 19 mg/kg/day(LOAEL) for males. | Chronic Feeding and Carcinogenicity in Rats 41750901 |
| | | Chronic RfD = 0.04 mg/kg/day cPAD = 0.04 mg/kg/day | |
| Cancer Risk | $Q_1^* = 0.33 \times 10^{-1} \text{ (mg/kg/day)}^{-1}$ | | |
| Short-Term ¹ (Dermal) | NOAEL= 6 MOE = 100 | Decreased maternal body weight gain at 8 mg/kg/day (LOAEL). | Developmental Toxicity in Rabbits 41336301 |
| Intermediate-Term ¹ (Dermal) | NOAEL= 4 MOE = 100 | Reduction in body weight gain and food consumption at 20 mg/kg/day (parental LOAEL). | Reproductive Toxicity in Rats 41352401 |
| Long-Term ¹ (Dermal) | NOAEL= 4 MOE = 100 | Decreased body weight / body weight gain and increased mortality at 20 mg/kg/day (LOAEL). | Chronic Feeding and Carcinogenicity in Rats 41750901 |
| Short Term ² (Inhalation) | LOAEL= 0.31mg/L or 50 mg/kg MOE = 1000 | Increased mortality at 0.31 mg/L (LOAEL) in males. | Acute Inhalation in Rats 42857003 |
| Intermediate Term ² (Inhalation) | | | |
| Long Term ² (Inhalation) | | | |

¹ A 14% dermal absorption factor will be used for risk assessment and an MOE of 100.

² An MOE of 1000 was selected for inhalation, including a 10X factor due to lack of a NOAEL, severity of effects at the lowest dose tested, and 4 hour duration.

Propargite is considered **corrosive** and has been placed in Category I for **both eye and dermal irritation** in rabbits. There have also been documented reports of dermal and eye irritation developing in workers exposed to propargite in the field. Evidence for its dermal sensitization potential have been noted; a study that provides conclusive results has not been possible due to the irritating properties of this chemical.

In a rabbit developmental toxicity study, propargite formulated as Omite (85% a.i.) was administered in corn oil by gavage to New Zealand White rabbits, 25 per dose, at levels of 0, 2, 4, 6, 8, or 10 mg/kg/day on gestation days (GD) 7-19. A reduction in maternal body weight gain occurred at doses of 8 and 10 mg/kg/day during GD 7-20 (gain of 9 g and loss of 20 g, respectively, versus a gain of 114, 165 and 119 g for control, 2 and 4 mg/kg/day, respectively). Only the incidence of fetuses with fused sternebrae at 10 mg/kg/day was considered to be significantly greater than that observed in concurrent and historical controls. The maternal LOAEL is 8 mg/kg/day, based on decreased body weight gain. The maternal NOAEL is 6 mg/kg/day. The developmental LOAEL is 10 mg/kg/day, based on increased incidence of fused sternebrae. The developmental NOAEL is 8 mg/kg/day.

In a chronic toxicity/carcinogenicity study, propargite formulated as Omite (87.2%, a.i.) was administered to 50 Sprague-Dawley Crl:CD BR rats/sex/dose (an additional 10 rats/sex/dose were sacrifice at 53 weeks) in 0.5% corn oil in the diet at dose levels of 0, 50, 80, 400 and 800 ppm (0, 2, 4, 19 and 39 mg/kg/day for males and 0, 3, 5, 24 and 49 mg/kg/day for females) for 24 months. Mortality for males (8/50 and 20/50 at 400 and 800 ppm, respectively) and for females (7/50 at 800 ppm) appeared to be related to the increased incidence of undifferentiated sarcoma in the GI tract. There were dose-related increases in incidence of jejunum tumors in both sexes. The incidences were 0, 0, 0, 10 and 15 tumors (0, 0, 0, 17% and 25%) in males and 0, 1, 0, 1, and 9 tumors (0, 2%, 0, 2% and 15%) in females for the control, 50, 80, 400, and 800 ppm dose groups, respectively (60 animals per group). They were not always associated with any increase in ulceration or other signs of irritation of the stomach or jejunum. Tumors of the jejunum were seen in males and females receiving the highest doses of 400 and 800 ppm. The dosing was considered to be adequate to assess the carcinogenic potential of propargite. The LOAEL is 400 ppm (19 mg/kg/day) for males due to increased mortality, decreased body weight and body weight gain, as well as decreases in total protein and calcium. The NOAEL is 80 ppm (4 mg/kg/day) for males. The LOAEL is 800 ppm (39 mg/kg/day) for females due to decreases in body weight and body weight gain. The NOAEL is 400 ppm (24 mg/kg/day) for females.

Propargite did not cause reproductive effects in rats but produced decreased parental and pup body weights. In a two-generation reproduction study, Omite (87.2 % a.i.) was administered to 25 Crl:CDBR rats/sex/dose in their diet at dose levels of 0, 80, 400, and 800 ppm (0, 4, 20, and 40 mg/kg/day) for 10 weeks then mated to produce the F_{1a} generation. They were mated a second time after a 2-week rest period to produce the F_{1b} generation. The F_{1b} generation were treated in a similar manner to produce the F_{2a} and F_{2b} generation. No compound-related clinical signs or reactions were observed for either parental group. A transient decrease in body weight gain occurred for animals in the high-dose and mid-dose groups. Both food consumption and food efficiency were reduced at 400 and 800 ppm. Necropsy revealed no compound related effects on gross or microscopic histological findings. There were no compound related adverse effects on the reproductive performance of any group. At the high-dose, there were decreases in mean pup weight at birth and during the period of lactation. The systemic LOAEL is 400 ppm (20 mg/kg/day), based on decreased parental body weight gain, and food consumption. The systemic NOAEL is 80 ppm (4 mg/kg/day). The offspring LOAEL is 800 ppm (40 mg/kg/day),

based on reduction of pup weight during lactation. The offspring NOAEL is 400 ppm (20 mg/kg/day). The reproductive LOAEL and NOAEL are > 800 ppm (40 mg/kg/day).

In an acute inhalation toxicity study (MRID 42857003), CD CrI:CD BR rats (5/sex/dose) were exposed by inhalation route (nose only) to propargite (85%, a.i.) at concentrations of 0.31, 0.80, and 1.3 mg/L for 4 hours. Mortality at the lowest level was observed within 24 hours of exposure (1/10). At the 0.80 mg/L dose, deaths occurred on day 2 and 3 (2/10). At the 0.31 mg/L dose, animals recovered and were sacrificed at day 15, the animals at 0.80 mg/L were observed an additional week and showed incomplete recovery. At 1.3 mg/L, all animals (10/10) died between days 2 and 17. Signs of toxicity included labored respiration, decreased activity, nasal discharge, anogenital staining, matted coats, at all levels. The animals at 0.80 and 1.3 mg/L showed moist rales, grasping, perioral encrustation as well. Weight loss was observed in all animals; however, the survivors exceeded their pretest weights at termination. Necropsy revealed discoloration of the lungs. Some showed signs of gastrointestinal distress and discoloration of the skin.

3.2 FQPA Considerations

The FQPA Safety Factor Committee met on August 9, 1999 to evaluate hazard and exposure data for propargite and recommend application of the FQPA Safety Factor (as required by Food Quality Protection Act of August 3, 1996), to ensure the protection of infants and children from exposure to propargite. Based on the lack of increased susceptibility following *in utero* exposure to rats and rabbits and pre/post natal exposure to rats, adequacy of the database, no currently registered residential uses, and because the exposure assessments will not underestimate the potential dietary (food and drinking water) exposures for infants and children from the use of propargite, the FQPA committee recommended reduction of the FQPA Safety Factor to 1X.

3.3 Endocrine Disruption

EPA is required under the FFDCA, as amended by FQPA, to develop a screening program to determine whether certain substances (including all pesticide active and other ingredients) "may have an effect in humans that is similar to an effect produced by a naturally occurring estrogen, or other such endocrine effects as the Administrator may designate." Following the recommendations of its Endocrine Disruptor Screening and Testing Advisory Committee (EDSTAC), EPA determined that there was scientific bases for including, as part of the program, the androgen and thyroid hormone systems, in addition to the estrogen hormone system. EPA also adopted EDSTAC's recommendation that the Program include evaluations of potential effects in wildlife. For pesticide chemicals, EPA will use FIFRA and, to the extent that effects in wildlife may help determine whether a substance may have an effect in humans, FFDCA authority to require the wildlife evaluations. As the science develops and resources allow, screening of additional hormone systems may be added to the Endocrine Disruption Screening Program (EDSP).

4.0 EXPOSURE ASSESSMENT

4.1 Summary of Registered Uses

Propargite is currently registered for ornamentals and a variety of field, fruit, and vegetable crops. Registered propargite end-use products include emulsifiable concentrate (EC) and wettable powder (WP) formulations. Depending on the crops, these formulations may be applied as broadcast, banded or directed spray or chemigation foliar treatments pre- or postharvest using ground or aerial equipment. Single application rates range from 0.8 to 4.5 pounds per acre. Preharvest intervals range from 7 to 60 days.

Manufacturing-Use Products

A search of the Reference Files System (REFS) conducted 12/16/99 identified a single propargite manufacturing-use product (MP) registered under PC Code 097601: the Uniroyal Chemical Company Inc. 90.6% T (EPA Reg. No. 400-95). Only the 90.6% T is subject to a reregistration eligibility decision.

According to a REFS search, conducted on 12/16/99, there are seven active end-use products (EPs) registered under FIFRA Section 3. These EPs, including the associated Special Local Need (SLN) registrations under FIFRA Section 24(c), are listed in Table 2. For the purpose of generating this Reregistration Eligibility Document (RED), the Agency examined the registered food/feed use patterns and reevaluated the available residue chemistry database for adequacy in supporting these use patterns, based on the product labels registered to Uniroyal Chemical Company.

Table 2. Propargite EPs with Food/Feed Uses Registered to Uniroyal Chemical Company

| EPA Reg. No. | Label Acceptance Date ¹ | Formulation | Product Name |
|----------------------|------------------------------------|----------------|--|
| 400-82 ² | 7/9/99 | 32% WP | Omite® - 30W Agricultural Miticide |
| 400-89 | 5/28/98 | 6 lb/gal EC | Omite® - 6E Agricultural Miticide |
| 400-104 ³ | 1/14/98 (3/98 in REFs) | 6.55 lb/gal EC | Comite® Agricultural Miticide |
| 400-154 ⁴ | 1/14/98 | 6 lb/gal EC | Comite® II Agricultural Miticide |
| 400-425 ⁵ | 5/27/98 | 32% WP | Omite® - CR Agricultural Miticide (For California Only) |
| 400-426 ⁵ | 5/28/98 | 32% WP | Omite® - CR Agricultural Miticide (Not For California) |
| 400-427 | 7/9/99 | 32% WP | Omite® - 30WS Agricultural Miticide |

¹ Date of the most recently EPA-approved label found by reviewer in the product jacket or Pesticide Product Label System (PPLS) unless specified otherwise.

² Including SLN No. CA810088 (avocado) and CA860070 (orange, grapefruit).

³ Including SLN Nos. AL910005, AR830015, AZ810022, AZ970004, CA780167, CA820083, CA8300024, CA920011, CA940031, GA910003, ID770005, ID910015, ID940011, ID960016, ID970015, IN990002, MS830024, MT890010, MT900001, NC910007, NV870009, NV880007, OR770013, OR790034, OR910019, OR940012, OR940013, OR970012, SC910003, TX830028, UT790015, UT960006, VA910006, WA770012, WA870029, WA890020, WA910033, WA970010, WI990016, and WY960001.

⁴ Including SLN Nos. CO940006, KS950001, NM940001, TX940005, and TX940006.

⁵ Including SLN Nos. ID950014, OR940021, and WA940007.

4.2 Food Exposure

The directions for use that were considered in the risk assessment are indicated in the Chemistry Chapter (J. Stokes, 1/20/00, D250257). Label revisions are needed. In addition, per Table 1 of OPPTS GLN 860.1000, the feeding restriction on cotton trash (presumably meaning cotton gin byproducts) is not practical. Once adequate residue data are submitted for cotton gin byproducts and a tolerance established, this feeding restriction should be deleted from the label.

The nature of the residue in plants and animals is adequately understood. The HED Metabolism Assessment Review Committee (MARC) concluded the residue of concern in plants and animals is propargite *per se* but required additional data on the metabolism of the propynyl sulfite side chain (N. Dodd, D256182, 6/7/99). The registrant submitted additional data on the metabolism of the propynyl sulfite side chain in rats and this study was classified as acceptable and propynyl sulfite metabolites should not be included as residues of concern (S. Shallal, D259994, 11/4/99). Approximately 56-65 % of the administered dose was eliminated via urine and/or feces within the first 24 hours. Only 2-2.6% of the administered dose was recovered from the carcasses of rats and mice; individual organs were, therefore, not analyzed for radioactivity. Six major metabolites were isolated from rat urine. The proposed metabolic pathway suggests that

following the cleavage of the 2-propynyl sulfite side chain of the propargite molecule, it is further detoxified via glutathione conjugation with further degradation leading to formation of the major metabolites.

Analytical methods available for enforcing propargite tolerances include Methods II, V, and VI for plant commodities and Methods III and IV for animal commodities in PAM, Volume II (Sec. 180.259). The preferred enforcement analytical method for plant commodities is Method V. All are gas liquid chromatography (GLC) methods with either sulfur-specific microcoulometric detection (Method II), microcoulometric detection (Method III), or flame photometric detection (Methods IV, V, and VI). Limits of quantitation are 0.08 (milk) and 0.1 ppm (plant and animal commodities).

In frozen storage, propargite is stable in/on avocados for 422 days, corn for 366 days, strawberries for 236 days, dried hops, apples, oranges, and sorghum grain for one year, and in plucked tea leaves, dried green tea, and dried black tea for 259 days. Additional confirmatory storage stability data are required for an oily commodity to support residue studies on peanut and walnut, and storage stability data are required to support corn and peanut processing studies. Propargite is stable in frozen storage for 90 days in milk, beef liver and beef fat, eggs, and chicken fat, and 180 days in beef kidney. Residues were stable for 30 days in beef muscle and declined by 17% after 90 days and 39% after 180 days.

Adequate field trials are available pending submission of required storage stability data, sample storage information, or required label amendments. Data on oranges indicate that residues up to 8.3 ppm may occur from registered use and that the 5 ppm tolerance is inadequate. This tolerance has been reassessed at 10 ppm. In sorghum grain, maximum propargite residues were 3.8 ppm, supporting a decrease in the current 10 ppm tolerance. Although one sample of cottonseed showed a residue of 0.11 ppm, based on the residue data for other samples after treatment at higher rates, HED considers the existing 0.1 ppm tolerance adequate to cover the current label use. This 0.1 ppm tolerance is in harmony with Codex. For all other crops the residue data support the established tolerances. Additional field trials are needed on cotton to determine a tolerance for propargite residues in/on cotton gin byproducts.

Adequate processing studies have been submitted for potatoes, citrus, field corn, grapes and peanuts. Storage stability data are required to support the corn and peanut processing studies. The corn processing study indicated that a tolerance is required for residues in aspirated grain fractions. The citrus processing study did not detect residue concentration in dried pulp, indicating that the current 40 ppm tolerance should be revoked. Residues concentrated in orange oil by 7x; based on a HAFT of 4 ppm (residue range 1.6-8.3 ppm; n=6) in oranges, a tolerance of 30 ppm is required. Although residues concentrated in raisins by 1.7x, this factor applied to the HAFT of 4.7 ppm yields a concentration in raisins of 8 ppm, which is lower than the 10 ppm tolerance for residues in/on the RAC. Therefore, a tolerance for raisins is not required.

The reregistration requirements for animal feeding studies are fulfilled. Acceptable ruminant and poultry feeding studies have been submitted and evaluated. In cows dosed with propargite at 50 ppm (approximately 2x) residues of propargite *per se* were <0.01-0.011 ppm in milk, 0.086-0.2 ppm in fat, and <0.01-0.02 ppm in liver, muscle, and kidney. In a poultry feeding study, propargite residues were <0.01 ppm (nondetectable) in eggs from hens dosed at 5, 15, or 50 ppm (1, 3, and 10x). Propargite residues in fat were <0.01 ppm in hens dosed at 5 ppm and 0.013-0.082 ppm in from hens dosed at 15 or 50 ppm. Propargite was not analyzed in tissues. In the poultry metabolism study, the parent compound was not detected in muscle, liver, or kidney.

The metabolism of propargite in rotated crops is similar to that in primary crops. Based on adequate confined and limited field rotational crop studies, the Agency concluded that a six-month plantback interval (PBI) for root crops and a two-month PBI for all other crops are acceptable (J. Stokes, 5/31/00, D230867).

4.2.1 Tolerance Reassessment Summary

Effective 10/19/99 EPA has revoked the following tolerances: propargite residues in/on apples, apricots, succulent beans, cranberries, figs, peaches, pears, plums, and strawberries [established under §180.259(a)] and dried figs (§186.5000) [FR 64 39068-39072, 7/21/99]. Uses of propargite on these crops have been canceled for over 3 years. The final rule will remove §186.5000, transferring the tolerances for residues in hops, dried and tea, dried to §180.259.

4.2.2 Codex Harmonization

The U.S. tolerances for propargite residues and Codex MRLs are identical with respect to the residue regulated; both are defined as the parent compound. A numerical comparison of the Codex MRLs and the corresponding reassessed U.S. tolerances is presented in the propargite product and residue chemistry chapter.

4.2.3 Dietary Exposure Reassessment

Consumption Data

HED conducts dietary risk assessments using the Dietary Exposure Evaluation Model (DEEM™), which incorporates consumption data generated in USDA's Continuing Surveys of Food Intakes by Individuals (CSFII), 1989-1992. For acute dietary risk assessments, the entire distribution of single day food consumption events is combined with either a single residue level (deterministic analysis, risk at 95th percentile of exposure reported) or a distribution of residues (probabilistic analysis, referred to as "Monte Carlo," risk at 99.9th percentile of exposure reported) to obtain a distribution of exposure in mg/kg/day. For chronic dietary risk assessments, the three-day average of consumption for each sub-population is combined with residues in commodities to determine average exposure in mg/kg/day.

Propargite Residue Data

Revised anticipated residues (ARs) (T. Morton, D266001, 5/24/00) were calculated and used in the revised dietary exposure analyses. The Biological and Economic Analysis Division (OPP/BEAD) has provided usage information for propargite (Jihad Alsadek, 5/22/00). Field trial data and USDA Pesticide Data Program (PDP) data were used in calculation of ARs. For all PDP analyses the ½ Limit of Detection (LOD) value was a weighted average of all laboratory limits of detection.

4.2.4 Acute Dietary Exposure Assessment

Estimated acute dietary exposure is below HED’s level of concern for the U.S. Population and all subpopulations at the 99.9th percentile. Use of PDP monitoring data, field trial data, and calculated livestock ARs results in a risk estimate of 2 % of the acute PAD (% aPAD) for the subpopulation Females (13-50 years) at the 99.9th percentile.

Table 3. Population Adjusted Dose Acute Dietary Exposure Results for Propargite (PAD = 0.08 mg/kg/day).

| Subgroups | 95 th Percentile Exposure mg/kg/day (%aPAD) | 99 th Percentile Exposure mg/kg/day (%aPAD) | 99.9 th Percentile Exposure mg/kg/day (%aPAD) |
|---------------------|---|---|---|
| Females (13-50 yrs) | 0.00005 (<1%) | 0.0001 (<1%) | 0.001 (2%) |

4.2.5 Chronic Dietary Exposure Assessment

Estimated chronic dietary exposure is below HED’s level of concern. Use of PDP monitoring data, field trial data, and calculated livestock ARs results in a maximum risk of <1 % of the chronic PAD (% cPAD) for the U.S. Population and all subpopulations.

Table 4. Propargite Chronic Dietary Exposure/Risk.

| Population Subgroup | Chronic | |
|-----------------------|-------------------------|--------|
| | Exposure (mg/kg/day) | % cPAD |
| U.S. Population | 0.00001 | <1 |
| All infants (<1 yr) | 0.00001 | <1 |
| Children (1-6 yrs) | 0.00001 | <1 |
| Children (7-12 yrs) | 0.00001 | <1 |
| Females (13-50 years) | 0.00001 | <1 |
| Males (20+ yrs) | 0.00001 | <1 |

4.2.6 Cancer Dietary Exposure Assessment

The cancer dietary risk estimate for propargite is 1.8×10^{-7} . Results of the analyses indicate potential residues in milk contribute to estimated exposure and risk. There were no detections of propargite in PDP data for milk and the highest residue in the 2x feeding study was at the 0.011 ppm. Therefore, a sensitivity analysis was performed by inserting zeroes for the milk commodities resulting in a cancer dietary risk estimate of 1.6×10^{-7} .

Table 5. Propargite Cancer Dietary Exposure/Risk.

| Population Subgroup | Cancer | | Cancer (with zeroes inserted for milk commodities) | |
|---------------------|-------------------------|-------------------------------------|---|-------------------------------------|
| | Exposure (mg/kg/day) | Lifetime Risk ($Q_1^* - .033$) | Exposure (mg/kg/day) | Lifetime Risk ($Q_1^* - .033$) |
| U.S. Population | 0.000006 | 1.8×10^{-7} | 0.000005 | 1.6×10^{-7} |

4.2.7 Drinking Water Exposure

Environmental Fate and Effects Division (EFED) provided HED with estimated environmental concentrations (EECs) for propargite in surface water and groundwater. EFED model estimates include two applications of propargite. PRZM-EXAMS and SCI-GROW EECs are as follows:

Table 6. PRZM-EXAMS and SCI-GROW EECs (ug/L) for propargite use.

| Model | EECs |
|----------------------------|---|
| Surface Water (PRZM-EXAMS) | Cotton 1 in 10 year Peak Concentration = 34.4 ppb 1 in 10 year Annual Mean Concentration = 8.7 ppb 36-year Annual Mean Concentration = 4.8 ppb |
| Groundwater (SCI-GROW) | 0.006 ppb |

In addition, EFED provided time weighted averages for propargite concentrations in surface water samples from the USGS NAWQA (Oristimba Creek Watershed) for the years 1992 and 1993 were 0.30 and 1.24 ppb, respectively.

4.2.7.1 DWLOCs for Acute Exposure

Acute DWLOCs were calculated based on the acute dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were less than the acute DWLOCs, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The acute DWLOC for Females 13-50 years is 2300 ppb. The PRZM-EXAMS surface water value is 34 ppb.

The EECs for groundwater (SCI-GROW) were less than the acute DWLOC's, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). To calculate the DWLOC, the acute dietary food exposure was subtracted from the acute PAD using the equation:

$$DWLOC_{acute} (ug/L) = \frac{[acute\ water\ exposure\ (mg/kg/day) \times (body\ weight\ in\ kg)]}{[consumption\ (L/day) \times 10^{-3}\ mg/\mu g]}$$

where acute water exposure (mg/kg/day) = [aPAD - (acute food (mg/kg/day))]

Table 7. Drinking Water Levels of Comparison for Acute Dietary Exposure.

| Population Subgroup | Acute PAD (mg/kg/day) | Food Exposure (mg/kg/day) | Max Water Exposure (mg/kg/day) | DWLOC _{acute} (ug/L) | PRZM-EXAMS (ug/L) | SCI-GROW (ug/L) |
|---------------------|-----------------------|---------------------------|--------------------------------|-------------------------------|-------------------|-----------------|
| Females 13-50 yrs. | 0.08 | 0.001 | 0.08 | 2400 | 34 | 0.006 |

4.2.7.2 DWLOCs for Chronic Exposure

Chronic DWLOCs were calculated based on the chronic dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were less than the chronic DWLOCs, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the chronic DWLOCs, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). To calculate the chronic DWLOC, the chronic dietary food exposure was subtracted from the chronic PAD using the equation:

$$DWLOC_{chronic} (ug/L) = \frac{[chronic\ water\ exposure\ (mg/kg/day) \times (body\ weight\ in\ kg)]}{[consumption\ (L/day) \times 10^{-3}\ mg/\mu g]}$$

where chronic water exposure (mg/kg/day) = [cPAD - (chronic food (mg/kg/day))]

Table 8. Drinking Water Levels of Comparison for Chronic Dietary Exposure.

| Population Subgroup | Chronic PAD (mg/kg/day) | Food Exposure (mg/kg/day) | Max. Water Exposure (mg/kg/day) | DWLOC _{chronic} (ug/L) | GENEEC (ug/L) | SCI-GROW (ug/L) |
|---------------------|-------------------------|---------------------------|---------------------------------|---------------------------------|---------------|-----------------|
| US Population | 0.04 | 0.00001 | 0.04 | 1400 | 8.7 | 0.006 |
| All Infants | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Children 1-6 | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Children 7-12 | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Females 13-50 yrs. | 0.04 | 0.00001 | 0.04 | 1200 | 8.7 | 0.006 |
| Males 20+ yrs | 0.04 | 0.00001 | 0.04 | 1400 | 8.7 | 0.006 |

4.2.7.3 DWLOCs for Cancer Exposure

A cancer DWLOC was calculated based on the cancer dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were greater than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is greater than HED’s level of concern. The EECs for groundwater (SCI-GROW) were less than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is less than HED’s level of concern. The Agency’s default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). The PRZM-EXAMS and SCI-GROW EECs were 4.8 and 0.006 ppb, respectively. Surface water concentrations below 1.0 ppb would result in cancer risks below 1×10^{-6} for drinking water alone when back calculated. Time weighted average propargite concentration in surface water samples from the USGS NAWQA (Oristimba Creek Watershed) for the years 1992-1993 were 0.30 and 1.24 ppb, respectively. Therefore, even when monitoring data are used cancer exposure to propargite from surface water sources is greater than HED’s level of concern.

Table 9. Summary of Cancer DWLOC Calculations for Propargite.

| Population | Q _r * | Negligible Risk Level ¹ | Target Max Exposure ² mg/kg/day | Chronic Food Exposure mg/kg/day | Max Water Exposure ³ mg/kg/day | Groundwater EEC (ug/l) | Surface Water EEC (ug/L) | Cancer DWLOC ⁴ (ug/L) |
|------------|-----------------------|------------------------------------|--|---------------------------------|---|------------------------|--------------------------|----------------------------------|
| U.S. Pop. | 0.33x10 ⁻¹ | 0.000001 | 0.0000303 | 0.00001 | 0.0000203 | 0.006 | 4.8 | 0.71 |

¹ DWLOC_{CANCER} was calculated for U.S. population only. Default body weight and consumption value for calculation of the DWLOC were: 2L/70 kg

² Target Maximum Exposure (mg/kg/day) = [negligible risk/Q*]

³ Maximum Water Exposure (mg/kg/day) = [Target Maximum Exposure - (Chronic Food Exposure + Residential Exposure (Lifetime Average Daily Dose))]

⁴ Cancer DWLOC(ug/L) = $\frac{[\text{maximum water exposure (mg/kg/day)} \times \text{body weight (kg)}]}{[\text{water consumption (L)} \times 10^{-3} \text{ mg/}\mu\text{g}]^2}$

4.3 Non-Dietary Exposure

4.3.1 Occupational Handler Exposure Scenarios

EPA has determined that there are potential exposures to mixers, loaders, applicators, or other handlers during usual use-patterns associated with propargite. Based on the use patterns and potential exposures described above, 14 major agricultural exposure scenarios are identified in this document to represent the extent of propargite uses.

Agricultural exposure scenarios include: (1a) mixing/loading liquids for aerial application, (1b) mixing/loading liquids for chemigation, (1c) mixing/loading liquids for groundboom application, (1d) mixing/loading liquids for orchard airblast sprayer application, (1e) mixing/loading liquids for application of high pressure handwand, (2a) mixing/loading wettable powder for aerial application, (2b) mixing/loading wettable powder for groundboom application, (2c) mixing/loading wettable powder for orchard airblast sprayer application, (2d) mixing/ loading wettable powder for application of high pressure handwand, (3) applying sprays with fixed-wing aircraft, (4) applying sprays using a groundboom sprayer, (5) applying sprays with an airblast sprayer, (6) applying liquids with a high pressure handwand and (7) flagging during aerial spray application.

In most cases, HED assesses the exposure and risk to mixer/loaders and applicators separately for tractor drawn applications (i.e., airblast, groundboom, and granular spreaders). This practice has evolved, not because it is believed that there are always separate job functions, but rather because of the limited amount of information regarding these practices along with limited exposure data.

HED has adopted a methodology to present the risks separately for some scenarios and combine others. Most of the hand-held equipment such as backpack sprayers, and push type granular spreaders are assessed as a combined function. With these types of small operations the mixing, loading, and applying are almost always carried out by the same individual and there are data available to estimate exposure from these activities. For equipment such as fixed-wing-aircraft, groundboom tractors, and airblast sprayers the applications are assessed separately from the individual who mixes and loads the formulated product. HED assumes that the pilots are rarely involved in the mixing/loading. By separating the two job functions, HED can determine the most appropriate PPE or engineering control without requiring the handler to wear PPE throughout the entire workday or engineering controls that are not needed.

The potential handler exposures are assessed using the toxicological endpoints and uncertainty factors associated with the active ingredient. Therefore, the PPE and engineering controls are determined by the assessment of the active ingredient and not the currently required risk mitigation measures on propargite labels. This distinction of determining risk mitigation measures based on the active ingredient instead of the label required PPE is also important because of the nature of the end-use products. For example, some end-use products require additional PPE that are not necessary for the active ingredient because of the end-use product's

potential for eye and/or skin irritation based on inerts. Conversely, the Agency does not want to mandate additional PPE (e.g., heat stress issues) if it is not required based on the endpoint and uncertainty factors. Baseline attire (long pants, long sleeved shirt, and no gloves) is not presented in this chapter because of the need for additional PPE and/or engineering controls for all scenarios. There are some PPE, such as chemical-resistant aprons and/or head gear, that the Agency uses as qualitative measures because there are no recognized protection factors (PF) to assess their effectiveness.

4.3.1.1 Occupational Handler Exposure Data Sources and Assumptions

Uniroyal submitted applicator exposure studies in support of the reregistration process for propargite. These studies include:

- Airblast applicator exposure studies (MRID Nos. 418486-05 and 420997-02)
- Groundboom applicator exposure study (MRID No. 418486-05)

It is HED's policy to combine chemical-specific studies with similar surrogate data from the Pesticide Handlers Exposure Database (PHED) to assess handler exposures for regulatory actions. In addition, the exposure estimates from PHED (V1.1) are used to assess exposure where no chemical specific data are available.

Table 10. Exposure Variables for Agricultural Uses of Propargite.

| Exposure Scenario (Scenario #) | Are Chemical Specific Monitoring Data Available? ^a | Application Rates (lb ai/acre) ^b | Daily Acres Treated ^c |
|--|---|--|----------------------------------|
| Mixer/Loader Exposure | | | |
| Mixing/Loading Liquids for Aerial Application (1a) | No | 1.5 min / 2.5 max carrot, sugar beet, potatoes, dry beans, mint, corn (field, pop, sweet), sorghum grain, alfalfa, clover, peanut, jojoba | 350 |
| | | 2.5 grapefruit, orange | 125 |
| | | 2.5 min/4.5 max almond, walnut | |
| | | 1.6 cotton | 350-1200 |
| | | hops max 2.5 | 80 |
| Mixing/Loading Liquids for Chemigation (1b) | No | 2.0 min /2.5 max Potatoes, corn (sweet) | 350 |
| Mixing/Loading Liquids for Groundboom Application (1c) | No | 1.5 min /2.5 max potatoes, corn (field, pop, sweet) sorghum grain, alfalfa, clover, cotton, peanut, jojoba and mint | 80 |
| | | 2.5 hops | 40 |
| Mixing/Loading Liquids for Airblast Sprayer Application (1d) | No | 1.5 quince, cherry, prunes, orange, grapefruit, lemon, lime, tangerine, boysenberry, current, raspberry, hops, date, persimmons, | 40 |

| Exposure Scenario (Scenario #) | Are Chemical Specific Monitoring Data Available?* | Application Rates (lb ai/acre) ^b | Daily Acres Treated ^c |
|--|---|--|----------------------------------|
| | | 2.5 Xmas tree plantations, conifers, shade trees | 40 |
| | | 1.5 min/max 3.0 almond, filbert, macadamia nut, pecan, pistachio | 40 |
| | | 4.5 walnut | 40 |
| Mixing/Loading Liquids for Application of High Pressure Handwand (1e) | No | 1.5 non-bearing nursery stock | 5 |
| Mixing/Loading Wettable Powder for Aerial Application (2a) | No | 3.0 nectarine | 125 |
| | | 4.0 walnut | 125 |
| Mixing/Loading Wettable Powder for Groundboom Application (2b) | No | 1.6 peanut | 80 |
| Mixing/ Loading Wettable Powder for Airblast Sprayer Application (2c) | No | 4.5 max grapefruit, orange, lemon, avocado | 40 |
| | | 3.0 grapes | 40 |
| Mixing/ Loading Wettable Powder for Application of High Pressure Handwand (2d) | No | 0.5 min / 2.5 max non-bearing nursery stock | 5 |
| Applicator Exposure | | | |
| Applying Sprays with Fixed-Wing Aircraft –Enclosed Cockpit (3) | No | 1.5 min / 2.5 max carrot, sugar beet, potatoes, dry beans, mint, corn (field, pop, sweet), sorghum grain, alfalfa, clover, | 350 |
| | | 2.5 grapefruit, orange | 125 |
| | | 2.5 min/4.5 max almond, walnut | |
| | | 1.5 peanut, jojoba | 125 |
| | | 1.6 cotton | 350-1200 |
| | | hops max 2.5 | 80 |
| Applying Sprays with a Groundboom Sprayer (4) | Yes 41848606 | 1.5 min / 2.5 max potatoes, corn (field, pop, sweet) sorghum grain, alfalfa, clover, cotton, peanut, jojoba and mint | 80 |
| Applying Sprays with an Airblast Sprayer (5) | Yes 41848605 42099702 | 1.5 min quince, cherry, prunes, orange, grapefruit, lemon, lime, tangerine, boysenberry, current, raspberry, hops, date, persimmons, | 40 |
| | | 2.5 Christmas tree plantations, conifers, shade trees | 40 |
| | | 1.5min / max 3.0 almond, filbert, macadamia nut, pecan, pistachio | 40 |
| | | 4.5 walnut | 40 |
| Applying Liquids with a High Pressure Handwand (6) | No | 0.5 min / 2.5max non-bearing nursery stock | 5 |

| Exposure Scenario (Scenario #) | Are Chemical Specific Monitoring Data Available? ^a | Application Rates (lb ai/acre) ^b | Daily Acres Treated ^c |
|--|---|--|----------------------------------|
| Flagger Exposure | | | |
| Flagging During Aerial Spray Application (7) | No | 1.5 min / 2.5 max carrot, sugar beet, potatoes, dry beans, mint, corn (field, pop, sweet), sorghum grain, alfalfa, clover, peanut, jojoba | 350 |
| | | 2.5 grapefruit, orange | 125 |
| | | 2.5 min/4.5 max almond, walnut | |
| | | 1.6 cotton | 350-1200 |
| | | 2.5 max hops | 80 |

^a Available chemical-specific passive dosimetry data have been combined with PHED (V1.1).

^b Application rates are the maximum or range found on EPA Reg. Nos. 400-82, 400-83, 400-89, 400-104, 400-154, 400-425, 400-426, 400-427.

^c Daily acres treated are based on HED's estimates of acreage that would be reasonably expected to be treated in a single day for each exposure scenario of concern.

The handler exposure assessments encompass all of the major uses of propargite throughout the country. It is difficult to assess all of the "typical" agricultural uses (i.e., actual or predominant application rates and farm sizes), and therefore, an assessment has been developed that is believed to be realistic and yet provides a reasonable certainty that the exposures are not underestimated. The assumptions and uncertainties are identified below to be used in risk management decisions:

- *Application Rates:* The application rates are the maximum allowable that were identified on the available product labels. A range of application rates were used when the maximum application rates for various crops varied widely. Application rates have been rounded off.
- *Amount Handled:* The daily acres treated are HED standard values (see Table 9). Deviations from these standard values include the aerial acreage for orchard fruits, tree nuts, and Christmas trees. The orchard acreage is assessed at 125 acres because fruit orchards are grown in smaller plots, and cotton field is assessed at 350 to 1200 acres.
- *Unit Exposures:* The unit exposure values calculated by PHED generally range from the geometric mean to the median of the selected data set. To add consistency and quality control to the values produced from this system, the PHED Task Force has evaluated all data within the system and has developed a set of grading criteria to characterize the quality of the original study data. The assessment of data quality is based on the number of observations and the available quality control data. These evaluation criteria and the caveats specific to each exposure scenario are summarized in Appendix A, Table A4 of the Occupational Exposure Chapter, D276513, S. Tadayon, 8/xx/01. While data from PHED provides the best available information on handler exposures, it should be noted that some aspects of the included studies (e.g., duration, acres treated, pounds of active ingredient handled) may not accurately represent labeled uses in all cases.

- BEAD provided data for both commercial applicator and private grower; therefore, calculations were performed for both, where applicable. Two exposure frequencies were used in the calculations, the first represented the maximum number of applications per site per season to represent private use (7), and the second frequency applied a factor of two to the first frequency to represent commercial handlers making multiple applications per site per season (14).

Handler exposure assessments are completed by EPA using a baseline exposure scenario and, if required, increasing levels of risk mitigation (PPE and engineering controls) to achieve a margin of exposure of 100 for dermal exposure and 1,000 for inhalation exposure or cancer risk of $1.0E-4$. Appendix A of the Occupational Exposure Chapter, D276513, S. Tadayon, 7/31/01 presents the short-term and intermediate term MOE calculations for baseline exposure plus the risk mitigation measures of personal protective equipment (PPE) and engineering controls using the passive dosimetry results from the chemical-specific studies combined with surrogate data from PHED for the agricultural uses of propargite. Table 10 presents the cancer risk calculations for baseline exposure plus the risk mitigation measures of personal protective equipment (PPE) and engineering controls.

EPA calculated the baseline MOE (short-term and intermediate-term) and cancer for each of the exposure scenarios using the following **baseline** PPE assumptions:

- all occupational handlers are wearing footwear (socks plus shoes or boots);
- occupational mixers and loaders using open mixing techniques are wearing long-sleeved shirts, long pants, and no gloves;
- occupational applicators who use open cab airblast or tractor-driven application equipment and handlers flagging for aerial applications are wearing long-sleeved shirts, long pants, and no gloves; and
- occupational handlers (mixers, loaders, and applicators) who use hand-held application equipment are wearing long-sleeve shirts, long pants, and no gloves.

If the baseline short-term or intermediate-term MOE calculated using this baseline PPE was 100 or greater (since the NOAEL is based on data from animal studies) for an exposure scenario, then no further calculations were made. If the baseline short-term or intermediate-term MOE was less than 100 for any exposure scenario, an additional short-term or intermediate-term MOE was calculated based on increasing the level of PPE over the baseline PPE. HED calculated the additional PPE short-term or intermediate-term MOE for each occupational exposure scenario with a baseline total MOE of less than 100, using the following additional **PPE** assumptions:

- all occupational handlers are wearing footwear (socks plus shoes or boots);

- occupational mixers and loaders using open mixing techniques are wearing long-sleeved shirts and long pants and gloves;
- occupational applicators who use open cab airblast or tractor-driven application equipment and handlers flagging for aerial applications are wearing (except flaggers- no gloves) long-sleeved shirts and long pants (coveralls and chemical resistant head gear for open cab airblast);
- Also, if necessary, a dust/mist mask represented by a 10-fold protection factor is added to mitigate the risks.

If the additional-PPE short-term or intermediate-term MOE calculated using this additional-PPE was 100 or greater (the NOAEL is based on data from animal studies) for an exposure scenario, then no further calculations were made. If the additional-PPE short-term or intermediate-term MOE remained less than 100 for any occupational exposure scenario, an additional short-term or intermediate-term MOE was calculated based on mandatory use of engineering controls where feasible. Engineering controls are not available for occupational handlers (mixers, loaders, and applicators) who use hand-held application equipment. HED calculated the engineering-control short-term or intermediate-term MOE for each occupational exposure scenario with an additional-PPE short-term or intermediate-term MOE of less than 100, using the following **engineering control** assumptions:

- all occupational handlers are wearing footwear (socks plus shoes or boots);
- occupational mixers and loaders handling liquid formulations using a closed system are wearing chemical-resistant gloves plus long-sleeved shirts and long pants;
- occupational mixers and loaders handling wettable powders using a closed system (water-soluble packages) are wearing long-sleeved shirts and long pants, and chemical-resistant gloves; and
- occupational applicators who use aerial, airblast, or tractor-driven application equipment and handlers flagging for aerial applications are located in enclosed cabs or cockpits and are wearing long-sleeved shirts and long pants, and no gloves.

4.3.1.2 Occupational Handler Risk Characterization

Table 10 summarizes the numeric MOE values for both the short- and intermediate-term exposure durations as well as cancer risk estimates. In the majority of cases, it is dermal exposure rather than the inhalation exposure contributing the most to the exposure estimate (dermal and inhalation exposures were not combined). The MOEs are presented for baseline, PPE and engineering controls. Cancer risk estimates are also summarized at different levels of mitigation. Baseline represents long pants, long-sleeved shirts and no gloves; PPE represents

exposure while wearing long pants, long-sleeved shirts and chemical resistant gloves, and an organic vapor respirator (10-fold protection factor) while using open mixing/loading systems and open cab tractors. The engineering controls represent exposure while wearing long pants, long-sleeved shirts and no gloves (except chemical resistant gloves for closed loading systems) while using closed mixing/loading systems and enclosed cabs/cockpits.

The results of the **short-term** exposure duration indicate that the MOEs range from less than 1 to 2,570. A total of 68 MOEs were calculated for the various application rates assessed in each scenario. Based on the maximum level of protection (e.g., various levels of PPE or engineering controls) all MOEs are greater than 100.

The results of the **intermediate-term** exposure duration indicate that the total MOEs range from 1 to 2,000. A total of 68 MOEs were calculated for the various application rates assessed in each scenario. Based on the maximum level of protection (e.g., various levels of PPE or engineering controls) all MOEs are greater than 100.

The baseline cancer risk estimates for handlers ranged from $3.2\text{E-}3$ to $2.7\text{E-}6$. When engineering controls were added the cancer risk was mitigated to $6.1\text{E-}5$ to $6.9\text{E-}8$.

Table 11. Exposure Variables, MOEs and Cancer for Agricultural use of Propargite.

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb ai/acre) | Daily Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | |
|--|-----------------------------|--|--------------------------------|---------------------|------------------------|-----|--------------|-------------------------------|-----|--------------|-----------------|-----|-------------------|-------------------|-------------------|-------------------|
| | | | | | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control |
| Mixer/Loader Exposure | | | | | | | | | | | | | | | | |
| Mixing/Loading Liquids for Aerial Application (1a) | Roots and Tuber Vegetable | carrot, sugar beet, potatoes, dry beans, mint | Min 2.0 | 350 | 1 | 160 | NA | <1 | 125 | NA | 4135 | NA | NA | 1.3E-3/ 2.7E-3 | 1.4E-5/ 2.8E-5 | 4.0E-6/ 8.0E-6 |
| | Legume Vegetable | | Max 2.5 | | 1 | 130 | NA | <1 | 100 | NA | 3305 | NA | NA | 1.6E-3/ 3.2E-3 | 1.7E-5/ 3.5E-5 | 5.0E-6/ 9.9E-6 |
| | Herbs & Spices | hops | Max 2.5 | 80 | 4 | 560 | NA | 3 | 435 | NA | 14465 | NA | NA | 3.6E-4/ 7.3E-4 | 4.0E-6/ 7.9E-5 | 1.2E-6/ 2.3E-6 |
| | Citrus Fruits | grapefruit, orange | Max 2.5 | 125 | 3 | 360 | NA | 2 | 280 | NA | 9260 | NA | NA | 5.6E-4/ 1.1E-3 | 6.3E-6/ 1.3E-5 | 1.8E-6/ 3.6E-6 |
| | Tree Nuts | almond, walnut | Min 2.5 | 125 | 3 | 360 | NA | 2 | 280 | NA | 9260 | NA | NA | 5.6E-4/ 1.1E-3 | 6.3E-6/ 1.3E-5 | 1.8E-6/ 3.6E-6 |
| | | | Max 4.5 | | 2 | 200 | NA | 1 | 155 | NA | 5145 | NA | NA | 1.0E-3/ 2.0E-3 | 1.1E-5/ 2.2E-5 | 3.3E-6/ 6.5E-6 |
| | Cereal Grains | corn (field, pop, sweet), sorghum grain, alfalfa, clover | Min 1.5 | 350 | 2 | 215 | NA | 1 | 165 | NA | 5510 | NA | NA | 9.6E-4/ 1.9E-3 | 1.1E-5/ 2.1E-5 | 3.1E-6/ 6.1E-6 |
| | Non-grass Animal Feed | | Max 2.5 | | 1 | 130 | NA | <1 | 100 | NA | 3305 | NA | NA | 1.6E-3/ 3.2E-3 | 1.7E-5/ 3.5E-5 | 5.0E-6/ 9.9E-6 |
| | Oil Seed | cotton | Max 1.6 | 350 | 2 | 200 | NA | 1 | 155 | NA | 5165 | NA | NA | 1.0E-3/ 2.0E-3 | 1.1E-5/ 2.2E-5 | 3.3E-6/ 6.5E-6 |
| | | | | | <1 | 60 | 155 | <1 | 45 | 120 | 1505 | NA | NA | 3.6E-4/ 7.3E-4 | 4.0E-5/ 7.9E-5 | 1.1E-5/ 2.2E-5 |
| | | peanut, jojoba | Min 1.5 | 350 | 2 | 215 | NA | 1 | 165 | NA | 5510 | NA | NA | 9.6E-4/ 1.9E-3 | 1.1E-5/ 2.1E-5 | 3.1E-6/ 6.1E-6 |
| | | | Max 2.5 | | 1 | 130 | NA | <1 | 100 | NA | 3305 | NA | NA | 1.6E-3/ 3.2E-3 | 1.7E-5/ 3.5E-5 | 5.0E-6/ 9.9E-5 |
| Ornamental plants | Christmas Tree conifer seed | Max 2.5 | 125 | 3 | 360 | NA | 2 | 280 | NA | 9260 | NA | NA | 5.6E-4/ 1.1E-3 | 6.3E-6/ 1.3E-5 | 1.8E-6/ 3.6E-6 | |
| Mixing/Loading Liquids for Chemigation (1b) | Roots and vegetable | potatoes, corn (sweet) | Min 2.0 | 350 | 1 | 160 | NA | 1 | 125 | NA | 4135 | NA | NA | 1.3E-3/ 2.6E-3 | 1.4E-5/ 2.8E-5 | 4.0E-6/ 7.9E-6 |
| | Cereal Grains | | Max 2.5 | | 1 | 130 | NA | <1 | 100 | NA | 3305 | NA | NA | 1.6E-3/ 3.2E-3 | 1.7E-5/ 3.5E-5 | 5.0E-6/ 9.9E-6 |

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb ai/acre) | Daily Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | | |
|---|---------------------------------|--|--------------------------------|---------------------|------------------------|------|--------------|-------------------------------|-------|--------------|-----------------|------|--------------|-------------------|-------------------|-------------------|--|
| | | | | | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. control | |
| Mixing/Loading Liquids for Groundboom Application (1c) | Roots and Vegetable | potatoes, corn (field, pop, sweet) sorghum grain, alfalfa, clover, cotton, peanut, jojoba and mint | Min 1.5 | 80 | 7 | 930 | NA | 6 | 725 | NA | 24110 | NA | NA | 2.2E-4/ 4.4E-4 | 2.4E-6/ 4.8E-6 | 6.9E-7/ 1.4E-6 | |
| | Cereal Grains | | | | | | | | | | | | | | | | |
| | Non-grass Animal Feed | | | | | | | | | | | | | | | | |
| | Oil Seed | | Max 2.5 | | 4 | 560 | NA | 3 | 435 | NA | 14465 | NA | NA | 3.6E-4/ 7.3E-4 | 4.0E-6/ 7.9E-6 | 1.2E-6/ 2.3E-6 | |
| | Herbs and Spices | | | | | | | | | | | | | | | | |
| Mixing/Loading Liquids for Airblast Sprayer Application (1d) | Pome Fruits | quince, cherry, prunes, orange, grapefruit lemon, lime, tangerine, boysenberry, current, date raspberry, hops, persimmons. | 1.5 | 40 | 15 | 1865 | NA | 10 | 1450 | NA | 48220 | NA | NA | 1.1E-4/ 2.2E-4 | 1.2E-6/ 2.4E-6 | 3.6E-7/ 7.3E-7 | |
| | Stone fruits | | | | | | | | | | | | | | | | |
| | Citrus Fruits | | | | | | | | | | | | | | | | |
| | Berries | | | | | | | | | | | | | | | | |
| | Herbs and Spices | | | | | | | | | | | | | | | | |
| | Tropical and Subtropical Fruits | | | | | | | | | | | | | | | | |
| | Tree Nuts | almond, filbert, macadamia nut, pecan, pistachio | Min 1.5 | | 15 | 1865 | NA | 10 | 1450 | NA | 48220 | NA | NA | 1.1E-4/ 2.2E-4 | 1.2E-6/ 2.4E-6 | 3.6E-7/ 7.3E-7 | |
| | | | Max 3.0 | | 7 | 930 | NA | 6 | 725 | NA | 24110 | NA | NA | 2.2E-4/ 4.4E-4 | 2.4E-6/ 4.8E-6 | 6.9E-7/ 1.4E-6 | |
| | | walnut | Max 4.5 | | 5 | 620 | NA | 4 | 485 | NA | 16075 | NA | NA | 3.3E-4/ 6.6E-4 | 3.6E-6/ 7.3E-6 | 1.1E-6/ 2.1E-6 | |
| | Ornamental plants | Christmas Tree plantation, conifers, shade trees | Max 2.5 | | 9 | 1120 | NA | 7 | 870 | NA | 28935 | NA | NA | 1.9E-4/ 3.7E-4 | 2.0E-6/ 4.0E-6 | 5.9E-7/ 1.2E-6 | |
| Mixing/Loading Liquids for Application of High Pressure Handwand (1e) | Non-bearing nursery stock | all crops | Max 1.5 | 5 | 120 | NA | NA | 90 | 11595 | NA | 38578 0 | NA | NA | 1.4E-5/ 2.8E-5 | 5.0E-6/ 9.9E-6 | 6.9E-8/ 1.4E-7 | |
| Mixing/Loading Wettable Powder for Aerial Application (2a) | Stone fruits | nectarine | Max 3.0 | 125 | 2 | 40 | 325 | 1 | 30 | 255 | 215 | 2155 | NA | 9.6E-4/ 1.9E-3 | 1.0E-4/ 2.0E-4 | 5.3E-6/ 1.1E-5 | |
| | Tree Nuts | walnut | Max 4.0 | | 1 | 30 | 245 | 1 | 25 | 190 | 160 | 1615 | NA | 1.3E-3/ 2.6E-3 | 1.4E-4/ 2.8E-4 | 7.9E-6/ 1.6E-5 | |

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb ai/acre) | Daily Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | | |
|--|---------------------------------|---|--------------------------------|---------------------|------------------------|------|--------------|-------------------------------|------|--------------|-----------------|--------|--------------|-------------------|-------------------|-------------------|-------------------|
| | | | | | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. control | |
| | Ornamental plants | Christmas Tree | Max 2.5 | 125 | 2 | 50 | 390 | 2 | 40 | 305 | 255 | 2585 | NA | 7.9E-4/ 1.6E-3 | 8.6E-5/ 1.7E-4 | 4.6E-6/ 9.2E-6 | |
| Mixing/Loading Wettable Powder for Groundboom Application (2b) | Oil Seed | peanut | Max 1.6 | 80 | 5 | 120 | NA | 4 | 90 | 745 | 625 | 6310 | NA | 3.2E-6/ 6.5E-4 | 4.0E-5/ 7.9E-5 | 1.8E-6/ 3.6E-6 | |
| Mixing/ Loading Wettable Powder for Airblast Sprayer Application (2c) | Citrus fruits | grapefruit, orange, lemon, avocado | Min 3.0 | 40 | 6 | 125 | NA | 5 | 100 | NA | 665 | 6730 | NA | 3.0E-4/ 6.0E-4 | 3.6E-5/ 7.3E-5 | 1.7E-6/ 3.4E-6 | |
| | Tropical and subtropical fruits | | Max 4.5 | | 4 | 85 | 680 | 3 | 65 | 530 | 445 | 4485 | NA | 4.6E-4/ 9.2E-4 | 5.3E-5/ 1.1E-4 | 2.6E-6/ 5.1E-6 | |
| | Herbs & spices | hops | Min 2.0 | | 9 | 190 | NA | 7 | 145 | NA | 1000 | NA | NA | NA | 2.0E-4/ 4.0E-4 | 2.4E-5/ 4.8E-5 | 1.2E-6/ 2.3E-6 |
| | | | Max 2.5 | | 7 | 150 | NA | 5 | 120 | NA | 800 | 8075 | NA | 2.5E-4/ 5.0E-4 | 3.0E-5/ 5.9E-5 | 1.5E-6/ 2.9E-6 | |
| | Small Fruits | grapes | Max 3.0 | | 6 | 125 | NA | 5 | 100 | NA | 665 | 6730 | NA | NA | 3.0E-4/ 6.0E-4 | 3.6E-5/ 7.3E-5 | 1.7E-6/ 3.4E-6 |
| Mixing/ Loading Wettable Powder for Application of High Pressure Handwand (2d) | Non- bearing Nursery Stock | all crops | Min 0.5 | 5 | 280 | NA | NA | 215 | NA | NA | 32000 | NA | NA | 6.6E-6/ 1.3E-5 | 2.7E-7/ 5.3E-7 | NA | |
| | | | Max 2.5 | | 55 | 1210 | NA | 45 | 940 | NA | 6400 | NA | NA | 3.2E-5/ 6.3E-5 | 1.4E-6/ 2.7E-6 | NA | |
| Applicator Exposure | | | | | | | | | | | | | | | | | |
| Applying Sprays with Fixed-Wing Aircraft-Enclosed Cockpit (3) | Roots and Tuber Vegetable | carrot, sugar beet, potatoes, dry beans, mint | Min 2.0 | 350 | NA | NA | 735 | NA | NA | 570 | NA | NA | 72940 | NA | NA | 2.4E-6/ 4.9E-6 | |
| | | | Max 2.5 | | NA | NA | 585 | NA | NA | 455 | NA | NA | 58355 | NA | NA | 3.0E-5/ 6.1E-5 | |
| | Legume Vegetable | | | | | | | | | | | | | | | | |
| | Herbs and Spices | hops | Max 2.5 | 80 | NA | NA | 2570 | NA | NA | 2000 | NA | NA | 255295 | NA | NA | 6.9E-7/ 1.4E-6 | |
| | Citrus fruits | grapefruit, orange | Max 2.5 | 125 | NA | NA | 1645 | NA | NA | 1280 | NA | NA | 163390 | NA | NA | 1.1E-6/ 2.2E-6 | |
| Tree Nuts | almond, walnut | Min 2.5 | | NA | NA | 1645 | NA | NA | 1280 | NA | NA | 163390 | NA | NA | 1.1E-6/ 2.2E-6 | | |

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb/acre) | Daily Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | |
|--------------------------------|---|--|--|---------------------|------------------------|------|--------------|-------------------------------|------|--------------|-----------------|-------|--------------|-------------------|-------------------|-------------------|
| | | | | | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control | Base line | PPE | Eng. Control |
| | | | Max 4.5 | | NA | NA | 915 | NA | NA | 710 | NA | NA | 90770 | NA | NA | 1.9E-6/ 3.9E-6 |
| | Cereal Grains | corn (field, pop. sweet), sorghum grain, alfalfa, clover | Min 1.5 | 350 | NA | NA | 980 | NA | NA | 760 | NA | NA | 97255 | NA | NA | 1.8E-6/ 3.6E-6 |
| | Non-Grass animal Feed | | Max 2.5 | | NA | NA | 590 | NA | NA | 455 | NA | NA | 58355 | NA | NA | 3.0E-6/ 6.1E-6 |
| | Oil Seed | peanut, jojoba | Min 1.5 | 350 | NA | NA | 980 | NA | NA | 760 | NA | NA | 97255 | NA | NA | 1.9E-6/ 3.9E-6 |
| | | | Max 2.5 | | NA | NA | 590 | NA | NA | 455 | NA | NA | 583550 | NA | NA | 6.6E-6/ 1.3E-5 |
| | | cotton | Max 1.6 | 350 | NA | NA | 920 | NA | NA | 715 | NA | NA | 91175 | NA | NA | 1.8E-6/ 3.6E-6 |
| | | | | 1200 | NA | NA | 270 | NA | NA | 210 | NA | NA | 26595 | NA | NA | 3.0E-6/ 6.1E-6 |
| | Stone fruit | nectarine | Max 3.0 | 125 | NA | NA | 1370 | NA | NA | 1065 | NA | NA | 136155 | NA | NA | 1.3E-6/ 2.6E-6 |
| | Ornamental plants | Christmas tree, conifer seed | Max 2.5 | 125 | NA | NA | 1645 | NA | NA | 1280 | NA | NA | 163390 | NA | NA | 1.1E-6/ 2.2E-6 |
| | Applying Sprays with a Groundboom Sprayer (4) | Roots and Vegetable | potatoes, corn (field, pop. sweet) sorghum grain, alfalfa, clover, cotton, peanut, jojoba and mint | Min 1.5 | 80 | 1530 | NA | NA | 1190 | NA | NA | 39100 | NA | NA | 1.4E-6/ 2.7E-6 | 1.2E-6/ 2.4E-6 |
| Cereal Grain | | Max 2.5 | | 920 | | NA | NA | 715 | NA | NA | 23460 | NA | NA | 2.4E-6/ 4.9E-6 | 2.0E-6/ 4.1E-6 | 6.9E-7/ 1.4E-6 |
| Non-grass animal feed | | | | | | | | | | | | | | | | |
| oil seed | | | | | | | | | | | | | | | | |
| herbs and spices | | | | | | | | | | | | | | | | |

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb ai/acre) | Duties Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | | |
|--|-------------------------------|---|--------------------------------|----------------------|------------------------|-----|-------------|-------------------------------|-----|-------------|-----------------|-----|-------------|-------------------|-------------------|-------------------|--|
| | | | | | Base line | PPE | Eng Control | Base line | PPE | Eng Control | Base line | PPE | Eng Control | Base line | PPE | Eng control | |
| Applying Sprays with an Airblast Sprayer (5) | pome fruits | quince, cherry, prunes, orange, grapefruit, lemon, lime, tangerine, boysenberry, current, hops, raspberry, date, persimmons, almond, filbert, macadamia nut, pecan, pistachio, walnut, Christmas Tree plantation, conifers, shade trees | Min 1.5 | 40 | 120 | NA | NA | 95 | 140 | NA | 12860 | NA | NA | 1.5E-5/ 3.0E-5 | 1.1E-5/ 2.1E-5 | 2.0E-6/ 4.0E-6 | |
| | stone fruits | | | | 40 | 120 | 750 | 30 | 90 | 570 | 4285 | NA | NA | 4.6E-5/ 9.2E-5 | 3.1E-5/ 6.2E-5 | 6.9E-6/ 1.4E-5 | |
| | citrus fruits | | Max 4.5 | | | | | | | | | | | | | | |
| | berries | | | | | | | | | | | | | | | | |
| | tropical & subtropical fruits | | | | | | | | | | | | | | | | |
| | small fruits | | | | | | | | | | | | | | | | |
| | tree nuts | | | | | | | | | | | | | | | | |
| ornamental plants | | | | | | | | | | | | | | | | | |
| Applying Liquids with a High Pressure Handwand (6) | non-bearing nursery stock | all crops | Min 0.5 | 5 | 570 | NA | NA | 445 | NA | NA | 17580 | NA | NA | 3.6E-6/ 7.3E-6 | 1.9E-6/ 3.8E-6 | NA | |
| | | | Max 2.5 | | 115 | NA | NA | 90 | 250 | NA | 3515 | NA | NA | 1.9E-5/ 3.8E-5 | 9.6E-6/ 1.9E-5 | NA | |
| Flagger Exposure | | | | | | | | | | | | | | | | | |
| Flagging During Aerial Spray Application (7) | Roots & tuber Vegetable | carrot, sugar beet, potatoes, dry beans, mint | Min 2.0 | 350 | 335 | NA | NA | 260 | NA | NA | 14170 | NA | NA | 5.9E-6/ 1.2E-5 | 4.6E-6/ 9.1E-6 | 1.2E-7/ 2.4E-7 | |
| | legume vegetable | | Max 2.5 | | 265 | NA | NA | 210 | NA | NA | 11335 | NA | NA | 7.6E-6/ 1.5E-5 | 5.7E-6/ 1.1E-5 | 1.5E-7/ 3.0E-7 | |
| | herbs and spices | hops | Ma 2.5 | 80 | 1170 | NA | NA | 910 | NA | NA | 49600 | NA | NA | 1.7E-6/ 3.4E-6 | 1.3E-6/ 2.6E-6 | 3.4E-8/ 6.9E-8 | |
| | Citrus fruits | grapefruit, orange | Max 2.5 | 125 | 750 | NA | NA | 580 | NA | NA | 31745 | NA | NA | 2.7E-6/ 5.4E-6 | 2.0E-6/ 4.1E-6 | 5.3E-8/ 1.1E-7 | |
| | Tree Nut | almond, walnut | Min 2.5 | 125 | 750 | NA | NA | 580 | NA | NA | 31745 | NA | NA | 2.7E-6/ 5.4E-6 | 2.0E-6/ 4.1E-6 | 5.3E-8/ 1.1E-7 | |
| | | | Max 4.5 | | 415 | NA | NA | 325 | NA | NA | 17635 | NA | NA | 5.0E-6/ 9.9E-6 | 3.7E-6/ 7.3E-6 | 9.6E-8/ 1.9E-7 | |
| | cereal grain | (field, pop, sweet), sorghum grain, alfalfa, clover | Min 1.5 | 350 | 445 | NA | NA | 345 | NA | NA | 18895 | NA | NA | 4.6E-6/ 9.2E-6 | 3.4E-6/ 6.8E-6 | 9.0E-8/ 1.8E-7 | |
| | non-grass animal feed | | Max 2.5 | | 265 | NA | NA | 210 | NA | NA | 11335 | NA | NA | 7.6E-6/ 1.5E-5 | 5.7E-6/ 1.1E-5 | 1.5E-7/ 3.0E-7 | |
| | oil seed | cotton | Max 1.6 | 350 | 415 | NA | NA | 325 | NA | NA | 17715 | NA | NA | 5.0E-6/ 9.9E-6 | 3.6E-6/ 7.3E-6 | 9.6E-8/ 1.9E-7 | |

| Exposure Scenario (Scenario #) | Crop Group | Crop | Application Rates (lb ai/acre) | Daily Acres Treated | Short-Term Dermal MOEs | | | Intermediate-Term Dermal MOEs | | | Inhalation MOEs | | | Cancer | | |
|--------------------------------|-------------------|-----------------------------|--------------------------------|---------------------|------------------------|-----|--------------|-------------------------------|-----|--------------|-----------------|-----|--------------|-------------------|-------------------|-------------------|
| | | | | | Baseline | PPE | Eng. Control | Baseline | PPE | Eng. Control | Baseline | PPE | Eng. Control | Baseline | PPE | Eng. Control |
| | | | | 1200 | 120 | NA | NA | 100 | NF | NA | 5165 | NA | NA | 1.7E-5/ 3.3E-5 | 1.2E-5/ 2.5E-5 | 3.3E-7/ 6.6E-7 |
| | | Peanut, jojoba | Min 1.5 | 350 | 445 | NA | NA | 345 | NA | NA | 18895 | NA | NA | 4.6E-6/ 9.2E-6 | 3.4E-6/ 6.8E-6 | 9.0E-8/ 1.8E-7 |
| | | | Max 2.5 | | 265 | NA | NA | 210 | NA | NA | 11335 | NA | NA | 7.6E-6/ 1.5E-5 | 5.7E-6/ 1.1E-5 | 1.5E-7/ 3.0E-7 |
| | Stone fruits | nectarine | Max 3.0 | 125 | 625 | NA | NA | 485 | NA | NA | 26455 | NA | NA | 3.2E-6/ 6.4E-6 | 1.1E-6/ 2.2E-6 | 6.4E-8/ 1.3E-7 |
| | Ornamental plants | Christmas tree conifer seed | Max 2.5 | 125 | 750 | NA | NA | 580 | NA | NA | 31745 | NA | NA | 2.7E-5/ 5.4E-5 | 2.0E-6/ 4.1E-6 | 5.3E-8/ 1.1E-7 |

- ^a Baseline Total Daily Dose = [Baseline Daily Dermal Exposure (mg/day) * 0.14 (Dermal Absorption Factor) + Baseline Daily Inhalation Exposure (mg/day)]/Body Weight (70 kg).
- ^b Baseline LADD (mg/kg/day) = Baseline Total Daily Dose (mg/kg/day) * (Number of days exposure per year (7 private applicator and 14 for commercial applicator) /365 days per year) * 35 years worked/70 year lifetime.
- ^c Baseline Total Cancer Risk = Baseline LADD (mg/kg/day) * (Q₁*), where 3.3x10⁻² (mg/kg/day)⁻¹.
- ^d PPE Total Daily Dose = [PPE Daily Dermal Exposure (mg/day) * 0.14 (Dermal Absorption Factor) + baseline Daily Inhalation Exposure (mg/day)]/Body Weight (70 kg).
- ^e PPE LADD (mg/kg/day) = PPE Total Daily Dose (mg/kg/day) * (Number of days exposure per year (7 private applicator and 14 for commercial applicator) /365 days per year) * 35 years worked/70 year lifetime.
- ^f PPE Total Cancer Risk = PPE LADD (mg/kg/day) * (Q₁*), where 3.3x10⁻² (mg/kg/day)⁻¹.
- ^g Eng. Control Total Daily Dose = [Eng. Control Daily Dermal Exposure (mg/day) * 0.14 (Dermal Absorption Factor) + baseline Daily Inhalation Exposure (mg/day)]/Body Weight (70 kg).
- ^h Eng. Control LADD (mg/kg/day) = Eng. Control Total Daily Dose (mg/kg/day) * (Number of days exposure per year (7 private applicator and 14 for commercial applicator) /365 days per year) * 35 years worked/70 year lifetime.
- ⁱ Eng. Control Total Cancer Risk = Eng. Control LADD (mg/kg/day) * (Q₁*), where Q₁* = 3.3x10⁻² (mg/kg/day)⁻¹.

4.3.2 Occupational Postapplication Exposure Scenarios

EPA has determined that there are potential short and intermediate-term postapplication exposures to individuals entering treated fields for the purpose of postapplication activities.

For the purpose of conducting this assessment, indicator crop groups/activities, and assumptions regarding application rates and dermal transfer coefficients for these crop groups were selected that are likely to be representative of postapplication exposures to propargite. The crop groups/activities listed below were chosen because appropriate residue data were available, and exposure assumptions could be made that would be inclusive of other similar crop types/activities. Although several studies have been submitted, it was still necessary to use standard transfer coefficients and crop-specific residues as substitutes to represent other crops. Also, the development of these exposure scenarios followed the guidance provided in the Science Advisory Council for Exposure Policy Memo Number 003.1 (Revised August, 2000). The postapplication exposure scenarios include the following:

- All activities associated with **legume vegetable, roots and tuber vegetable and non grass animal feed** groups. This scenario is assumed to be representative of exposures from typical weeding and irrigation activities. DFR and passive dosimetry data for dry beans were used, based on studies using an application rate of 2.46 lb ai/acre. This application rate is consistent with the application rates for most crops in these groups. A dermal transfer coefficient of 60 cm²/hr was calculated from a weeder reentry study (MRID No. 426891-04) to represent weeding and hoeing activities for dry beans. For irrigation and scouting a dermal transfer coefficient of 1500 cm²/hr, and for hand harvesting transfer coefficient of 2500 cm²/hr were used from an ARTF study (ARF021 -- scouting dry peas).
- All activities associated with **citrus** fruits. This scenario is assumed to be representative of exposures from all activities. DFR data for navel oranges were used, based on a study using maximum application rates of 3.15 lb ai/acre and 4.5 lb ai/acre. Dermal transfer coefficients of 1000 cm²/hr for irrigation, scouting, and hand weeding from ARF023 -- scouting table grapes, 3000 cm²/hr for pruning from a citrus hand pruning study (MRID NO. 430627) and 8000 cm²/hr for hand harvesting from an apple thinning study (MRID NO.424281).
- All activities associated with **stone fruits, pome fruits, tropical and subtropical fruits, ornamental plants**. This scenario is assumed to be representative of exposures from all activities. DFR data for apple were used, based on a study using application rates of 3.6 lb ai/acre and 1.7 lb ai/acre in states of Vermont and Washington. Dermal transfer coefficients of 1000 cm²/hr for irrigation, scouting, and hand weeding from ARF023 -- scouting table grapes study, 3000 cm²/hr for pruning from a peach hand harvesting study (MRID NO. 428300) and 8000 cm²/hr for hand harvesting from an apple thinning study (MRID NO.424281).

- All activities associated with **berries**. This scenario is assumed to be representative of exposures from all activities. DFR data for grape were used, based on a study using a maximum application rate of 2.7 lb ai/acre. Dermal transfer coefficients of 500 cm²/hr for irrigation, and hand weeding from an ARTF study (ARF023 -- scouting table grapes), 1000 cm²/hr for scouting from an ARTF study (ARF023 -- scouting table grapes) and 5000 cm²/hr for hand harvesting, pruning, tying from a hand harvesting raisin grapes study (MRID NO.409856).
- All activities associated with **small fruits**. This scenario is assumed to be representative of exposures from all activities. DFR and passive dosimetry data for grapes were used, based on studies using a maximum application rate of 2.7 lb ai/acre. Dermal transfer coefficients of 500 cm²/hr for irrigation, and hand weeding from ARF023 -- scouting table grapes study, 1000 cm²/hr for scouting from an ARTF study (ARF023 -- scouting table grapes), 5000 cm²/hr for hand harvesting, pruning, tying from a hand harvesting raisin grapes study (MRID NO.409856) and for cane turning 10,000 cm²/hr from a cane turning study in table grapes (MRID NO. 409753).
- All activities associated with **tree nuts**. This scenario is assumed to be representative of exposures from all activities. DFR data for almonds were used, based on a study using a maximum application rate of 3.0 lb ai/acre. Dermal transfer coefficient of 48 cm²/hr was calculated from a worker reentry study (MRID NO. 418486-04) which represents tree shaker scenario. Transfer coefficients of 500 cm²/hr for scouting weeding and irrigation from an ARTF study (ARF023 -- scouting table grapes) and 2500 cm²/hr for pruning and hand harvesting from citrus hand pruning study (MRID NO. 430627).
- All activities associated with **cereal grains**. This scenario is assumed to be representative of exposures from all activities. DFR data for corn were used, based on a study using an application rate of 2.46 lb ai/acre. Dermal transfer coefficients of 1000 cm²/hr for irrigation and scouting from an ARTF study (ARF009--scouting sweet corn) and 17000 cm²/hr for detasseling and hand harvesting from an ARTF study (ARF010--hand harvesting sweet corn) were used.
- All activities associated with **herbs and spices**. This scenario is assumed to be representative From all activities. DFR data for hops were used, based on a study using an application rate of 1.35 lb ai/acre. For mint dermal transfer coefficient of 1500 cm²/hr from ARF021-- scouting dry peas study was used. For hops dermal transfer coefficients of 100 cm²/hr for irrigation, hand weeding and scouting from a cotton and beans hoeing study (MRID NO.426891) and 2000 cm²/hr from harvesting and training from an ARTF study (ARF024-- hand harvesting tobacco) were used.
- All activities associated with **oil seed crop** group. This scenario is assumed to be representative of all activities. DFR data for cotton were used, based on studies using an application rate of 1.64 lb ai/acre. This application rate is consistent with the application

rates for most crops in these groups. Dermal transfer coefficients of 63 cm²/hr was calculated from a weeder reentry study (MRID No.426891-03) to represent weeding activities and 2500 cm²/hr to represent harvesting activities from an ARTF study (ARF021–scouting dry peas) associated with cotton. Dermal transfer coefficient of 1500 cm²/hr for irrigation and scouting from an ARTF study (ARF021–scouting dry peas) was also used.

Currently, HED conducts post-application exposure and risk assessments assuming that the workers are wearing “typical” work clothing – long-sleeved shirt, long pants, shoes and socks. Additional personal protective equipment is not considered in these assessments.

4.3.2.1 Occupational Postapplication Exposure Data Sources and Assumptions

For the purposes of this assessment, regression analysis were conducted using the natural log-transformed DFR data from the above studies to estimate residue levels on various crops on various days for postapplication using the following equation:

$$y = mx + b$$

where:

- x = days postapplication;
- m = slope of the regression line;
- b = constant; and
- y = residue on day x .

The summary of regression analysis on submitted studies is presented in Table 12.

Table 12. Summary of Data used for Post Application of Propargite

| Crop grouping | DFR Study | Formulation Type | Study Application Rate (lb ai/acre) | R ² | Initial DFR as a % of Appl. Rate (Day 0 measured values) | Transfer Coefficient | Dissipation (% per day) | ½ Life (days) | |
|---|---|------------------|-------------------------------------|----------------|--|----------------------|-------------------------|---------------|------|
| Legume vegetable, roots and tuber vegetable and non-grass animal feed | Dry beans (MRID 420118-01, 426891-04) | Comite EC | 2.46 | 0.96 | 18.9 | 60 | 11 | 5.97 | |
| Citrus | Navel oranges (MRID 409090-03) | Omite CR | 3.15 | 0.78 | 9.2 | NA | 6 | 12.1 | |
| | | Omite 30W | 3.15 | 0.65 | 4.9 | NA | 4 | 15.5 | |
| | | Omite CR | 4.5 | 0.74 | 10.4 | NA | 5 | 12.9 | |
| Stone fruits, Pome fruits, Tropical and subtropical fruits, ornamental plants | Apple (MRID 409090-04) | Omite CR | V T | 3.6 | 0.98 | 4.2 | NA | 10 | 6.8 |
| | | | W A | 3.6 | 0.21 | 4.5 | NA | 4 | 18.1 |
| | | Omite 30W | V T | 3.6 | 0.99 | 4.2 | NA | 10 | 6.5 |
| | | | W A | 3.6 | 0.37 | 2.1 | NA | 3 | 20.6 |
| Small fruits | Grape (MRID 409753-01, MRID 409753-04) | Omite 30W | | 2.7 | 0.72 | 2.7 | 878 | 6 | 10.7 |
| | | | | | | | 1895 | | |
| | | | | | | | 3713 | | |
| | | | | | | | 10246 | | |
| Tree nuts | Almond (MRID 418486-04, MRID 418486-03) | Omite 6E | 3.0 | 0.79 | 6.0 | 48 | 5 | 13.6 | |
| Cereal grain | Corn (MRID 416803-02) | Comite EC | 2.46 | 0.72 | 6.4 | NA | 18 | 3.5 | |
| Herbs and spices | Hops (MRID 426891-03) | Comite EC | 1.35 | 0.72 | 14.5 | NA | 4 | 18.6 | |
| Oil seed | Cotton (MRID 426891-03, MRID 414578-06) | Comite EC | 1.64 | 0.92 | 11 | 63 | 11 | 6.2 | |

The postapplication exposure assessment encompasses all of the major uses of propargite throughout the country. It is difficult to assess all of the “typical” agricultural uses for propargite (i.e., actual or predominate application rates and climatological conditions), and therefore, an

assessment has been developed which is believed to be realistic and yet provides a reasonable certainty that the exposures are not underestimated. The assumptions and uncertainties are identified below to be used in risk management decisions:

- *Crop Specific Residues:* A multitude of crops are treated with propargite and crop specific residue data are not available for all situations. Therefore, the use of the available data to “simulate” residues on other crops introduces uncertainties in the setting of reentry intervals. It is reasonable to believe that the residues monitored in the available studies approximate the residues on other crops, but the extent that these residues might be an under- or overestimate is unknown. The DFR results from these crops may alter the surrogate assessment for determining REIs.
- *Transfer Coefficients:* The transfer coefficients selected are based on the activities monitored in the submitted studies and on HED’s policy which is based on data submitted to the Agency or are from published literature studies. Usually, a “central” value from the selected study was used. However, to account for some uncertainty inherent in translating data between crops, activities, and groups, higher transfer coefficient values were sometime used. These values are believed to be reasonable estimates that would not underestimate the risks.
- *Exposure Duration:* The amount of time (e.g., days) that a worker would be involved in postapplication activities is not available. Therefore, both short-term and intermediate-term exposure durations are provided and the intermediate-term duration is believed to be most representative for the postapplication exposures. Furthermore, the REIs are calculated at the residue level predicted on a specific day after treatment; subsequent declining residue levels (i.e., average residues under the dissipation curve) are not incorporated into the assessment because of the lack of exposure duration data (including the fact that harvesters may travel to multiple fields). Note: Scouts are assumed to be exposed eight hours per day, which may be an overestimation.
- Fully Mechanized is defined as activities that eliminates the potential of pesticide exposure by physically separating the worker from anything that has been treated with the pesticide to which the restricted-entry interval applies, including, but not limited to, soil, water, air, or surfaces of plants. **These mechanized processes must meet the criteria described in the Worker Protection Standard for entry during an REI for activities with “No Contact.”** Examples of “no contact” mechanical processes include harvesting small grains or other crops using combines with closed cabs and cultivating crops (mechanical weed control) with closed cab tractors. Note that if, as is typical, these activities raise significant dust, the closed cab must provide respiratory protection to prevent inhalation exposure. Exposure data for these activities are not required because the triggers for conditional data requirements under 40 CFR §158.390 are not met.

If the workers must exit the closed cab while in the treated area (e.g., to unclog equipment), then they are considered to have potentially come into contact with treated surfaces. During an REI, the workers exiting the cab may use the Worker Protection Standard §170.112 *Exception for short-term activities*, which allows entry into treated areas for a maximum of one hour per day to perform tasks (other than hand labor tasks), as long as early-entry personal protective equipment is worn and the other early-entry requirements are met (i.e., training, decontamination sites, labeling instructions, etc.). However, it should be noted that the early-entry PPE is established for dermal protection only and presumes that pesticide residues have settled out of the air. If the mechanical activity has caused dusts that contain residues to become airborne, the exiting worker will not have respiratory protections, since early-entry PPE does not include a respirator.

4.3.2.2 Occupational Postapplication Risk Characterization

Short-term Risks

The target dermal MOE is 100 for propargite. The results of the occupational postapplication assessment are presented in Tables 13-33, and are summarized below:

- Propargite MOEs equal or exceed 100 for hand harvesting of roots and tuber vegetable (carrot, sugar beet, potato) on day 7 at 1.5 lb ai/A and on day 11 at 2.5 lb ai/A. MOEs for All other activities equal or exceed 100 on day 0.
- Propargite MOEs equal or exceed 100 for hand harvesting of legume vegetable (dry beans) on day 7 at 1.5 lb ai/A and on day 11 at 2.5 lb ai/A. MOE,s equal or exceed 100 for irrigation and scouting on day 3 at 1.5 lb ai/A and on day 7 at 2.5 lb ai/A and for weeding and hoeing on day 0.
- Propargite MOEs equal or exceed 100 for irrigation and scouting of non-grass animal feed (alfalfa, clover) on day 3 at 1.5 lb ai/A and on day 7 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of citrus (orange, lemon, lime, tangerine and grapefruit) on day 0 at 2.5 lb ai/A, on day 0 at 3.15 lb ai/A and on day 4 at 4.5 lb ai/A. MOE,s equal or exceed 100 for Pruning on day 12 at 2.5 lb ai/A, on day 16 at 3.15 lb ai/A and on day 21 at 4.5 lb ai/A. MOE,s equal or exceed 100 for harvesting on day 28 at 2.5 lb ai/A, on day 32 at 3.15 lb ai/A and on day 37 at 4.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand hand weeding of pome fruits, stone fruits, tropical & subtropical fruits and ornamental plants (Quince, Cherry, Nectarine, Prune, Avocado, Date, Persimmons, X mas Tree, Ornamental and/or shade trees Ornamental, Herbaceous Plants) on day 0 at 0.5 lb ai/A, on day 0 at 1.5 lb ai/A, on day 0 at 2.5 lb ai/A and on day 0 at 4.5 lb ai/A. MOE,s equal or exceed 100 for Pruning

on day 0 at 0.5 lb ai/A, on day 0 at 1.5 lb ai/A, on day 3 at 2.5 lb ai/A and on day 8 at 4.5 lb ai/A. MOE,s equal or exceed 100 for harvesting on day 0 at 0.5 lb ai/A, on day 7 at 1.5 lb ai/A, on day 12 at 2.5 lb ai/A and on day 18 at 4.5 lb ai/A.

- Propargite MOEs equal or exceed 100 for irrigation, hand weeding and scouting of berries(boysenberry, currant and raspberry) on day 0 at 2.0 lb ai/A. MOE,s equal or exceed 100 for harvesting, pruning and tying on day 6 at 2.0 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, hand weeding and scouting of small fruits(grapes) on day 0 at 3.0 lb ai/A. MOE,s equal or exceed 100 for harvesting, pruning and tying on day 12 at 3.0 lb ai/A. MOE,s equal or exceed 100 for cane turning day 17 at 3.0 lb ai/A.
- Propargite MOEs equal or exceed 100 for tree shaking, irrigation, hand weeding and scouting of tree nuts(Almond, Filbert, Macadamia ,Pecan,Pistachio, Walnut) on day 0 at 3.0 lb ai/A and 4.5 lb ai/A.. MOE,s equal or exceed 100 for hand harvesting and pruning on day 17 at 3.0 lb ai/A and on day 25 at 4.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, and scouting of cereal grain (Corn (unspecified), Corn, field, Corn, Pop Corn, Sweet, Sorghum, grain) on day 0 at 2.5 lb ai/A. MOE,s equal or exceed 100 for hand harvesting and detasseling on day 11 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of herbs and spices (mint) on day 7 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of herbs and spices (hops) on day 0 at 2.5 lb ai/A MOE,s equal or exceed 100 for harvesting and training on day 27 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation and scouting oil seed (cotton, peanut and jojoba) on day 0 at 1.5 lb ai/A MOE,s equal or exceed 100 for harvesting (cotton) on day 3 at 1.5 lb ai/A and for weeding and hoeing on day 0 at 1.5 lb ai/A.

Intermediate-term Risks

The target MOE is 100 for propargite. The resulting occupational postapplication assessments, as shown in Tables 13 through 33, indicate that:

- Propargite MOEs equal or exceed 100 for hand harvesting of roots and tuber vegetable (carrot, sugar beet, potato) on day 9 at 1.5 lb ai/A and on day 13 at 2.5 lb ai/A. MOEs for All other activities equal or exceed 100 on day 0.

- Propargite MOEs equal or exceed 100 for hand harvesting of legume vegetable (dry beans) on day 9 at 1.5 lb ai/A and on day 13 at 2.5 lb ai/A. MOE,s equal or exceed 100 for irrigation and scouting on day 5 at 1.5 lb ai/A and on day 9 at 2.5 lb ai/A and for weeding and hoeing on day 0.
- Propargite MOEs equal or exceed 100 for irrigation and scouting of non-grass animal feed (alfalfa, clover) on day 5 at 1.5 lb ai/A and on day 9 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of citrus (orange, lemon, lime, tangerine and grapefruit) on day 0 at 2.5 lb ai/A, on day 2 at 3.15 lb ai/A and on day 8 at 4.5 lb ai/A. MOE,s equal or exceed 100 for Pruning on day 16 at 2.5 lb ai/A, on day 20 at 3.15 lb ai/A and on day 26 at 4.5 lb ai/A. MOE,s equal or exceed 100 for harvesting on day 32 at 2.5 lb ai/A, on day 36 at 3.15 lb ai/A and on day 42 at 4.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand hand weeding of pome fruits, stone fruits, tropical & subtropical fruits and ornamental plants (Quince, Cherry, Nectarine, Prune, Avocado, Date, Persimmons, X mas Tree, Ornamental and/or shade trees Ornamental, Herbaceous Plants) on day 0 at 0.5 lb ai/A, on day 0 at 1.5 lb ai/A, on day 0 at 2.5 lb ai/A and on day 0 at 4.5 lb ai/A. MOE,s equal or exceed 100 for Pruning on day 0 at 0.5 lb ai/A, on day 0 at 1.5 lb ai/A, on day 5 at 2.5 lb ai/A and on day 11 at 4.5 lb ai/A. MOE,s equal or exceed 100 for harvesting on day 0 at 0.5 lb ai/A, on day 9 at 1.5 lb ai/A, on day 14 at 2.5 lb ai/A and on day 20 at 4.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, hand weeding and scouting of berries(boysenberry, currant and raspberry) on day 0 at 2.0 lb ai/A. MOE,s equal or exceed 100 for harvesting, pruning and tying on day 10 at 2.0 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, hand weeding and scouting of small fruits(grapes) on day 0 at 3.0 lb ai/A. MOE,s equal or exceed 100 for harvesting, pruning and tying on day 16 at 3.0 lb ai/A. MOE,s equal or exceed 100 for cane turning day 27 at 3.0 lb ai/A.
- Propargite MOEs equal or exceed 100 for tree shaking, irrigation, hand weeding and scouting of tree nuts(Almond, Filbert, Macadamia ,Pecan, Pistachio, Walnut) on day 0 at 3.0 lb ai/A and 4.5 lb ai/A.. MOE,s equal or exceed 100 for hand harvesting and pruning on day 22 at 3.0 lb ai/A and on day 30 at 4.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, and scouting of cereal grain (Corn (unspecified), Corn, field, Corn, Pop Corn, Sweet, Sorghum, grain) on day 0 at 2.5 lb ai/A . MOE,s equal or exceed 100 for hand harvesting and detasseling on day 13 at 2.5 lb ai/A.

- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of herbs and spices (mint) on day 9 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation, scouting and hand weeding of herbs and spices (hops) on day 0 at 2.5 lb ai/A MOE,s equal or exceed 100 for harvesting and training on day 33 at 2.5 lb ai/A.
- Propargite MOEs equal or exceed 100 for irrigation and scouting of oil seed (cotton, peanut and jojoba) on day 0 at 1.5 lb ai/A MOE,s equal or exceed 100 for harvesting (cotton) on day 6 at 1.5 lb ai/A and for weeding and hoeing on day 1 at 1.5 lb ai/A.

Table 13. Summary of the Short- and Intermediate-Term Reentry Intervals (REIs) for the Contact Rates and Crop Grouping Matrix.

| Crop grouping | Crop | Activity | Short Term- REI (days) | | | Intermediate -Term REI (days) | | | | |
|---------------------------|-------------------------------|--|----------------------------------|--------------|----------------------------------|----------------------------------|-------------|----------------------------------|-------------|-------------|
| | | | 1.5 lb ai/A | | 2.5 lb ai/A | 1.5 lb ai/A | | 2.5 lb ai/A | | |
| Roots and Tuber Vegetable | Potato | Hand Harvesting | Not available Special concern | | Not available Special concern | Not available Special concern | | Not available Special concern | | |
| | Carrot, Sugar beet | Hand Harvesting | 7 | | 11 | 9 | | 13 | | |
| | Potato, Carrot, Sugar beet | all other activities | 0 | | 0 | 0 | | 0 | | |
| Legume Vegetables | Dry beans | Weeding and Hoeing | 0 | | 0 | 0 | | 0 | | |
| | | Irrigation, Scouting | 3 | | 7 | 5 | | 9 | | |
| | | Hand Harvesting | 7 | | 11 | 9 | | 13 | | |
| Non-grass Animal Feed | Alfalfa, Clover | Hand Harvesting, Mechanized harvesting | Not available Special concern | | Not available Special concern | Not available Special concern | | Not available Special concern | | |
| | | Irrigation, Scouting | 3 | | 7 | 5 | | 9 | | |
| Citrus Fruits | Crop | | 2.5 lb ai/A | 3.15 lb ai/A | 4.5 lb ai/A | | 2.5 lb ai/A | 3.15 lb ai/A | 4.5 lb ai/A | |
| | | Orange, Lemon, Lime, Tangerine, Grapefruit | 0 | 0 | 4 | | 0 | 2 | 8 | |
| | | Pruning | 12 | 16 | 21 | | 16 | 20 | 26 | |
| | | Harvesting | 28 | 32 | 37 | | 32 | 36 | 42 | |
| Pome Fruits | Crop | | 0.5 lb ai/A | 1.5 lb ai/A | 2.5 lb ai/A | 4.5 lb ai/A | 0.5 lb ai/A | 1.5 lb ai/A | 2.5 lb ai/A | 4.5 lb ai/A |
| | | Quince, Cherry, Nectarine, Prune, Avocado, Date, Persimmons, X mas Tree, Ornamental and/or shade trees | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | Stone Fruits | 0 | 0 | 3 | 8 | 0 | 0 | 5 | 11 |
| | | Tropical and Subtropical Fruits | 0 | 7 | 12 | 18 | 0 | 9 | 14 | 20 |
| Ornamental Plants | Ornamental, Herbaceous Plants | | | | | | | | | |
| Berries | Crop | | 2.0 lb ai/A | | | | 2.0 lb ai/A | | | |
| | | Boysenberry, Currant, Raspberry | 0 | | | | 0 | | | |

| | | | | | | |
|------------------|--|--|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| | | Scouting | 0 | | 0 | |
| | | Harvesting, Pruning, tying | 6 | | 10 | |
| Small fruits | Crop | | 3.0 lb ai/A | | 3.0 lb ai/acre | |
| | Grape | Irrigation, Hand weeding | 0 | | 0 | |
| | | Scouting | 0 | | 0 | |
| | | Harvesting, Pruning, tying | 12 | | 16 | |
| | | Cane turning | 23 | | 27 | |
| Tree Nuts | Crop | | 3.0 lb ai/acre | 4.5 lb ai/A | 3.0 lb ai/A | 4.5 lb ai/A |
| | Almond, Filbert, Macadamia, Pecan Pistachio, Walnut | Sweeping and blowing the nuts | Not available Special concern | Not available Special concern | Not available Special concern | Not available Special concern |
| | | Tree shakers | 0 | 0 | 0 | 0 |
| | | Scouting, Weeding, Irrigation | 0 | 0 | 0 | 0 |
| | | Pruning, Hand harvesting | 17 | 25 | 22 | 30 |
| | Crop | | 2.5 lb ai/A | | 2.5 lb ai/A | |
| Cereal grains | Corn (unspecified), Corn, field, Corn, Pop Corn, Sweet, Sorghum, grain | Irrigation, Scouting | 0 | | 0 | |
| | | Corn (unspecified), Corn, field, Corn, Pop Corn, Sweet | Hand harvesting, Detasseling | 11 | | 13 |
| Herbs and Spices | Crop | | 2.5 lb ai/A | | 2.5 lb ai/A | |
| | Mint | Irrigation, Scouting, Hand weeding | 7 | | 9 | |
| | Hops | Irrigation, Hand weeding, Scouting | 0 | | 0 | |
| | | Harvesting, Training | 27 | | 33 | |
| Oil seed | Crop | | 1.5 lb ai/A | | 1.5 lb ai/A | |
| | Cotton | Weeding and hoeing | 0 | | 0 | |
| | | Harvesting | 3 | | 6 | |
| | Cotton Peanut, Jojoba | Irrigation, Scouting | 0 | | 1 | |

Cancer Risks

REIs have been estimated using the short- and intermediate-term endpoints. Additionally, the cancer endpoint was used to estimate REIs. HED's target range for cancer probabilities are $1E-4$ to $1E-6$ for occupational assessments. Historically, setting REIs on cancer endpoints has been difficult because of the need for lifetime use assumptions. To estimate the LADD (Life time Average Daily Dose) the typical application rate, the number of days worked per year, and the number of years one would be exposed during a working lifetime are needed. Each one of these variables are dependent upon many factors. For example, the number of days worked per year must correspond to the days worked when the pesticide of concern has been applied.

Additionally, the residue dissipation over the work interval should be estimated. Without an estimate for residue dissipation one needs to assume (unrealistically) that the worker travels from one treated field to another so that the highest residue value is always found. In the case of propargite, a screening estimate was developed because lifetime use data are not available. The screening level estimate assumed: (1) that workers would be exposed for 30 days; (2) no residue dissipation; (3) range of application rates; and (4) a worker would be exposed for 35 years. Based on these assumptions, the cancer probabilities on the day the REIs were estimated using the subchronic endpoints, ranged from $9.74E-6$ to $1.00E-4$.

Table 14. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Roots and Tuber Vegetable at 1.5 lb ai/acre.

| All activities- Potato, Carrot, Sugar beet Transfer Coefficient = 300 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.0155 | 0.0133 | 385 | 300 | 5.47E-4 | 1.81E-5 |
| Hand Harvesting activities- Carrot, Sugar beet Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.1295 | 0.1110 | 45 | 35 | 4.56E-3 | 1.50E-4 |
| 7 | 1.227 | 0.0573 | 0.0491 | 105 | 80 | 2.02E-3 | 6.67E-5 |
| 9 | 0.972 | NA | 0.0389 | NA | 100 | 1.60E-3 | 5.28E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 15. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Roots and Tuber Vegetable at 2.5 lb ai/acre.

| All activities- Potato, Carrot, Sugar beet Transfer Coefficient = 300 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.624 | 0.0259 | 0.0222 | 230 | 180 | 9.12E-4 | 3.01E-5 |
| Hand Harvesting activities- Carrot, Sugar beet Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.624 | 0.2158 | 0.1849 | 30 | 20 | 7.60E-3 | 2.51E-4 |
| 11 | 1.283 | 0.0599 | 0.0513 | 100 | 80 | 2.11E-3 | 6.96E-5 |
| 13 | 1.016 | NA | 0.0407 | NA | 100 | 1.67E-3 | 5.51E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 16. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Non-Grass Animal Feed at 1.5 lb ai/acre.

| Irrigation/Scouting-Alfalfa, Clover Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.0777 | 0.0666 | 75 | 60 | 2.74E-3 | 9.04E-5 |
| 3 | 2.197 | 0.0615 | 0.0527 | 100 | 75 | 2.17E-3 | 7.16E-5 |
| 5 | 1.549 | NA | 0.0372 | NA | 105 | 1.53E-3 | 5.05E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 17. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Non-Grass Animal Feed at 2.5 lb ai/acre.

| Irrigation/Scouting- Alfalfa, Clover Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.624 | 0.1295 | 0.1110 | 45 | 35 | 4.56E-3 | 1.50E-4 |
| 7 | 2.045 | 0.0573 | 0.0491 | 105 | 80 | 2.02E-3 | 6.67E-5 |
| 9 | 1.620 | NA | 0.0389 | NA | 100 | 1.60E-3 | 5.28E-5 |

^a DAT = days after treatment.
^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)
^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].
^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.
^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.
^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]. , where adult body weight = 70 kg.
^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 18. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Dry Beans applied at 1.5 lb ai/acre.

| Weeding/hoeing activities- Dry Beans Transfer Coefficient = 60.0 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.0031 | 0.0027 | 1930 | 1500 | 1.09E-4 | 3.60E-6 |
| Irrigation/scouting activities- Dry Beans Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.0777 | 0.0666 | 75 | 60 | 2.74E-3 | 9.04E-5 |
| 3 | 1.956 | 0.0022 | 0.0469 | 110 | 85 | 1.93E-3 | 6.37E-5 |
| 5 | 1.549 | NA | 0.0372 | NA | 110 | 1.53E-3 | 5.05E-5 |
| Hand Harvesting- Dry Beans Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.774 | 0.1295 | 0.1110 | 45 | 35 | 4.56E-3 | 1.50E-4 |
| 7 | 1.227 | 0.0573 | 0.0491 | 105 | 80 | 2.02E-3 | 6.67E-5 |
| 9 | 0.972 | NA | 0.0389 | NA | 100 | 1.60E-3 | 5.28E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.,

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 19. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Dry Beans applied at 2.5 lb ai/acre.

| Weeding/hoeing activities- Dry Beans Transfer Coefficient = 60.0 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.55 | 0.0051 | 0.0044 | 1200 | 915 | 1.79E-4 | 5.91E-6 |
| Irrigation/scouting activities- Dry Beans Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.55 | 0.1274 | 0.1092 | 45 | 35 | 4.49E-3 | 1.48E-4 |
| 7 | 2.012 | 0.0563 | 0.0483 | 105 | 85 | 1.98E-3 | 6.53E-5 |
| 9 | 1.595 | NA | 0.0383 | NA | 105 | 1.57E-3 | 5.18E-5 |
| Hand Harvesting- Dry Beans Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.550 | 0.2123 | 0.1820 | 30 | 20 | 7.48E-3 | 2.47E-4 |
| 11 | 1.263 | 0.0589 | 0.0505 | 100 | 80 | 2.08E-3 | 6.86E-5 |
| 13 | 1.000 | 0.0467 | 0.0400 | NA | 100 | 1.64E-3 | 5.41E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 20. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Citrus at 2.5 lb ai/acre.

| Irrigation/scouting activities - Citrus Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.214 | 0.0413 | 0.0354 | 145 | 115 | 1.46E-3 | 4.82E-5 |
| Pruning activities - Citrus Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.214 | 0.124 | 0.1063 | 50 | 40 | 4.70E-3 | 1.55E-4 |
| 10 | 1.192 | 0.0668 | 0.0572 | 90 | 70 | 2.35E-3 | 7.76E-5 |
| 12 | 1.054 | 0.0590 | 0.0506 | 100 | 80 | 2.08E-3 | 6.86E-5 |
| 16 | 0.823 | NA | 0.0395 | NA | 100 | 1.62E-3 | 5.35E-5 |
| Hand Harvesting - Citrus Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.214 | 0.3306 | 0.2834 | 20 | 15 | 1.16E-2 | 3.83E-4 |
| 10 | 1.192 | 0.1781 | 0.1526 | 35 | 25 | 6.27E-3 | 2.07E-4 |
| 12 | 1.054 | 0.1573 | 0.1349 | 40 | 30 | 5.54E-3 | 1.83E-4 |
| 16 | 0.823 | 0.1228 | 0.1053 | 50 | 40 | 4.33E-3 | 1.43E-4 |
| 28 | 0.391 | 0.0585 | 0.0501 | 102 | 80 | 2.06E-3 | 6.80E-5 |
| 32 | 0.306 | 0.0456 | 0.0391 | NA | 100 | 1.61E-3 | 5.31E-5 |

^a DAT = days after treatment.

^b Based on DFR (μg/cm²) data from a study of postapplication of Propargite (Ornite CR[®]) on navel oranges using an application rate of 3.15 lb ai/acre (MRID # 409090-03). Data normalized to an application rate of 2.5 lb ai/acre. (labeled application rate for the rest of crops within citrus group)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 21. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Citrus at 3.15 lb ai/acre.

| Irrigation/scouting activities - Citrus Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.789 | 0.0521 | 0.0446 | 115 | 90 | 1.83E-3 | 6.04E-5 |
| 2 | 2.465 | NA | 0.0394 | NA | 100 | 1.62E-3 | 5.35E-5 |
| Pruning activities - Citrus Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.789 | 0.1562 | 0.1339 | 40 | 30 | 5.50E-3 | 1.82E-4 |
| 16 | 1.036 | 0.0193 | 0.0497 | 105 | 80 | 2.04E-3 | 6.73E-5 |
| 20 | 0.809 | NA | 0.0388 | NA | 105 | 1.60E-3 | 5.28E-5 |
| Hand Harvesting - Citrus Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.789 | 0.4165 | 0.3570 | 15 | 10 | 1.47E-2 | 4.85E-4 |
| 16 | 1.036 | 0.1548 | 0.1327 | 40 | 30 | 5.45E-3 | 1.81E-4 |
| 32 | 0.385 | 0.0575 | 0.0493 | 105 | 80 | 2.03E-3 | 6.70E-5 |
| 36 | 0.301 | NA | 0.0385 | NA | 105 | 1.58E-3 | 5.21E-5 |

^a DAT = days after treatment.

^b Based on DFR (μg/cm²) data from a study of postapplication of Propargite (Omite CR[®]) on navel oranges using an application rate of 3.15 lb ai/acre (MRID # 409090-03). Data normalized to an application rate of 2.5 lb ai/acre. (labeled application rate for the rest of crops within citrus group)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 22. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Citrus at 4.5 lb ai/acre.

| Irrigation/scouting activities- Citrus Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.985 | 0.0744 | 0.0638 | 80 | 65 | 2.62E-3 | 8.65E-5 |
| 4 | 3.111 | 0.0581 | 0.0498 | 105 | 80 | 2.05E-3 | 6.77E-5 |
| 8 | 2.429 | NA | 0.0389 | NA | 100 | 1.60E-3 | 5.28E-5 |
| Pruning activities- Citrus Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.985 | 0.2231 | 0.1913 | 25 | 20 | 7.86E-3 | 2.59E-4 |
| 21 | 3.111 | 0.0608 | 0.0522 | 100 | 75 | 2.14E-3 | 7.06E-5 |
| 26 | 2.429 | NA | 0.0383 | NA | 105 | 1.57E-3 | 5.18E-5 |
| Hand Harvesting-Citrus Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.985 | 0.5950 | 0.5100 | 10 | 8 | 2.10E-2 | 6.93E-4 |
| 37 | 0.404 | 0.0603 | 0.0517 | 100 | 80 | 2.12E-3 | 7.00E-5 |
| 42 | 0.296 | NA | 0.0378 | NA | 105 | 1.56E-3 | 5.15E-5 |

^a DAT = days after treatment.

^b Based on DFR (μg/cm²) data from a study of postapplication of Propargite (Ornite CR[®]) on navel oranges using an application rate of 3.15 lb ai/acre (MRID # 409090-30). Data normalized to an application rate of 2.5 lb ai/acre. (labeled application rate for the rest of crops within citrus group)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 23. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants at 0.5 lb ai/acre.

| Irrigation/scouting/Hand weeding activities-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.300 | 0.0051 | 0.0044 | 1170 | 910 | 1.81E-4 | 5.97E-6 |
| Pruning activities -Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0300 | 0.0154 | 0.0132 | 1300 | 300 | 5.42E-4 | 1.79E-5 |
| Hand Harvesting-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0300 | 0.0410 | 0.0351 | 150 | 115 | 1.44E-3 | 4.75E-5 |

^a DAT = days after treatment
^b Based on DFR data from a study of postapplication of Propargite (Omite 30 w[®]) on apple using an application rate of 3.6 ai/acre (MRID # 409090-04).
^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].
^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.
^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.
^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.
^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 24. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Pome Fruits, Stone Fruits, Tropical and Subtropical Fruits and Ornamental plants at 1.5 lb ai/acre.

| Irrigation/scouting/Hand weeding activities-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^e (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.80 | 0.0154 | 0.0132 | 390 | 300 | 5.42E-4 | 1.79E-5 |
| Pruning activities-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^e (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.80 | 0.0461 | 0.0395 | 130 | 100 | 1.63E-3 | 5.38E-5 |
| Hand Harvesting-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^e (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.80 | 0.1230 | 0.1054 | 50 | 40 | 4.33E-3 | 1.43E-4 |
| 7 | 0.39 | 0.0588 | 0.0504 | 100 | 80 | 2.07E-3 | 6.83E-5 |
| 9 | 0.32 | NA | 0.0409 | NA | 100 | 1.68E-3 | 5.54E-5 |

^a DAT = days after treatment

^b Based on DFR data from a study of postapplication of Propargite (Omite 30 w[®]) on apple using an application rate of 3.6 ai/acre (MRID # 409090-04).

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 25. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants at 2.5 lb ai/acre.

| Irrigation/scouting/Hand weeding activities-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.4 | 0.0256 | 0.0220 | 235 | 180 | 9.03E-4 | 2.98E-5 |
| Pruning activities -Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.4 | 0.0769 | 0.0659 | 80 | 60 | 2.7E-3 | 8.91E-5 |
| 3 | 1.11 | 0.0623 | 0.0480 | 105 | 85 | 1.97E-3 | 6.50E-5 |
| 5 | 0.81 | NA | 0.0389 | NA | 100 | 1.60E-3 | 5.28E-5 |
| Hand Harvesting-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.4 | 0.2050 | 0.1757 | 30 | 25 | 7.22E-3 | 2.38E-4 |
| 12 | 0.39 | 0.0579 | 0.0496 | 105 | 80 | 2.04E-3 | 6.73E-5 |
| 14 | 0.31 | NA | 0.0402 | NA | 100 | 1.65E-3 | 5.45E-5 |

^a DAT = days after treatment

^b Based on DFR data from a study of postapplication of Propargite (Omite 30 w[®]) on apple using an application rate of 3.6 ai/acre (MRID # 409090-04).

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 26. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants at 4.5 lb ai/acre.

| Irrigation/scouting/Hand weeding activities-Pome Fruits, Stone Fruits, Tropical and Subtropical Fruits and Ornamental plants Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.5 | 0.0461 | 0.0395 | 130 | 100 | 1.63E-3 | 5.38E-5 |
| Pruning activities -Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 3000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.5 | 0.1384 | 0.1186 | 45 | 35 | 4.88E-3 | 1.61E-4 |
| 8 | 1.06 | 0.0596 | 0.0511 | 100 | 80 | 2.10E-3 | 6.93E-5 |
| 11 | 0.78 | NA | 0.0372 | NA | 105 | 1.53E-3 | 5.05E-5 |
| Hand Harvesting-Pome Fruits, Stone Fruits, Tropical & Subtropical Fruits and Ornamental plants Transfer Coefficient = 8000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 2.5 | 0.3691 | 0.3163 | 20 | 15 | 1.3E-2 | 4.29E-4 |
| 18 | 0.37 | 0.0544 | 0.0475 | 110 | 85 | 1.95E-3 | 6.44E-5 |
| 20 | 0.30 | NA | 0.0385 | NA | 105 | 1.58E-3 | 5.21E-5 |

^a DAT = days after treatment
^b Based on DFR data from a study of postapplication of Propargite (Omite 30 w[®]) on apple using an application rate of 3.6 ai/acre (MRID # 409090-04).
^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 mg) x ET (hrs) / BW (kg)].
^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.
^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.
^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]. , where adult body weight = 70 kg.
^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 27. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Berries at 2.0 lb ai/acre.

| Irrigation/Hand weeding activities-Berries Transfer Coefficient = 500 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.9 | 0.0084 | 0.0072 | 720 | 560 | 2.95E-4 | 9.74E-6 |
| Scouting activities-Berries Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.9 | 0.0167 | 0.0143 | 360 | 280 | 5.90E-4 | 1.95E-5 |
| Harvesting/Pruning/Tying activities-Berries Transfer Coefficient = 5000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 0.9 | 0.0837 | 0.0717 | 70 | 55 | 2.95E-3 | 9.74E-5 |
| 6 | 0.62 | 0.0577 | 0.0495 | 105 | 80 | 2.03E-3 | 6.70E-5 |
| 10 | 0.48 | NA | 0.0386 | NA | 105 | 1.59E-3 | 5.25E-5 |

^a DAT = days after treatment
^b Based on DFR data from a study of postapplication of Propargite (Ornite 6E[®]) on grape using an application rate of 2.88 ai/acre (MRID # 418486-03).
^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].
^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.
^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.
^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]. , where adult body weight = 70 kg.
^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 28. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for grapes at 3.0 lb ai/acre.

| Irrigation/Hand weeding activities-grapes Transfer Coefficient = 500 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.3 | 0.0126 | 0.0108 | 475 | 370 | 4.42E-04 | 1.46E-5 |
| Scouting activities-grapes Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.3 | 0.0251 | 0.0215 | 240 | 215 | 8.84E-04 | 2.92E-5 |
| Harvesting/Pruning activities-grapes Transfer Coefficient = 5000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.3 | 0.1255 | 0.1076 | 50 | 35 | 4.42E-3 | 1.46E-4 |
| 12 | 0.64 | 0.1195 | 0.0500 | 100 | 80 | 2.10E-3 | 6.93E-5 |
| 16 | 0.50 | NA | 0.0400 | NA | 100 | 1.64E-3 | 5.41E-5 |
| Cane turning activities-grapes Transfer Coefficient = 10000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.3 | 0.2511 | 0.2152 | 25 | 20 | 8.84E-3 | 2.92E-4 |
| 23 | 0.32 | 0.0605 | 0.0519 | 100 | 75 | 2.13E-3 | 7.03E-5 |
| 27 | 0.25 | NA | 0.0405 | NA | 100 | 1.66E-3 | 5.48E-5 |

^a DAT = days after treatment

^b Based on DFR data from a study of postapplication of Propargite (Ormite 6 E[®]) on grape using an application rate of 2.88 ai/acre (MRID # 418486-03)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.,

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 29. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Tree nuts at 3.0 lb ai/acre.

| Tree Shaking activities-Tree nuts Transfer Coefficient = 48 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.10 | 0.0027 | 0.0023 | 2235 | 1740 | 9.46E-5 | 3.12E-6 |
| Scouting/Weeding/Irrigation activities-Tree nuts Transfer Coefficient = 500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.1 | 0.0286 | 0.0245 | 210 | 165 | 1.01E-3 | 3.33E-5 |
| Hand Harvesting/Pruning activities-Tree nuts Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 3.10 | 0.1428 | 0.1224 | 40 | 35 | 5.03E-3 | 1.66E-4 |
| 17 | 1.28 | 0.0597 | 0.0512 | 100 | 80 | 2.10E-3 | 6.93E-5 |
| 22 | 0.99 | NA | 0.0396 | NA | 100 | 1.63E-3 | 5.38E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication propargite residues on almond using an application rate of 3.0 lb ai/acre (MRID # 418486-03)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day)⁻¹, where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 30. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Tree nuts at 4.5 lb ai/acre.

| Tree Shaking activities-Tree nuts Transfer Coefficient = 48 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.6 | 0.0040 | 0.0035 | 1500 | 1160 | 1.42E-4 | 4.69E-6 |
| Scouting/Weeding/Irrigation activities-Tree nuts Transfer Coefficient = 500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.6 | 0.0428 | 0.0367 | 140 | 110 | 1.51E-3 | 4.98E-5 |
| Hand Harvesting/Pruning activities-Tree nuts Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.6 | 0.2142 | 0.1836 | 30 | 20 | 8.80E-3 | 2.90E-4 |
| 25 | 1.27 | 0.0594 | 0.0509 | 100 | 80 | 2.09E-3 | 6.90E-5 |
| 30 | 0.99 | NA | 0.0394 | NA | 100 | 1.62E-3 | 5.35E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication propargite residues on almond using an application rate of 3.0 lb ai/acre (MRID # 418486-03)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr], where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day)⁻¹, where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 31. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Cereal Grains at 2.5 lb ai/acre.

| Irrigation/ Scouting activities-Cereal grains Transfer Coefficient = 1000 cm ² /hr | | | | | | | |
|---|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.8 | 0.0330 | 0.0282 | 180 | 140 | 1.16E-3 | 3.83E-5 |
| Hand Harvesting/ Detasseling activities-Cereal grains Transfer Coefficient = 17000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.8 | 0.5602 | 0.4802 | 10 | 8 | 1.97E-2 | 6.50E-4 |
| 11 | 0.20 | 0.0631 | 0.0541 | 100 | 75 | 2.22E-3 | 7.33E-5 |
| 13 | 0.13 | NA | 0.0364 | NA | 110 | 1.50E-3 | 4.95E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication of Propargite (Cornite EC[®]) on corn using an application rate of 2.46b ai/acre (MRID # 416803-02)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]. , where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 32. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Hops at 2.5 lb ai/acre.

| Irrigation/Scouting/Hand weeding activities-Hops Transfer Coefficient = 100 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.9 | 0.0091 | 0.0078 | 660 | 515 | 3.21E-4 | 1.06E-5 |
| Harvesting/Trailing activities-Cereal grains Transfer Coefficient = 2000 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.9 | 0.1820 | 0.1560 | 35 | 25 | 6.41E-3 | 2.12E-4 |
| 27 | 1.62 | 0.0605 | 0.0518 | 100 | 75 | 2.13E-3 | 7.03E-5 |
| 33 | 1.27 | NA | 0.0406 | NA | 100 | 1.67E-3 | 5.51E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication propargite residues (Ornite CR) on hops using an application rate of 1.35 lb ai/acre (MRID # 413996-01)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 33. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Mint at 2.5 lb ai/acre.

| All activities-Mint Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b ($\mu\text{g}/\text{cm}^2$) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 4.624 | 0.1295 | 0.1110 | 45 | 35 | 4.56E-3 | 1.50E-4 |
| 7 | 2.045 | 0.0573 | 0.0491 | 105 | 80 | 2.02E-3 | 6.67E-5 |
| 9 | 1.620 | NA | 0.0389 | NA | 105 | 1.60E-3 | 5.28E-5 |

^a DAT = days after treatment.

^b Based on DFR data from a study of postapplication Propargite residues on dry-beans using an application rate of 2.46 lb ai/acre (MRID #426891-04)

^c Dose (mg/kg/day) = [DFR ($\mu\text{g}/\text{cm}^2$) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR ($\mu\text{g}/\text{cm}^2$) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.,

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

Table 34. Propargite Short-Term and Intermediate-Term Occupational Postapplication Assessment for Oil Seed Crop at 1.5 lb ai/acre.

| Weeding and Hoeing activities-Cotton Transfer Coefficient = 63 cm ² /hr | | | | | | | |
|--|---|---|---------|-----------------------------|------------------------------------|-------------------|---------------------|
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.8 | 0.0022 | 0.0019 | 2760 | 2145 | 7.66E-5 | 2.53E-6 |
| Irrigation/Scouting activities- Oil Seed crop Transfer Coefficient = 1500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.8 | 0.0518 | 0.0444 | 115 | 90 | 1.82E-3 | 6.01E-5 |
| 1 | 1.65 | NA | 0.0395 | NA | 100 | 1.62E-3 | 5.35E-5 |
| Harvesting activities-Cotton Transfer Coefficient = 2500 cm ² /hr | | | | | | | |
| DAT ^a | DFR ^b (μg/cm ²) | Dermal Dose ^c (mg/kg/day) | | Short-term MOE ^d | Intermediate-term MOE ^e | LADD ^f | Cancer ^g |
| | | BW (60) | BW (70) | | | | |
| 0 | 1.8 | 0.0863 | 0.0740 | 70 | 55 | 3.04E-3 | 1.00E-4 |
| 3 | 1.30 | 0.0608 | 0.0522 | 100 | 75 | 2.14E-4 | 7.06E-6 |
| 6 | 0.92 | NA | 0.0368 | NA | 110 | 1.51E-3 | 4.98E-5 |

^a DAT = days after treatment

^b Based on DFR data from a study of postapplication of Propargite (Cornite EC ®) on cotton using an application rate of 1.64 ai/acre (MRID # 414578-06)

^c Dose (mg/kg/day) = [DFR (μg/cm²) x TC (cm²/hr) x CF (1 mg/1,000 μg) x ET (hrs) / BW (kg)].

^d Short-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 6 mg/kg/day.

^e Intermediate-term MOE = NOAEL (mg/kg/day) / Dose (mg/kg/day); where NOAEL = 4 mg/kg/day.

^f For agricultural scenarios, LADD = [DFR (μg/cm²) x Tc (cm²/hr) x mg/1,000 μg x hours exposed/day x exposure days/year x years of exposure x dermal absorption factor] / [body weight in kg x 70 yr x 365 days/yr]., where adult body weight = 70 kg.

^g Cancer Risk = LADD (mg/kg/day) x Q1* (mg/kg/day), where Q1* = 3.3x10⁻² (mg/kg/day)⁻¹

4.3.3 Incident Information

PROPARGITE REVIEW

Incident Data System

The following cases from the OPP Incident Data System (IDS) do not have documentation confirming exposure or health effects unless otherwise noted.

Incident#1280-23

A pesticide incident occurred in 1994, when a spray applicator got the chemical in his eyes. Specific symptoms were not mentioned. No further information on the disposition of the case was reported.

Incident#4066-12

A pesticide incident occurred in California in 1996, when 49 field workers lifted canes in grape fields that were wet with dew. Many of the workers clothes became soaking wet and they experienced burning, itching, and a rash on their arms, neck, chest, and stomach. From information collected by the Agricultural Commissioner's staff it appeared that the label was followed. There was evidence of non-compliance with the re-entry interval of 30 days. Results from analysis of foliage samples confirmed residues of propargite on the grape foliage. All of these workers were seen at the primary medical care center. No further information on the disposition of the case was reported.

Incident#5995-1

A pesticide incident occurred in 1997, when five workers experienced skin and eye irritations after formulating and packaging a chemical which was caused by abnormally high levels of dust being generated in the pack-room. No further information on the disposition of the case was reported.

Incident#7346-1

A pesticide incident occurred in 1985, when a worker was inadvertently drenched with spray from an air blast sprayer used to treat a grape vineyard. The worker experienced vomiting within thirty minutes and later developed chronic asthma and other respiratory problems. These symptoms were not consistent with exposure to propargite and there may have been exposure to a second pesticide that was responsible for these symptoms. No further information on the disposition of the case was reported.

Poison Control Center Data - 1993 through 1996

From 1993 through 1996 there were 62 exposures to propargite reported to Poison Control Centers participating in the Toxic Exposure Surveillance System. A total of 40 of these exposures were reported to be non-occupational including 33 adults and children six years old and over and seven children under age six. Twenty-two cases of exposure were reported to be occupationally related. Twenty-one of these cases occurred in California and therefore may also be reported in the section below concerning California data. No detailed analysis is performed because there were too few cases in any one category. Of the total cases 23 were reported to have a minor medical outcome and three cases were reported to have a moderate medical outcome. There were no fatalities or life-threatening cases. The most common symptoms reported included nausea, oral irritation, chest pain, dizziness, headache, and eye and dermal effects. A total of 25 of these cases were seen in a health care facility, however, none were admitted for hospitalization.

California Data - 1982 through 1996

Detailed descriptions of 923 cases submitted to the California Pesticide Illness Surveillance Program (1982-1996) were reviewed. In 671 of these cases, propargite was judged to be responsible for the health effects. Only cases with a definite, probable or possible relationship were reviewed. Propargite ranked 44th as a cause of systemic poisoning in California for the years 1982-1994. All of the systemic cases reported in this period were in an agricultural setting with roughly one-third occurring among handlers and two-thirds among field workers.

Table 30 presents the types of illnesses reported by year for the time period 1982 through 1996. Table 31 gives the total number of workers that took time off work as a result of their illness and how many were hospitalized and for how long.

Table 35. Cases Due to Propargite Exposure in California Reported by Type of Illness and Year, 1982-1996.

| Year | Illness Type | | | | Total |
|-------|-----------------------|-----|---------------------------|------|-------|
| | Systemic ^b | Eye | Combination. ^c | Skin | |
| 1982 | 2 | 9 | 2 | 40 | 53 |
| 1983 | 6 | 18 | 5 | 24 | 53 |
| 1984 | 3 | 13 | 4 | 63 | 83 |
| 1985 | 1 | 9 | - | 37 | 47 |
| 1986 | - | 7 | 1 | 143 | 151 |
| 1987 | 1 | 5 | 4 | 25 | 35 |
| 1988 | 3 | 7 | - | 81 | 91 |
| 1989 | 3 | 3 | - | 6 | 12 |
| 1990 | 5 | 4 | 1 | 7 | 17 |
| 1991 | - | 3 | - | 3 | 6 |
| 1992 | - | 5 | - | 15 | 20 |
| 1993 | 2 | 4 | - | 4 | 10 |
| 1994 | 3 | 2 | 1 | 5 | 11 |
| 1995 | 2 | - | - | 70 | 72 |
| 1996 | 2 | 2 | 1 | 5 | 10 |
| Total | 33 | 91 | 19 | 528 | 671 |

^b Category includes cases where skin, eye, or respiratory effects were also reported.

^c Category includes combined irritative effects to eye, skin, and respiratory system.

Table 36. Number of Persons Disabled (taking time off work) or Hospitalized for Indicated Number of Days After Propargite Exposure in California, 1982-1996.

| | Number of Persons Disabled | Number of Persons Hospitalized |
|-------------------|----------------------------|--------------------------------|
| One day | 55 | - |
| Two days | 25 | - |
| 3-5 days | 50 | - |
| 6-10 days | 18 | - |
| more than 10 days | 4 | - |
| Unknown | 161 | 5 |

A total of 528 persons had skin illnesses or 79% of 671 persons. Data covering the years 1982-1989 found that propargite was the leading cause of skin-related injuries among all pesticides. For the years 1990-1994, propargite dropped to seventh place among specific active ingredients. Worker activities associated with exposure to propargite are presented in Table 36 below.

Table 37. Illnesses by Activity Categories for Propargite Exposure in California, 1982-1996.

| Activity Category | Illness Category | | | | Total |
|-------------------|-----------------------|-----|------|--------------------------|-------|
| | Systemic ^b | Eye | Skin | Combination ^c | |
| Applicator | 7 | 45 | 64 | 10 | 126 |
| Mixer/Loader | 3 | 22 | 35 | 4 | 64 |
| Coincidental | 2 | 4 | 9 | 2 | 17 |
| Field Residue | 13 | 14 | 411 | 3 | 441 |
| Drift | 5 | - | 3 | - | 8 |
| Other | 2 | 6 | 7 | - | 15 |
| Total | 32 | 91 | 529 | 19 | 671 |

^a Coincidental=accidental exposure to application strength dilution but not directly involved in pesticide handling activity; Drift= exposure to pesticide that has drifted from intended targets.

^b Category includes cases where skin, eye, or respiratory effects were also reported

^c Category includes combined irritative effects to eye, skin, and respiratory system

According to the above activity categories, field residue was associated with the majority (66%) of the exposures. These illnesses included symptoms of chest tightness, shortness of breath, headache, sore throat, coughing, dermatitis, rash on arms, neck, chest and eyes, and eye irritation. In 1988, 26 workers harvesting nectarines developed rashes in orchards treated with propargite and two other pesticides. Samples of foliar dislodgeable residues suggested that propargite was the cause of the dermatitis cases.

Dermatitis developed in 114 orange pickers in a single incident in 1986. One-third of the workers developed peeling indicating severe dermatitis (Saunders et al. 1987). As a result of this and other large outbreaks the reentry interval was extended from 2-7 days (depending on crop) to 14-42 days in 1989 resulting in a significant reduction in propargite-related illness (Mehler et al., 1992).

National Pesticide Telecommunications Network

On the list of the top 200 chemicals for which NPTN received calls from 1984-1991 inclusively, propargite was ranked 116th with 28 incidents in humans reported and three incidents in animals (mostly pets).

According to California data, it appears that a majority of cases involved skin illnesses some of which can be quite severe requiring extensive time off work to recover. A large proportion of cases resulted from field reentry and worker activities involving extensive contact with treated foliage such as turning cane for grapes and harvesting citrus. Both eye and skin problems are commonly reported among applicators who handle propargite without proper protection.

5.0 AGGREGATE RISK ASSESSMENTS AND RISK CHARACTERIZATION

5.1 Acute Aggregate Risk

There are no registered residential uses of propargite so acute aggregation will include only food and water.

Acute DWLOCs were calculated based on the acute dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were less than the acute DWLOCs, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The acute DWLOC for Females 13-50 years is 2400 ppb. The PRZM-EXAMS surface water value is 34 ppb.

The EECs for groundwater (SCI-GROW) were less than the acute DWLOC's, indicating that acute aggregate exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). To calculate the DWLOC, the acute dietary food exposure was subtracted from the acute PAD using the equation:

$$\text{DWLOC}_{\text{acute}}(\text{ug/L}) = \frac{[\text{acute water exposure (mg/kg/day)} \times (\text{body weight in kg})]}{[\text{consumption (L/day)} \times 10^{-3} \text{ mg/} \mu\text{g}]}$$

where acute water exposure (mg/kg/day) = [aPAD - (acute food (mg/kg/day))]

Table 38. Drinking Water Levels of Comparison for Acute Dietary Exposure.

| Population Subgroup | Acute PAD (mg/kg/day) | Food Exposure (mg/kg/day) | Max. Water Exposure (mg/kg/day) | DWLOC _{acute} (ug/L) | PRZM-EXAMS (ug/L) | SCI-GROW (ug/L) |
|---------------------|-----------------------|---------------------------|---------------------------------|-------------------------------|-------------------|-----------------|
| Females 13-50 years | 0.08 | 0.001 | 0.08 | 2400 | 34 | 0.006 |

5.2 Short- and Intermediate-Term Aggregate Risks

There are no registered residential uses of propargite, therefore, short- and intermediate-term aggregation is not appropriate.

5.3 Chronic Aggregate Risk

There are no registered residential uses of propargite, therefore, chronic aggregation will include only food and water.

Chronic DWLOCs were calculated based on the chronic dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were less than the chronic DWLOCs, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the chronic DWLOC's, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The PRZM-EXAMS surface water value is 8.7 ppb. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). To calculate the chronic DWLOC, the chronic dietary food exposure was subtracted from the chronic PAD using the equation:

$$DWLOC_{chronic}(ug/L) = \frac{[chronic\ water\ exposure\ (mg/kg/day) \times (body\ weight\ in\ kg.)]}{[consumption\ (L/day) \times 10^{-3}\ mg/\mu g]}$$

where chronic water exposure (mg/kg/day) = [cPAD - (chronic food (mg/kg/day))]

Table 37. Drinking Water Levels of Comparison for Chronic Dietary Exposure.

| Population Subgroup | Chronic PAD (mg/kg/day) | Food Exposure (mg/kg/day) | Max. Water Exposure (mg/kg/day) | DWLOC _{chronic} (ug/L) | PRZM-EXAMS (ug/L) | SCI-GROW (ug/L) |
|---------------------|-------------------------|---------------------------|---------------------------------|---------------------------------|-------------------|-----------------|
| US Population | 0.04 | 0.00001 | 0.04 | 1400 | 8.7 | 0.006 |
| All Infants | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Children 1-6 | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Children 1-12 | 0.04 | 0.00001 | 0.04 | 400 | 8.7 | 0.006 |
| Females 13-50 yrs. | 0.04 | 0.00001 | 0.04 | 1200 | 8.7 | 0.006 |
| Males 20+ yrs | 0.04 | 0.00001 | 0.04 | 1400 | 8.7 | 0.006 |

5.4 Cancer Aggregate Risk

There are no registered residential uses of propargite so cancer aggregation will include only food and water.

A cancer DWLOC was calculated based on the cancer dietary (food) exposure and default body weights and water consumption figures. The EECs for surface water (PRZM-EXAMS) were greater than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is greater than HED's level of concern. The EECs for groundwater (SCI-GROW) were less than the cancer DWLOC, indicating that chronic exposure to propargite in food and water is less than HED's level of concern. The Agency's default body weights and water consumption values used to calculate DWLOCs are as follows: 70 kg/2L (adult male), 60 kg/2L (adult female), and 10 kg/1L (child). The PRZM-EXAMS and SCI-GROW EECs were 4.8 and 0.006 ppb, respectively. Surface water concentrations below 0.2 ppb would result in cancer risks below 1×10^{-6} for drinking water alone when back calculated. Time weighted average propargite concentration in surface water samples from the USGS NAWQA (Oristimba Creek Watershed) for the years 1992-1993 were 0.30 and 1.24 ppb, respectively. Therefore, even when monitoring data are used cancer exposure to propargite from surface water sources is greater than HED's level of concern.

Table 38. Summary of Cancer DWLOC Calculations for Propargite.

| Population | Q _i [*] | Negligible Risk Level ¹ | Target Max Exposure ² mg/kg/day | Chronic Food Exposure mg/kg/day | Max Water Exposure ³ mg/kg/day | Groundwater EEC (ug/L) | Surface Water EEC (ug/L) | Cancer DWLOC ⁴ (ug/L) |
|------------|-----------------------------|------------------------------------|---|------------------------------------|--|---------------------------|-----------------------------|-------------------------------------|
| U.S. Pop. | 0.33x10 ⁻¹ | 0.000001 | 0.0000303 | 0.00001 | 0.0000203 | 0.006 | 4.8 | 0.71 |

¹ DWLOC_{CANCER} was calculated for U.S. population only. Default body weight and consumption value for calculation of the DWLOC were: 2L/70 kg

² Target Maximum Exposure (mg/kg/day) = [negligible risk/Q*]

³ Maximum Water Exposure (mg/kg/day) = [Target Maximum Exposure - (Chronic Food Exposure + Residential Exposure (Lifetime Average Daily Dose))]

⁴ Cancer DWLOC(ug/L) = [maximum water exposure (mg/kg/day) x body weight (kg)]
[water consumption (L) x 10⁻³ mg/ug]²

6.0 DATA NEEDS

Additional data requirements have been identified in the attached Science Chapters and are summarized below.

Toxicology Data for OPPTS Guidelines: None required.

Product and Residue Chemistry Data for OPPTS Guidelines:

OPPTS GLN 830.7050 (UV/Visible absorption)

OPPTS GLN 860.1200 (Directions for Use) - Label revisions are required.

OPPTS GLN 860.1380 - Additional storage stability data are required for peanut, walnut, corn, and tea..

OPPTS GLN 860.1520 - Additional residue data are required for cotton gin byproducts.

Occupational Exposure Data for OPPTS Guidelines: None required.



13544

031352

Chemical: Propargite

PC Code: 097601

HED File Code 14000 Risk Reviews

Memo Date: 09/13/2001

File ID: DPD276544

Accession Number: 412-02-0006

HED Records Reference Center

12/10/2001