

BONIDE

1.00% ROTENONE

ACTIVE INGREDIENTS
Rotenone 1.00%
Other Chlorinated Hydrocarbons 2.00%
INERT INGREDIENTS 97.00%

EPA Reg. No. 4-17
1.00%
2.00%
97.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

(See back panel cautions)

1 LB. NET

1.00% ROTENONE

1.00% ROTENONE contains Toxic principles of Cube, combined with wetting agent and diluent to make an effective dust or spray. Recommended for general use on flowers, vines, plants, shrubs, raspberries, and vegetables, for controlling those insects against which Rotenone preparations are effective.

DIRECTIONS

Shake can to fluff powder. This package can be used as a dust or spray. Punch out one or more openings, press sides of container to form a cone for dusting motion.

FOR DUSTING: Apply 1/2 to 1 lb. per acre depending upon size of plants, crop and insect infestation.

FOR SPRAYING: Mix at the rate of 1 pound to 5 gallons of water. Spray as necessary. Apply thoroughly with a good pressure spray.

In all cases make repeat applications as often as necessary.

FOR USE ON BEANS: To control Mexican Bean Beetles and certain other insects, apply thoroughly, especially the underside of leaves as soon as the eggs begin to hatch.

FOR USE ON CABBAGE, CAULIFLOWER: To control Imported Cabbageworm, Diamond Back Moth, apply with a good duster to all parts of the plants as soon as they appear.

FOR USE ON POTATO AND EGGPLANTS: To control Colorado Potato Beetle, repel Potato Flea Beetle, apply as soon as insects appear. Repeat applications at 5 to 7 day intervals.

FOR USE ON CUCUMBER, MELON: To repel Cucumber Beetle, repel Flea Beetles, apply thoroughly to all parts of the plants as soon as they appear. Repeat applications and make repeat applications at 5 to 7 day intervals.

FOR USE ON TOMATOES: To repel Potato Flea Beetles and control Colorado Potato Beetle Larvae, apply 2 or 3 days after plants are set in the field. Repeat applications at 5 to 7 day intervals.

FOR USE ON FLOWERS, RASPBERRIES, SHRUBS: To control Flea Beetles, aphids, apply thoroughly as soon as insects appear.

Certain insects not controllable with Rotenone may be controlled with Bonide 5% Sevin Dust, Bonide Methoxychlor, 25% Miscible Concentrate.

CAUTION


Do not apply to food crops within 1 day of harvest.

Buyer's Guarantee Limited to Label Claims

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

4-17
3/20/72
P3182

282



~~ACCEPTED~~
 MAR 9 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

ACCEPTED
 MAR 20 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

4-17