



ACTIVE INGREDIENTS  
 Rotenone  
 Other Cilia Resins  
 Methyl Isocyanates  
 INERT INGREDIENTS

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 (See back panel cautions)

35%  
 65%  
 1.30%  
 97.70%

1 LB. NET

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS — CAUTION**  
 May be harmful if swallowed. Avoid breathing dust or spray. Avoid contact with eyes, skin, clothing. Wash thoroughly after use. In case of contact rinse with plenty of water.

**ENVIRONMENTAL HAZARDS**  
 This product is extremely toxic to fish and aquatic organisms. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply while bees are actively visiting the area. Additional protective information may be obtained from your Cooperative Extension Service.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Storage:** Store in a cool dry place which is inaccessible to children and pets.  
**Container Disposal:** Do not reuse empty container. Wrap container and put in trash collection.

**ROSYN** is used as dust or spray any time when conditions are favorable, applied at 7 to 10 day intervals, as soon as plants are above ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required (i.e., after rainfall, etc). Do not apply to food crops within one day of harvest.

**AS A DUST:** This can is a squeeze type duster. Shake can to fluff powder. Set top opening for desired amount (3 or 4 holes). Press sides of can with finger using rapid motion for fog effect. Cover foliage with a thin uniform film of dust, both top and undersides of leaves.

**AS A SPRAY:** Mix at the rate of 1 pound to 5 gallons of water or stronger if necessary (1 tablespoonful to 1 quart of water). Add powder to small quantity of water first making a paste, then add desired amount of water. Spray to cover all parts of exposed plant both top and undersides of leaves.

**CONTROLS** Mexican Bean Beetles and certain other insects on String Beans, Cabbage, Cauliflower, Eggplant, Broccoli, Brussels Sprouts, Potatoes, Tomatoes.

**FOR USE ON BEANS:** To control Mexican bean beetles and certain other leaf eating insects.

**FOR USE ON CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS:** To control imported cabbage worm and diamond back moth.

**FOR USE ON POTATOES AND EGGPLANTS:** To control Colorado potato beetle larvae and to repel potato flea beetle.

**FOR USE ON TOMATOES:** To repel potato flea beetle and control Colorado potato beetle larvae.

EPA Est. 4-NY-1 EPA Reg. No. 4-21

Buyer's Guarantee Limited to Label Claims  
 Mfg. by BONIDE CHEMICAL CO., INC., Yorkville, N.Y. 13495

105/1  
 pm 17  
 4.51