

leg # 4-21

PM-18

172

DEC 28 1991

Mr. James Wurz
Bonide Chemical Company, Inc.
2 Wurz Avenue
Yorkville, NY 13495

Dear Mr. Wurz:

Subject: Bonide Rotosyn Insect Dust
EPA Registration No. 4-21
Confidential Statement of Formula and Label Dated April 14,
1989 Submitted in Response to Registration Standard
Requirements

The label is acceptable provided that you submit five (5) copies of your final printed labeling before you release the product for shipment using this label.

A stamped copy of the labeling is enclosed for your records.

The Confidential Statement of Formula dated April 14, 1989 must be revised to include certified limits for the active ingredients and deliberately added inert ingredients. Further revisions will be required if changes are made in labels for Manufacturing Use Concentrates supplying the active ingredients for this product.

Sincerely yours,




Phil Hutton
Product Manager (18)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

52342:I:Schroeder:C.Disk:KEVRIC:06/11/91:CL:WO:CL

CONCURRENCES

SYMBOL							
SURNAME							
DATE							



BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS — CAUTION
 May be harmful if swallowed. Avoid breathing dust or spray. Avoid contact with eyes, skin, clothing. Wash thoroughly after use. In case of contact, rinse with plenty of water.

ENVIRONMENTAL HAZARDS
 This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage: Store in a cool dry place which is inaccessible to children and pets
Container Disposal: Do not reuse empty container. Wrap container and put in trash collection.

ROTSYN is used as dust or spray any time when conditions are favorable, applied at 7 to 10 day intervals, as soon as plants are above ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required (i.e., after rainfall, etc). Do not apply to food crops within one day of harvest.

AS A DUST: This can is a squeeze type duster. Shake can to fluff powder. Set top opening for desired amount (3 or 4 holes). Firm sides of can with finger using rapid motion for fog effect. Cover foliage with a thin uniform film of dust both top and undersides of leaves.

AS A SPRAY: Mix at the rate of 1 pound to 5 gallons of water or stronger if necessary (1 tablespoonful to 1 quart of water). Add powder to small quantity of water first making a paste, then add desired amount of water. Spray to cover all parts of exposed plant both top and undersides of leaves.

CONTROLS Mexican Bean Beetles and certain other insects on String Beans, Cabbage, Cauliflower, Eggplant, Broccoli, Brussels Sprouts, Potatoes, Tomatoes.

FOR USE ON BEANS: To control Mexican bean beetles and certain other leaf eating insects.

FOR USE ON CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: To control imported cabbage worm and diamond back moth.

FOR USE ON POTATOES AND EGGPLANTS: To control Colorado potato beetle larvae and to repel potato flea beetle.

FOR USE ON TOMATOES: To repel potato flea beetle and control Colorado potato beetle larvae.

EPA Est. 4 NY-1 EPA Reg. No. 4-21

Buyer's Guarantee Limited to Label Claims
 Mfg. by **BONIDE CHEMICAL CO., INC.**, Yorkville, N.Y. 13495

Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the purpose of the PPA, 4-21

DEC 28 1981

AC... with COMED... in EPA Letter Da.