

BONIDE

REG. U.S. PAT. OFF.

# BONTOX

VEGETABLE-FLORAL  
DUST or SPRAY



ACTIVE INGREDIENTS:	
Rotenone .....	.50%
Other Cube Resins .....	1.00%
Copper (In Basic Copper Sulphate) Expressed as Metallic .....	7.00%
Methyl Naphthalenes .....	.85%
Dichloro Diphenyl Trichloroethane .....	3.00%
INERT INGREDIENTS	
	87.65%
	100.00%

ACCEPTED  
 Feb. 20, 1961  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND ROENTICIDE ACT  
 OF ECONOMIC POISON REGISTRATION  
 UNDER NO. 4-29 SUB F  
 ATTACHED COMMENTS.

### CONTROLS

BONTOX is a micronized insecticide and fungicide especially recommended to control insects; such as flea beetles, leaf hoppers, Colorado potato beetles, tarnished plant bugs, aphids, tomato fruit worm, Mexican bean beetle, Japanese beetles, roller worms, rose chafer, tent caterpillars and diseases; such as early and late blights, certain leaf spots, downy mildew on vegetables such as potatoes, egg plants, beans and flowers; such as roses, gladiolus, mums, shrubs.

### DIRECTIONS FOR HOME USE



AS A DUST: Shake can to fluff powder, open 3 or 4 holes in top. Press sides of can with fingers rapidly for dust effect. Cover both tops and undersides of leaves with thin uniform film of dust.

AS A SPRAY: Using hand or tank sprayers: Mix 4 tablespoons per gallon of finished spray. Start by making gravy paste (adding a little water to the powder). Strain, then add rest of water: This in addition to keeping solution agitated will prevent nozzle-plugging. Spray thoroughly both tops and undersides of leaves until lightly dripping.

Repeat treatment as required, usually every 7 to 10 days. It is advisable to start dusting or spraying before insects and diseases appear as it is easier to prevent than cure certain diseases.

### CAUTION

Avoid inhalation, contamination of food stuffs and store out of reach of children. Do not apply to green or snap beans after the pods begin to form nor to food crops within 30 days of harvest; where control of insects and diseases closer to harvest is necessary switch to non-residual Bonide Garden Dust or for diseases only Bonide Copper Dust. BONTOX may be applied later to tomatoes up to 5 days before harvest if residue at harvest is removed by washing, wiping or other effective means. Poison if taken internally.

14 OZ. NET

MADE IN U.S.A.  
 BONIDE CHEMICAL CO., UTICA, N.Y.

4-29

62/20/1961

111