

BONIDE

REG. U.S. PAT. OFF.

BONTOX

VEGETABLE-FLORAL
DUST or SPRAY

REPTED
7-6-71

4-29

I
M
P
R
O
V
E
D

CONTAINS
SEVIN



ACTIVE INGREDIENTS:	USDA Reg. No. 4-29
*Carbaryl (1-naphthyl N-methylcarbamate)	1.25%
Rotenone50%
Other Cube Resins	1.00%
Copper (In Basic Copper Sulphate) Expressed as Metallic	7.00%
Methyl Naphthalenes85%
INERT INGREDIENTS	89.40%
*SEVIN REG. T.M. OF U.C.C.	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions) 4 LBS. NET

(READ THE LABEL BEFORE USING)

BONTOX

CONTROLS

"BONTOX" is a micronized insecticide and fungicide especially recommended to control flea beetles, leaf hoppers, Colorado potato beetles, tarnished plant bugs, aphid, tomato fruit worm, Mexican bean beetles, Japanese beetles, roller worms, rose chafer, tent caterpillars on tomatoes, potatoes, beans, roses, gladiolus, mums, shrubs, and early and late blights and downy mildew on potatoes, beans and tomatoes.

DIRECTIONS FOR HOME USE

AS A DUST: Shake can to fluff powder, open 3 or 4 holes in top. Press sides of can with fingers rapidly for dust effect. Cover both tops and undersides of leaves with thin uniform film of dust.

AS A SPRAY: Using hand or tank sprayers: Mix 4 tablespoons per gallon of finished spray. Start by making gravy paste (adding a little water to the powder). Strain, then add rest of water: This in addition to keeping solution agitated will prevent nozzle-plugging. Spray thoroughly both tops and undersides of leaves until lightly dripping.

Repeat treatment as required, usually every 7 to 10 days. It is advisable to start dusting or spraying before insects and diseases appear as it is easier to prevent than cure certain diseases.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using.

Avoid storage near food or feed products.

Do not apply to any of the named food crops within 1 day of harvest; where control of insects and diseases closer to harvest is necessary, switch to non-residual Bonide Garden Dust or for diseases only, Bonide Copper Dust. BONTOX may be applied later to tomatoes up to 1 day before harvest if residue at harvest is removed by washing, wiping or other effective means.

To protect honey bees, avoid use, if possible, during periods when bees are visiting the crop. If necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Buyer's Guarantee Limited to Label Claims

Manufactured by
BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

4-29

07/06/71

1/1