

OCT 17 1985

Annabel B. Olson
Bonide Chemical Co.
2 Wurz Avenue
Yorkville, NY 13495

Dear Ms. Olson:

Subject: Carbaryl Labeling Requirements
Bonide Rose and Floral Spray Bomb
EPA Registration No. 4-186
Bonide Vegetable-Floral Dust and Spray
EPA Registration No. 4-29
Your Letters of August 30,
and September 3, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

Include the following Bee Caution as part of the environmental precautionary paragraph.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

39422:Ellenberger:158417, 158416:RD-85/12;KENCO:10/9/85;Del. 10/22/85:eg

CONCURRENCES

SYMBOL ▶	158417	12						
SURNAME ▶	03	300						
DATE ▶								

BONIDE

VEGETABLE-FLORAL DUST OR SPRAY

IMPROVED CONTAINS SEVIN

ACTIVE INGREDIENTS:

	EPA Est. No. 4-NY-1	EPA Reg. No. 4-29
*Carbaryl (1-naphthyl N-methylcarbamate)		1.25%
Rotenone50%
Other Cube Resins		1.00%
Copper (In Basic Copper Sulphate) Expressed as Metallic		7.00%
Methyl Naphthalenes85%
INERT INGREDIENTS:		89.40%
		100.00%

*Sevin, Reg. T.M. of U.C.C.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statements.

NET WEIGHT: 14 Ounces

NET WEIGHT 4 LB.

NET WEIGHT 1 LB.

in letter case

OCT 17 1985

4-29

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water. Do not induce vomiting unless directed to do so by a physician. Remove person to fresh air.
If inhaled: Remove person to fresh air.
If in eyes: Flush with plenty of water. Get immediate attention.
If on skin: Remove contaminated clothing and wash affected areas.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to bees and aquatic organisms. Do not apply to or contaminate water by cleaning of equipment or disposal of wastes. Avoid visiting the area. Additional protective information may be obtained from:

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with these directions.

Storage: Keep pesticide in original container. Avoid contamination of food, feed, or dry place, preferably in a locked storage area.
Product Disposal: Securely wrap original container in several layers of newspaper. Do not reuse bag. Discard bag in trash.

(READ THE LABEL BEFORE USE)

BONIDE VEGETABLE-FLORAL DUST OR SPRAY is a contact insecticide and fungicide especially recommended to control Colorado potato beetles, tarnished plant bugs, Mexican bean beetles, Japanese beetles, roller caterpillars on tomatoes, potatoes, beans, roses, and early and late blights and downy mildew on tomatoes.

DIRECTIONS FOR HOME USE

AS A DUST: Shake can to fluff powder, open 3 or 4 inches and use with fingers rapidly for dust effect. Cover both sides of leaves with thin uniform film of dust.

AS A SPRAY: Using hand or tank sprayers: Mix 4 ounces of finished spray. Start by making gravy paste (add 1/2 cup of water to 1/2 cup of powder). Strain, then add rest of water. This in addition to agitation will prevent nozzle-plugging. Spray thoroughly on both sides of leaves until lightly dripping.

Repeat treatment as required, usually every 7 to 10 days. Start dusting or spraying before insects and diseases appear to prevent than cure certain diseases.

Do not apply to any of the named food crops within 14 days of harvest. Control of insects and diseases closer to harvest with non-residual Bonide Garden Dust or for diseases use Bonide VEGETABLE-FLORAL DUST OR SPRAY may be applied up to 1 day before harvest if residue at harvest is removed by wiping or other effective means.

Buyer's Guarantee Limited to Label Claims

© 1973 & Mfg. by BONIDE CHEMICAL CO., INC.