

PM04

4-29

7/6/98

1/4

ACCEPTED
JUL 6 1998
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-29

front panel

BONIDE
VEGETABLE - FLORAL
DUST OR SPRAY

EPA Est. No. 4-NY-1

EPA Reg. No. 4-29

ACTIVE INGREDIENTS:

* Carbaryl (CAS # 63-25-2).....	1.25%
Rotenone.....	.50%
Other Cube Resins.....	1.00%
Copper (in Basic Copper Sulphate) expressed as Metallic.....	7.00%
OTHER INGREDIENTS.....	90.25%
TOTAL.....	100.00%

* Sevin Reg. T.M. of Rhone Poulenc

KEEP OUT OF REACH OF CHILDREN
CAUTION
<u>See back panel for additional precautionary statements</u>

NET CONTENTS: _____

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION- Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wash thoroughly after handling. Wash contaminated clothing before reuse. May cause skin sensitization in certain individuals.

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.

Wear a hat and eye protection when making overhead applications.

Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.

Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Do not allow children or pets to contact treated area until sprays have dried or dust has settled.

STATEMENT OF PRACICAL TREATMENT

If Swallowed: Call a Physician or Poison Control Center. Drink (1) or (2) glasses of water and induce vomiting by touching finger to back of throat.

If inhaled: Remove person to fresh ait. . .

If in eyes: Flush with plenty of water. Get immediate attention.

If on skin: Remove contained clothing and wash affected areas with soap and water.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Additional protective information may be obtained from your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS

STORAGE AND DISPOSAL

Storage: Keep pesticide in original container. Avoid contamination of feed and foodstuffs. Store in a cool dry place, preferably in a locked storage area.

Product Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. Discard container in trash.

Bonide Vegetable-Floral Dust or Spray- To control certain insects and to control or prevent certain blight, leaf spots, mildew and certain other diseases on listed plants. Application for certain insects should be made as soon as insects appear. Applications against certain disease should begin well before they are expected to appear.

Use as a dust or spray any time when conditions are favorable, applied at 7 day intervals, as soon as plants are above the ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required, but not more often than once a week.

When dusting wear household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin. For plants above chest height consider using a liquid carbaryl product.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with dusted or wet surfaces. Allow spray to dry or dust to settle in treated areas before reentering.

Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. For trees taller than 10 feet, consider hiring a licensed professional.

Do not repeat applications more than once a week.

As a Dust: This can is a squeeze type duster. Shake can to fluff powder. Press sides of can with fingers, using rapid motion for fog effect. Depending on foliage density, one pound should cover approximately 1,000 sq.ft.

As a Spray: Mix 1 pound to 5 gallons of water, or 1 tablespoonful to 1 quart of water. Add powder to small quantity of water first making a paste, then add desired amount of water.

When to Apply: For insects apply when insects appear and continue application weekly. Young insects and larvae are usually easier to control than adults. When blossoms are in full bloom, protect pollinators by using a fungicide only, such as Bonide Copper Dust. For Disease, applications should begin well before they are expected to appear. Repeat weekly and after rains.

(PHI--Preharvest Interval)

CROP	DISEASES	INSECTS CONTROLLED OR REPELLED
Beans-Green Lima, Snap	Bacterial Blight Downy mildew	Mexican bean beetle, flea beetle, leafhopper, stink bug, tarnished plant bug, cutworm, armyworm (3) day PHI
Cabbage, Broccoli, Cauliflower Brussel sprout	Downy mildew, leaf spot	Imported cabbage worm, cabbage looper, diamond back moth, aphid. (3) day PHI
Carrots	Downy mildew, leaf spot	Flea beetle, thrips, aphid <u>carrots rust fly. (7)day PHI</u>
Celery	Early and late blight, leaf spot	Parsnip webworm, two-spotted spider mite, tarnished plant bug, carrot rust fly, aphid. (14) day PHI
Cucumber, Melons, Pumpkins, Squash	Angular leaf spot, downy mildew, powdery mildew, gummy stem blight, anthracnose, leaf spots, scab	Pickleworm, melonworm, cucumber beetle, young squash bug, aphid, two-spotted mite (3) day PHI
Potatoes	Early and late blights	Colorado potato beetle, flea beetle, leafhoper, young tomato hornworm, tomato pinworm, tarnished plant bug, stink bug, aphid, thrips, white fly, mealybugs, Japanese beetle Potato-(7) day PHI
Tomatoes, Eggplant	Early and late blight, anthracnose, bacterial spot, leaf spots, nail head rust, Setptoria and Stemphylium leaf mold	Tomatos, Eggplant-(3) day PHI

4/6

page 5`

For information on pesticide products (include health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunication Network at 1-800-858-7378.

Buyers guarantee limited to label claims

Manufactured by:

Bonide Products, Inc.

Yorkville, NY 13495