




# Bonide® Vegetable – Floral Dust or Spray

[Additional Brand Names: Bonide® Tomato-Potato Dust; Bonide® Tomato-Potato Spray or Dust; Bonide® Copper Dragon]

Controls both insects and diseases on tomatoes and vegetables.



### ACTIVE INGREDIENTS:

Carbaryl (CAS #63-25-2) .....	1.25%
Rotenone .....	0.50%
Other Cube Resins .....	1.00%
Copper (in Basic Copper Sulfate) expressed as Metallic .....	7.00%

<b>OTHER INGREDIENTS</b> .....	<b>90.25%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See back/side/other panel(s) for additional Precautionary Statements.

**NET WEIGHT:** \_\_\_\_\_

**BONIDE® Vegetable – Floral Dust or Spray**  
For Residential Use Only

**FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**Note to Physician:** Contains N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoximine methiodide) are NOT recommended.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Do not contaminate water, food or feed with this product.

**USER SAFETY REQUIREMENTS**

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.

Wear a hat and eye protection when making overhead applications.

Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.

Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Do not allow children or pets to contact treated area until sprays have dried or dust has settled.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Additional protective information may be obtained from your Cooperative Extension Service.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For home garden use only. Tinted green to blend with foliage.

### STORAGE AND DISPOSAL

**Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

**Disposal:** *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**Bonide® Vegetable – Floral Dust or Spray** – To control listed insects and to control or prevent blight, leaf spots, mildew and other listed diseases on plants. Application for insects should be made as soon as insects appear. Applications against disease should begin well before they are expected to appear.

Use as a dust or spray any time when conditions are favorable, applied at 7 day intervals, as soon as plants are above the ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required, but not more often than once a week.

When dusting, wear household latex or rubber gloves. Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin. Do not use the powder formulation on plants that are higher than chest height.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with dusted or wet surfaces. Allow spray to dry or dust to settle in treated areas before re-entering.

Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. For trees taller than 10 feet, consider hiring a licensed professional.

Do not repeat applications more than once a week.

**As a Dust:** This can is a squeeze type duster. Shake can to fluff powder. Press sides of can with fingers, using rapid motion for fog effect. Depending on foliage density, one pound should cover approximately 1,000 sq. ft.

**As a Spray:** Mix 1 pound to 5 gallons of water, or 1 tablespoonful to 1 quart of water. Add powder to small quantity of water first making a paste, then add desired amount of water.

**When to Apply:** For insects apply when insects appear and continue application weekly. Young insects and larvae are usually easier to control than adults. When blossoms are in full bloom, protect pollinators by using a fungicide only, such as Bonide® Copper Dust. For disease, applications should begin well before they are expected to appear. Repeat weekly and after rains.

(PHI – Preharvest Interval)

<b>CROP</b>	<b>DISEASES</b>	<b>INSECTS CONTROLLED OR REPELLED</b>
Beans – Green, Lima, Snap <b>(3) day PHI</b>	Bacterial blight, downy mildew	Mexican bean beetle, flea beetle, leafhopper, stink bug, tarnished plant bug, cutworm, armyworm.
Cabbage, Broccoli, Cauliflower, Brussels Sprout <b>(3) day PHI</b>	Downy mildew, leaf spot	Imported cabbage worm, cabbage looper, diamond back moth, aphid.
Carrots <b>(7) day PHI</b>	Downy mildew, leaf spot	Flea beetle, thrips, aphid, carrot rust fly.
Celery <b>(14) day PHI</b>	Early and late blight, leaf spot	Parsnip webworm, two-spotted spider mite, tarnished plant bug, carrot rust fly, aphid.
Cucumber, Melons, Pumpkins, Squash <b>(3) day PHI</b>	Angular leaf spot, downy mildew, powdery mildew, gummy stem blight, anthracnose, leaf spots, scab	Pickleworm, melonworm, cucumber beetle, young squash bug, aphid, two-spotted mite.
Potatoes <b>(7) day PHI</b>	Early and late blights	Colorado potato beetle, flea beetle, leafhopper, young tomato hornworm, tomato pinworm, tarnished plant bug, stink bug, aphid, thrips, white fly, mealybugs, Japanese beetle.
Tomatoes, Eggplant <b>(3) day PHI</b>	Early and late blight, anthracnose, bacterial spot, leaf spots, nail head rust, Septoria and Stemphylium leaf mold	

CROP	DISEASES	INSECTS CONTROLLED OR REPELLED
Strawberries (7) day PHI	Leafy spot, downy mildew, leaf blight, leaf scorch	Leaf roller, weevil, red spider, spittlebug.
Apricots, Cherries (sour) (3) day PHI	Brown rot	Black cherry aphid, mealy plum aphid, cherry fruit fly, Japanese beetle, tent caterpillar, orange tortrix, tussock moth, plum curculio.
Pears (3) day PHI	Scab, leaf blight	Codling moth, plum curculio, aphid, leafhopper, pear psylla. <b>Do not use dust on trees. Use only liquid formulation.</b>
Arbor vitae, Asters	Blight, leaf spots	Red spider, flea beetle, thrips, Japanese beetle, leaf roller, birch leaf miner, lace bugs, tent caterpillar. <b>Do not use dust on trees. Use only liquid formulation.</b>
Oaks	Leaf blister (Taphrina)	
Azaleas, Begonias, Rhododendrons	Leaf blight, leaf scorch, leaf spot	Aphids, mites, rose chafer, Japanese beetle, flea beetle, thrips, leafhopper, red spider, and repels bagworms. <b>Do not use dust on trees. Use only liquid formulations.</b>
Chrysanthemums	Botrytis blight, downy mildew, leaf spots	
Flowering Shrubs, Maples, Hawthorne, Gardenias, Marigolds, Phlox, stocks	Leaf spots	
Roses	Anthracnose, black spot, downy and powdery mildews, leaf spots, stem canker	

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN STRICT ACCORDANCE WITH DIRECTIONS.

Buyers guarantee limited to label claims

Manufactured by:

Bonide Products, Inc.

Oriskany, NY 13424