

B O N I D E
**ROTENONE-COPPER DUST FOR
POTATO-TOMATO-VEGETABLE
GARDENS**

4-30
JAN 10 1975

UNLESS OTHERWISE SPECIFIED
FUNCTIONAL EQUIVALENTS

ROTENONE-COPPER DUST

To control certain insects and to control or prevent certain blights, leaf spots, mildew, and certain other diseases on truck crops. Application for certain insects should be made as soon as insects appear. Applications against certain diseases should begin well before they are expected to appear.

DIRECTIONS

Use as dust or spray any time when conditions are favorable, applied at 7 to 10 day intervals, as soon as plants are above ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required (i.e., after rainfall, etc.)

AS A DUST: This can is a squeeze type duster. Shake can to fluff powder. Set top opening for desired amount (3 or 4 holes). Press sides of can with fingers, using rapid motion for fog effect. Cover foliage with a thin film of dust, both top and undersides of leaves.

AS A SPRAY: Mix 1 pound to 5 gallons of water, or 1 tablespoonful to 1 quart of water. Add powder to small quantity of water first making a paste, then add desired amount of water. Spray to cover all parts of exposed plant both top and undersides of leaves.

ON CUCURBITS: (Cucumber, muskmelon, pumpkin, squash, watermelon) To control striped cucumber beetle, spotted cucumber beetle, two-spotted mite, potato flea beetle, aphids, and to prevent or control scab, anthracnose, bacterial wilt, downy mildew. Apply when insects appear and continue application weekly. Applications against diseases should begin well before they are expected to appear.

ON BEANS: To control bean beetle, leafhopper, tarnished plant bug, to repel mites and aphids, and to control powdery and downy mildew, anthracnose and bacterial blight.

ON CELERY: To control swallowtail butterfly larvae and to prevent or control early, late and bacterial blights.

ON TOMATOES: To control potato flea beetle, Colorado potato beetle larvae, and russet mite, and to prevent or control early and late blight, fruit spot and anthracnose. To control horn worms, apply when worms are still small.

ON EGGPLANT AND POTATOES: To control Colorado potato beetle larvae, leafhoppers, and to repel potato flea beetles. To control phomopsis, alternaria and anthracnose on egg plant and early and late blight on potatoes.

ON ROSES: To control Japanese beetles, aphid, rose chafer and to prevent or control black spot.

CAUTION: Do not apply to food crops within 1 day of harvest. Poison if taken internally.

Do not reuse container. Destroy when empty.

Buyer's Guarantee Limited to Label Claims

© 1973 & Mfg. by BONIDE CHEMICAL CO., INC., Yorkville, N.Y. 13495

4-30

1-10-1975

1/1