

PM 18 4-30

PS 143

5399621

APR 14 1989

Mr. James J. Wurz  
Bonide Chemical Company, Inc.  
2 Wurz Avenue  
Yorkville, NY 13495

Dear Mr. Wurz:

Subject: Bonide Rotenone - Copper Dust for Potato-Tomato-Vegetable  
Gardens  
EPA Registration No. 4-30  
Confidential Statement of Formula and Label Dated April 14,  
1989 Submitted in Response to Registration Standard  
Requirements

The label is acceptable provided that "organic" is removed. It is an implied safety claim. This change must be made at the next printing of labels no longer than six (6) months after the date stamped on this letter. Submit five (5) copies of your final printed labeling before you release the product for shipment using this label.

A stamped copy of the label is enclosed for your records.

The Confidential Statement of Formula enclosed with the "Generic Data Exemption Statement" dated April 14, 1989, will be kept in the file for this product. Further, revisions will be required if changes are made in labels for Manufacturing Use Concentrates supplying the active ingredients for this product.

Sincerely yours,



Phil Hutton  
Product Manager (18)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

52342;I;Schroeder;C;Disk;KEVRIC;06/11/91;CL;NO;CL;Wotek

CONCURRENCES

SYMBOL								
SURNAME								
DATE								


# BONIDE®

**ROTENONE-COPPER DUST FOR  
POTATO-TOMATO-VEGETABLE  
GARDENS**

**INSECTICIDE-FUNGICIDE DUST**

EPA Est. 4-NY-1 EPA Reg. No. 4-30

<b>ACTIVE INGREDIENTS:</b>	
Copper (in basic Copper Sulphate) expressed as metallic	7.00%
Rotenone	75%
Other Cube Resins	1.50%
<b>INERT INGREDIENTS</b>	<b>90.75%</b>



ACCEPTED  
with COMMENTS  
by EPA on 4/30/68

Use only for control of insects,  
Fungicide, and Rotenone. Act as  
amended, for the pesticide  
registered under EPA Reg. No.  
4-30

**BEST AVAILABLE COPY**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS — CAUTION.**  
 May be harmful if swallowed. Avoid breathing dust or spray. Avoid contact with eyes, skin, clothing. Wash thoroughly after use. In case of contact with plenty of water. May cause skin sensitization in certain individuals.

**ENVIRONMENTAL HAZARDS**  
 This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Storage:** Store in original container in a cool dry place which is inaccessible to children and pets, preferably in a locked area.  
**Container Disposal:** Do not reuse empty container. Wrap container and put in trash collection.

## BEST AVAILABLE COPY

**BONIDE ROTENONE-COPPER DUST — To control certain insects and to control or prevent certain blights, leaf spots, mildew, and certain other diseases on truck crops.** Application for certain insects should be made as soon as insects appear. Applications against certain diseases should begin well before they are expected to appear.

### DIRECTIONS

Use as dust or spray any time when conditions are favorable, applied at 7 to 10 day intervals (as soon as plants are in the ground, or as soon as insects appear. Keep foliage covered, especially new growth. In all cases repeat treatment as required (i.e., after rainfall, etc.) Do not apply to food crops within one day of harvest.

**AS FOLIAGE:** To control aphids, leafhoppers, and other sucking insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.

**AS FOLIAGE:** To control leaf beetles, leafhoppers, and other chewing insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.

**AS FOLIAGE:** To control Colorado potato beetle larvae, leafhoppers, and other chewing insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.

**AS FOLIAGE:** To control Colorado potato beetle larvae, leafhoppers, and other chewing insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.

**ON CELERY:** To control swallowtail butterfly larvae and to prevent or control leaf spots and blights.

**ON CUCUMBERS:** To control cucumber beetles, leafhoppers, and other chewing insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.

**ON EGGPLANT AND POTATOES:** To control Colorado potato beetle larvae, leafhoppers, potato flea beetles. To control powdery mildew, anthracnose and bacterial blight on potatoes.

**ON PEPPERS:** To control Colorado potato beetle larvae, leafhoppers, and other chewing insects. Spray thoroughly, especially on the undersides of leaves. Repeat treatments as needed.