

BONIDE

CUKE and MELON DUST

FOR USE ON CUCURBIT PLANTS, such as melons, squash, pumpkins and cucumbers to control flea beetles, striped and twelve spotted cucumber beetles (insects which transmit bacterial wilt) and to prevent or control scab, anthracnose, downy and powdery mildew, early and late blights.

ACTIVE INGREDIENTS:	USDA Reg. No. 453
Rotenone	1.00%
Other Cube Extractives	2.00%
Copper (In basic Copper Sulphate) expressed as metallic	7.00%
INERT INGREDIENTS:	90.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

(Note back panel cautions) NET WEIGHT 1 LB.

CUKE AND MELON DUST

BONIDE CUKE AND MELON DUST is a special formula of micro-pulverized rotenone, neutral copper and wettable diluent. Recommended for use on cucumbers to repel or control potato flea beetles, striped and twelve-spotted cucumber beetles (insects which transmit bacterial wilt). To prevent or control scab, anthracnose and downy and powdery mildew.

BONIDE CUKE AND MELON DUST is suitable for application as a dust or spray. It requires no moisture to be effective. It is therefore possible to apply it to dry or wet foliage at any time, day or night, when dusting or spraying conditions are favorable.

DIRECTIONS

FOR DUSTING: Shake can to fluff powder, then use can as a duster — adjust top openings and squeeze sides with rapid motion for fog effect. Cover all parts of the plant with thin protective film, top and underside of leaves, also, around the vine stem.

FOR SPRAYING: Mix 1 tablespoon per quart water. Add water to powder, strain, add water and spray — agitate sprayer to keep nozzle from plugging.

WHEN TO APPLY: Start applications as soon as plants are above the ground, make repeat applications at 5 to 7 day intervals or as necessary, keep the foliage covered, especially new growth. Make sure to apply thoroughly around the vine stems.

CUKE AND MELON DUST may also be used on muskmelon, pumpkin, squash and watermelon to prevent or control such insects as striped cucumber beetle, spotted cucumber beetle, two-spotted mite, potato flea beetle and non-wooly aphids.

CAUTION: Harmful if swallowed. Avoid inhalation of dust. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not apply to food crops within 1 day of harvest.

Do not reuse container. Destroy when empty.

Buyer's Guarantee Limited to Label Claims

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

4-53

8/16/71

P182

RECEIVED
AUG 16 1971

ACCEPTED
AUG 16 1971
UNDER THE FEDERAL INSECTICIDE
FUNCTION AND FERTILIZER ACT
FOR ECONOMIC INJURY PREVENTION
ED UNDER NO. 7-53 SUBJECT
TO ATTACHED AGREEMENTS