

4-53

11/8/2001

1/4

Bonide Cuke and Melon Dust

Page 1 of 4

# BONIDE


## CUKE and MELON DUST

### FOR USE ON HOME GARDENS ONLY

FOR USE ON CUCURBIT VEGETABLES, Such as melons, muskmelon, watermelon, squash, pumpkins and cucumbers to control flea beetles, cucumber beetles (insects which transmit bacterial wilt) and non-wooly aphids and to prevent or control scab, anthracnose, downy and powdery mildew.

EPA Reg. No. 4-53

EPA Est. No. 4-NY-1



**ACTIVE INGREDIENTS:**

Rotenone .....	1.5%
Other Associated Resins .....	2.2%
Copper <sup>1</sup> .....	7.0%
<b>OTHER INGREDIENTS</b> .....	<b>89.3%</b>
<b>TOTAL</b>	<b>100.00%</b>

<sup>1</sup> In basic Copper Sulphate, expressed as metallic

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side/back panel for additional precautionary statements.

Net Weight 1lb., 4 lb.

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

May be harmful if swallowed, absorbed through the skin, or inhaled. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing. Avoid inhalation of dust or spray. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash thoroughly before reuse.

<b>FIRST AID</b>	
Have label with you when obtaining treatment advice.	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center, doctor, or 1-800-858-7378 immediately.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li></ul>
<b>In on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.</li></ul>

## ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Bonide Cuke and Melon Dust is a special formula of micro-pulverized rotenone, neutral copper and wettable diluent. Bonide Cuke and Melon Dust is suitable for application as a dust or spray. It requires no moisture to be effective. It is therefore possible to apply it to dry or wet foliage at any time, day or night, when dusting or spray conditions are favorable.

**USE RESTRICTIONS: FOR USE ON HOME GARDENS ONLY. FOR USE ON CUCURBIT BEGETABLES, such as melons, muskmelon, watermelon, squash, pumpkins and cucumbers to control flea beetles, striped and twelve-spotted cucumber beetles (insects which transmit bacterial wilt), and non-wooly aphids and to prevent or control scab, anthracnose, downy and powdery mildew.**

Do not apply to food crops within 1 day of harvest.

**WHEN TO APPLY:** At first appearance of pest or its damage after plants are above the ground.

## APPLICATION DIRECTIONS:

### AS A DUST:

#### Mixing Directions:

Use as is. No mixing required.

#### Application Directions:

At first appearance of pest or its damage, shake can to fluff powder, then use can as a duster-adjust top openings, and squeeze sides with rapid motion for fog effect. Cover all parts of the plant with thin protective film, top and underside of leaves, also, around the vine stem. Repeat at 5 to 7-day intervals or as needed, (especially new growth) until pests are under control.

4/4

**Mixing Directions:**

Mix 1 tablespoon per of quart of water. Add water to powder, strain, and add remainder of water and spray-

**Application Directions:**

At first appearance of pest or its damage, spray thoroughly, to point of runoff. Apply to upper and lower leaf surfaces and especially around the vine stem. Agitate sprayer to keep nozzle from plugging. Repeat at 5 to 7 day intervals or as needed, (especially new growth) until pests are under control.

**STORAGE AND DISPOSAL**

**Storage:** Store only in original container, in a dry place inaccessible to children and pets. Store and transport in an upright position.

**Disposal:** **If empty:** Do not reuse empty container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.  
Never place unused product down any indoor or outdoor drain.

For information on pesticide products (including health concerns, Medical emergencies, or pesticide incidents), call National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by:  
Bonide Products, Inc.  
6301 Sutliff Rd.  
Oriskany, NY 13424