

PM 23 4-59 P9175

MAR 31 1997

Ms. Anne Montalone
Bonide Products, Inc.
2 Wurz Ave.
Yorkville, NY 13495-1188

Dear Ms. Montalone:

Subject: Fruit Tree Spray
EPA Registration No. 4-59
Application To Amend Registration To Reinstate
Dust Use of Product for Roses, Evergreens,
Ornamentals and Flowers

The proposed label amendments have been reviewed and found acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended, provided that you:

Submit one (1) copy of the final printed label before releasing the product for shipment under the subject labeling.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL							
SURNAME	EWilson:Diskette:	Captan:	03-31-97				
DATE							

BONIDE

FRUIT TREE SPRAY

CONTROLS INSECTS AND DISEASES ON APPLES,
ROSES, EVERGREENS, ORNAMENTALS AND FLOWERS.

EPA Est. 4-NY-1

EPA Reg. No. 4-59

ACTIVE INGREDIENTS:

*Methoxychlor: 1,1,1-trichloro-2, 2-bis (p-methoxyphenyl) ethane.....	6.00%
**Captan.....	5.87%
Related derivatives.....	0.13%
Malathion: 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate.....	3.00%
Carbaryl (1-naphthyl N-methylcarbamate).....	0.50%
INERT INGREDIENTS.....	84.50%
TOTAL	100.00%

*Equivalent to 5.23% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.72% other isomers and related compounds.

**N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN
DANGER

See back panel for additional precautionary statements.

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 31 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

4-59

NET WEIGHT: _____

BEST AVAILABLE COPY

Buyer's Guarantee Limited to Label Claims.

Manufactured by:
BONIDE PRODUCTS, INC.
Yorkville, NY 13495

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing.

USER SAFETY REQUIREMENTS	
•	Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
•	Wear a hat and eye protection when making overhead applications.
•	Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
•	Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
•	Do not allow children or pets to contact treated area until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT

- If in Eyes:** Flush with plenty of water. Call a physician.
- If on Skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- If Swallowed:** Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.
- If Inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the means high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE: Store in a cool, dry area inaccessible to children or pets.
DISPOSAL: Do not reuse empty container. Wrap container and put in trash Collection.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Apply when air is calm to avoid drift and contact with eyes and skin.
- Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- For trees taller than 10 feet, consider hiring a licensed professional.
- Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- Do not repeat applications more than once a week.

BONIDE FRUIT TREE SPRAY is a combination of ingredients useful in controlling certain insect pests and plant diseases common to apples, roses, evergreens, ornamentals and flowers listed below. Spray is easily applied by tank, trombone, hand or hose end sprayers. Where nozzle plugging is a problem, use "Bonide Liquid Fruit Tree Spray".

DIRECTIONS

Select still periods for application (early morning or evening) to reduce waste by blow away and blow back on applicator. Make thin paste first with powder then dilute with balance of water to make quantity desired. During application, agitate sprayer to keep powder in suspension. Spray trees or plants lightly but thoroughly cover all parts of plant or tree, particularly undersides of leaves. Do not apply if rain is expected shortly after application. It is easier to prevent disease and scale damage than to cure same. Therefore, a preventive spray schedule is recommended.

BEST AVAILABLE COPY

SITE	PESTS	DISEASES	WHEN TO APPLY	APPLIC. CON. RATE	PRECAUTIONS AND RESTRICTIONS
Apples	Aphids, Apple maggot, Bagworm, Bud moth, Fruit fly & worm, Codling moth, Plum curculio, Flea beetles, Fruit tree leaf roller, Gypsy moth, Japanese beetle, Leaf hoppers, Mealy bugs, Mites (Clover, European-red, two-spotted and Willi-mette), Oriental fruit moth, Pear slugs, Psylla, Red banded leaf roller, Scale (Forbes, Putman, San Jose), Spittle bug, Tent caterpillars, Unspotted tentiform leaf miner, Yellow neck caterpillar	Bitter rot, Black spot, Black rot, Blossom blight, Botryosphaeria rot, Botrytis blossom end rot, Brooks fruit spot, Brown rot, Coryneum blight, Downy mildew, Fly speck, Frog eye, Fruit rot, Leaf spot, Sooty blotch	Start at greentip and repeat at seven (7) day intervals until bloom. At petal fall, repeat at ten (10) day intervals up to seven (7) days before harvest.	Use ten (10) level tbs. per gallon of water. Use 1 to 2 pints spray suspension on 5 to 8 ft. trees. Use 1 to 2 qts. spray suspension on 8 to 14 ft. trees. On large trees, use 1 to 3 gals. spray suspension.	To protect pollinating insects, do not spray during blossoming period. Use Bonide Captan, that contains only a fungicide, during blossoming period, if disease control is needed. Apply up to seven (7) days before harvest. Remove residue before eating or canning by washing or wiping.
Roses, Evergreens, Ornamentals, Flowers	Aphids, Bagworm, Bud moth, Flea beetles, Gypsy moth, Japanese beetle, Leaf hoppers, Mealy bugs, Mites (Clover, European-red, two-spotted and Willimette), Psylla, Red banded leaf roller, Scale (Forbes, Putman, San Jose), Spittle bug, Tent caterpillars, Unspotted tentiform leaf miner, Yellow neck caterpillar	Black spot, Blossom blight, Downy mildew, Leaf spot	Begin applications when pests or disease symptoms first appear or conditions favor their development and repeat at weekly intervals or as necessary to maintain control.	Use five (5) level tbs. per gallon of water. Use 1 to 2 pints spray suspension on 5 to 8 ft. trees. Use 1 to 2 qts. spray suspension on 8 to 14 ft. trees. On large trees, use 1 to 3 gals. spray suspension.	May cause damage to crassula or ferns.

***As a Dust for Roses, Evergreens, Ornamentals and Flowers:**

For control of insects such as aphids, bagworms, Japanese Beetles, mites, and other insects as listed above for Apples, which may attack roses, evergreens, ornamentals and flowers, apply this product on a calm day with a suitable dust applicator to cover all portions of the plant with a fine dust film on stems, leaves and buds. To protect bees, do not apply to blooms. May cause damage to Crassula or ferns. Do not apply to Cannariff Junipers. Do not apply as a dust to any food crops. Do not breath dust!

BEST AVAILABLE COPY