

4-59

07/08/2002

1/5

MASTER LABEL



FRUIT TREE SPRAY

[Alternate Brand Name(s): BONIDE® Rose and Flower Spray or Dust; FRANK'S Rose & Flower Dust or Spray]

Optional Marketing Statements

- Controls insects and disease on apples, evergreens, ornamentals and flowers.
- Wettable Powder
- Controls insects and diseases on apples, roses, evergreens and flowers.
- May be used as a spray or dust.
- Ready-to-Use (when marketed for application as a dust)

ACTIVE INGREDIENTS:


Captan (CAS No. 133-06-2)	5.87%
Related Derivatives	0.13%
Malathion (CAS No. 121-75-5)	3.00%
Carbaryl (CAS No. 63-25-2)	0.50%
OTHER INGREDIENTS:	90.50%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/other panel(s) for first aid and additional precautionary statements.

NET WEIGHT _____ LB. (____ g)



FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: Contains carbaryl, a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.	

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, socks and shoes.

USER SAFETY RECOMMENDATIONS
<ul style="list-style-type: none"> • Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

General Precautions and Restrictions

- **ENTRY RESTRICTIONS:** Do not allow people or pets to enter the treated area until sprays have dried or dust has settled.
- Apply when air is calm to avoid drift and contact with eyes and skin. Do not allow this product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not apply to lawns.
- Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- For trees taller than 10 feet, consider hiring a licensed professional.
- Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- Do not repeat applications more than once a week.

BONIDE FRUIT TREE SPRAY is a combination of ingredients useful in controlling certain insect pests and plant diseases common to apples, roses, evergreens, ornamentals and flowers listed below. Spray is easily applied by tank, trombone, hand or hose end sprayers. Where nozzle plugging is a problem, use Bonide Complete Fruit Tree Spray liquid.

DIRECTIONS

Select still periods for application (early morning or evening) to reduce waste by blow away and blow back on applicator. Make thin paste first with powder then dilute with balance of water to make quantity desired. During application, agitate sprayer to keep powder in suspension. Spray trees or plants lightly but thoroughly cover all parts of plant or tree, particularly underside of leaves. Do not apply if rain is expected shortly after application. It is easier to prevent disease and scale damage than to cure same. Therefore, a preventive spray schedule is recommended.

SITE	PESTS	DISEASES	WHEN TO APPLY	APPLICATION RATE	PRECAUTIONS & RESTRICTIONS
Apples	Aphid, Apple maggot, Bagworm, Bud moth, Fruit fly & worm, Codling moth, Plum curculio, Flea beetle, Fruit tree leaf roller, Gypsy moth, Japanese beetle, Leaf hopper, Mealybug, Mites (Clover, European red, two-spotted and Wilmette), Oriental fruit moth, Pear slug, Psylla, Red banded leaf roller, Scale (Forbes, Putman, San Jose), Spittle bug, Tent caterpillar, Unspotted tentiform leaf miner, Yellow neck caterpillar.	Alternaria leaf spot, Apple fruit rot, Bitter rot, Black rot, Blossom blight, Botryosphaeria rot, Botrytis blossom end rot, Brooks fruit spot, Brown rot, Fly speck, Frogeye, Sooty blotch.	Start at greentip and repeat at seven (7) day intervals until bloom. At petal fall, repeat at ten (10) day intervals up to seven (7) days before harvest.	Use ten (10) level Tbs. per gallon of water. Use 1 to 2 pints spray suspension on 5 to 8 ft. trees. Use 1 to 2 qts. spray suspension on 8 to 14 ft. trees. On large trees, use 1 to 3 gals. spray suspension.	To protect pollinating insects, do not spray during blossoming period. Use Bonide Captan, that contains only a fungicide, during blossoming period, if disease control is needed. Apply up to seven (7) days before harvest. Remove residue before eating or canning by washing or wiping.
Roses, Evergreens, Ornamentals, Flowers	Aphid, Bagworm, Bud moth, Flea beetle, Gypsy moth, Japanese beetle, Leaf hopper, Mealybug, Mites (Clover, European red, two-spotted and Wilmette), Psylla, Red banded leaf roller, Scale (Forbes, Putman, San Jose), Spittle bug, Tent caterpillar, Unspotted tentiform leaf miner, Yellow neck caterpillar.	Alternaria leaf spot, Black spot, Botrytis blossom blight.	Begin applications when pests or disease symptoms first appear or conditions favor their development and repeat at weekly intervals, or as necessary, to maintain control.	Use five (5) level Tbs. per gallon of water. Use 1 to 2 pints spray suspension on 5 to 8 ft. trees. Use 1 to 2 qts. spray suspension on 8 to 14 ft. trees. On large trees, use 1 to 3 gals. spray suspension.	May cause damage to Crassula or ferns.

To apply as a Dust for Roses, Evergreens, Ornamentals and Flowers:

For control of insects such as aphids, bagworms, Japanese beetles, Mites, and other insects listed above which may attack roses, evergreens, ornamentals and flowers, apply this product on a calm day with a suitable dust applicator to cover all portions of the plant with a fine dust film on stems, leaves and buds. To protect bees, do not apply to blooms. May cause damage to Crassula or ferns. Do not apply to Cannartii Junipers. Do not apply as a dust to evergreen or ornamental trees. Do not apply as a dust to any food crops. Do not breathe dust!

STORAGE: Store and transport in an upright position. Store in a cool, dry area inaccessible to children or pets.
DISPOSAL: *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partially filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused products down any indoor or outdoor drain.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents, call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by
Bonide Products, Inc.
Oriskany, NY 13424

