

5/19/72

LIVESTOCK BARN: As a residual spray for the control of flies, mosquitoes, spray at the rate of 4 ozs. in 2½ gals. of water. Apply to walls, ceilings, beams, window sills, and other surfaces where flies come to rest using approximately 2 gals. per 1000 sq. ft. of surface.

GLADIOLUS, DAFFODIL BULBS: For wireworm control — Dust dry bulbs at time of planting. Use 3 ozs. to each 100 pounds of bulbs.

WARNING: May be fatal if swallowed. Avoid contact with skin and eyes. Avoid breathing of vapor, dust or spray mist. May be absorbed through the skin. **Wash after handling.**

Avoid contamination of feed and foodstuffs. Avoid contamination of drinking cups and other utensils. Wash off if they are contaminated.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from area treated. Do not reuse container. Destroy when empty.

Do not use in poultry houses or in dairy barns.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer's guarantee limited to label claims.

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

4-74
BONIDE

LINDANE
INSECTICIDE SPRAY

ACTIVE INGREDIENT:
Gamma Isomer of Benzene Hexachloro-
(from Lindane)
INERT INGREDIENTS


WARNING: KEEP OUT OF
REACH OF CHILDREN
8 OUNCES

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer's guarantee limited to label claims

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502



LINDANE 25% W
INSECTICIDE SPRAY POWDER

ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride
(from Lindane)

INERT INGREDIENTS

EPA Reg. No. 176

25.0%
75.0%

WARNING: KEEP OUT OF REACH OF CHILDREN

8 OUNCES NET

Bonide LINDANE 25% W is a special micronized wettable powder containing 25% Lindane, recommended for certain insects on vegetable crops, shade trees, nursery and ornamental plants.

DIRECTIONS

CABBAGE, CAULIFLOWER, LETTUCE: For aphids, thrips — Spray at the rate of 1 tablespoonful in 1 gal. of water. Make weekly applications as needed. Do not apply to cabbage and cauliflower after edible parts start to form. Do not apply to leaf lettuce after heading stage. Do not apply to head lettuce after heads start to form.

CUCUMBER, PUMPKIN, SQUASH: For cucumber beetles, thrips, squash borer, aphids — Spray at the rate of 1 tablespoon in 1 gal. of water. Make weekly application. Begin when plants are about 1 inch high. On cucumbers and squash, do not apply after blossoms open. Pumpkins may be sprayed up to day of harvest.

SHADE TREES, NURSERY AND ORNAMENTAL PLANTS: Birch leaf miner, aphids, thrips, leaf hopper, weevil and flea beetles — Spray at the rate of 1 tablespoonful in 1 gal. of water. Apply as soon as insects appear, repeat as necessary. taxus