

LAWN, TURF & GARDEN INSECTICIDE is a finely formulated, atomized insecticide for use on lawn and turf, shade trees, ornamentals, flowers and vegetables against insects that attack them.

**AREA TREATMENT:** Use 2 pounds for each 1,000 square feet of area around the buildings infested with crickets, millipedes, ~~scorpions~~, scorpions, and wasps. Apply a 15 foot band around structure and extending to the foundation to prevent them from entering. Work into soil with coarse spray after treatment. Repeat two to three weeks later if necessary.

**TO CONTROL ANTS:** Spot apply liberally to ant hills and wash insecticide into soil.

**CAUTION:** Keep out of reach of children and animals. Harmful if inhaled or swallowed. Avoid breathing dust. Do not take internally. Avoid contact with skin. Wear regular long-sleeved clothing during application. If making daily commercial applications, change into protective clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after use.

**Note for physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, streams and ponds. Keep out of water areas until watered in by rainfall or hosing.

To protect honey bees, avoid use, if possible, during periods when bees are visiting flowers. If necessary to use during such periods, warn beekeepers well in advance to leave hives at a distance until 1 week after application.

Buyer's guarantee limited to label claims.

MANUFACTURED BY

**BONIDE CHEMICAL CO., INC., Utica, New York 13502**



### DIRECTIONS

**TO SPRAY:** Mix LAWN, TURF & GARDEN with water at the rate of 10 tablespoons per gallon of water (5 pounds to 50 gallons of water). Use a good pressure sprayer and apply thoroughly to all parts of the plants as soon as insects appear, repeat at 7 to 10 day intervals or as often as needed.

**SHADE TREES, ORNAMENTALS:** Japanese beetles, rose chafers, elm leaf beetles, leaf beetles, green striped maple worm, bud worms, leaf hoppers, tent caterpillars, ~~scorpions~~.

**ROSES, ANNUAL AND PERENNIAL FLOWERING PLANTS:** Japanese beetles, rose chafers, leaf beetles, leaf hoppers, flea beetles, tarnished plant bugs, leaf rollers, European corn borer, diamond borer, worms, exposed thrips, ~~scorpions~~.

**GARDEN VEGETABLES:** Cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, dandelion, bark moth, imported cabbage worm (allow 3 days between last application and harvest) of trees, Colorado potato beetles, flea beetles, leaf hoppers. Beans, Mexican bean beetle, flea beetle. Consult your experiment station for best time to apply in your locality.

**LAWN & TURF INSECTS:** Ants, bluegrass billbug, chinch bugs, cutworms, earwigs, fire ant, army worm, fleas, ~~scorpions~~, leaf hoppers, millipedes, mosquitoes, and sod webworms (lawn moths). Cut lawn close, remove clippings and water thoroughly before treatment. Apply uniformly and back up with water spray after treatment. Repeat two to three weeks later if necessary. Apply to surface at rate of 88 lbs. per acre or two pounds per 1,000 sq. ft. (a 50-foot by 70-foot area).

LAWN, TURF & GARDEN INSECT KILLER is a blended attritionized insecticide recommended for use on lawn and turf, shade trees, ornamentals, flowers and vegetables for the control of insects that attack them.

AREA TREATMENT: Use 2 pounds for each 1,000 square feet of area around homes and other buildings infested with crickets, millipedes, ~~termites~~, scorpions, and wasps. Treat at least a 15 foot band around structure and extending to the foundation to prevent entry into buildings. Work into soil with coarse spray after treatment. Repeat two to three weeks later if necessary.

TO CONTROL ANTS: Spot apply liberally to ant hills and wash insecticide into soil.

**CAUTION:** Keep out of reach of children and animals. Harmful if inhaled or swallowed. Avoid breathing dust. Do not take internally. Avoid contact with skin. Wear regular long sleeved work clothing during application. If making daily commercial applications, change to clean clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after handling.

**Note for physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, streams and ponds. Keep pets off treated areas until watered in by rainfall or hosing.

To protect honey bees, avoid use, if possible, during periods when bees are visiting the crop. If necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Buyer's guarantee limited to label claims.

MANUFACTURED BY

**BONIDE CHEMICAL CO., INC., Utica, New York 13502**

47  
ACCEPTED  
Nov 9, 1961

**ROSES, ANNUAL AND PERENNIAL FLOWERING PLANTS:** Japanese beetles, rose chafers, ants, leaf hoppers, flea beetles, tarnished plant bugs, leaf rollers, European corn borer, diamond back worms, exposed thrips, a [REDACTED]

**GARDEN VEGETABLES:** Cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, diamond back moth, imported cabbage worm (allow 3 days between last application and harvest). Potatoes, Colorado potato beetles, flea beetles, leaf hoppers. Beans, Mexican bean beetle, flea beetle. Consult your experiment station for best time to apply in your locality.

**LAWN & TURF INSECTS:** Ants, bluegrass billbug, chinch bugs, cutworms, earwigs, fall armyworm, fleas, [REDACTED], leaf hoppers, millipedes, mosquitoes, and sod webworms (lawn moths). Cut lawn close, remove clippings and water thoroughly before treatment. Apply uniformly and soak in with water spray after treatment. Repeat two to three weeks later if necessary. Apply to soil surface at rate of 88 lbs. per acre or two pounds per 1,000 sq. ft. (a 50-foot by 20-foot area).

Do not reuse container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.