

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (& DOMESTIC ANIMALS) CAUTION

Keep children and pets off treated areas until dry. Avoid contact with treated areas. Wash hands thoroughly after handling. Avoid breathing dust or mist. Avoid contact with eyes, nose, mouth, or skin. Wash clothing if contaminated. Do not eat, drink, or smoke while using this product. Do not use in enclosed areas. Do not use in areas where food is stored or prepared. Do not use in areas where food is eaten. Do not use in areas where food is stored or prepared. Do not use in areas where food is eaten.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink plenty of water. Do not induce vomiting. If Inhaled: Remove to fresh air. If on Skin: Wash with soap and water. If on Eyes: Flush with water for 15 minutes.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic life. Do not apply near water. Do not apply to lawns or gardens near water. Do not apply to lawns or gardens near water.

DIRECTIONS FOR USE

DISPOSAL

LAWN, TURF & GARDEN INSECT KILLER is a blended attractant insecticide recommended for use on lawn and turf, shade trees, ornamentals, flowers and vegetables for the control of insects that attack them and for control of fleas and ticks on cats and dogs.

TO SPRAY: Mix LAWN, TURF & GARDEN with water at the rate of 10 tablespoons to 1 gallon of water (5 pounds to 50 gallons of water). Use a good pressure sprayer and apply thoroughly to all parts of the plants as soon as insects appear. Repeat at 7 to 10 day intervals or as often as needed.

SHADE TREES, ORNAMENTALS: Japanese beetles, rose chafers, elm leaf beetles, leaf miners, green striped maple worm, bud worms, leaf hoppers, tent caterpillars.

ROSES, ANNUAL AND PERENNIAL FLOWERING PLANTS: Japanese beetles, rose chafers, ants, leaf hoppers, flea beetles, tarnished plant bugs, leaf rollers, European corn borer, diamond back worms, exposed thrips.

GARDEN VEGETABLES: Cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, diamond back moth, imported cabbage worm (allow 3 days between last application and harvest). Potatoes, Colorado potato beetles, flea beetles, leaf hoppers. Beans, Mexican bean beetle, flea beetle. Consult your experiment station for best time to apply in your locality.

LAWN & TURF INSECTS: Ants, bluegrass billbug, chinch bugs, cutworms, earwigs, fall armyworm, fleas, leaf hoppers, millipedes, mosquitoes, and sod webworms (lawn moths). Cut lawn close, remove clippings and water thoroughly before treatment. Apply uniformly and soak in with water spray after treatment. Repeat two to three weeks later if necessary. Apply to soil surface at rate of 88 lbs. per acre or two pounds per 1,000 sq. ft. (a 50-foot by 20-foot area).

TO CONTROL ANTS: Spot apply liberally to ant hills and wash insecticide into soil.

AREA TREATMENT: Use 2 pounds for each 1,000 square feet of area around homes and other buildings infested with crickets, millipedes, scorpions, and wasps. Treat at least a 15 foot band around structure and extending to the foundation to prevent entry into buildings. Work into soil with coarse spray after treatment. Repeat two to three weeks later if necessary.

CATS AND DOGS: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens or puppies under 4 weeks of age. Do not repeat application within 7 days.

Buyer's guarantee limited to label claims. Mfg. by Bonide Chemical Co., Inc., Yorkville, NY 14395

BONIDE

SEVIN® 10%
LAWN, TURF & GARDEN INSECT KILLER

CONTROLS bluegrass billbugs, chinch bugs, cutworms, earwigs, fleas, leafhoppers, fall armyworm, millipedes, mosquitoes and sod webworms (lawn moths), Japanese beetles, rose chafers, leaf miners, elm leaf beetles, exposed thrips, leafrollers, tent caterpillars and certain other insects as listed. For use on cats and dogs.

ACTIVE INGREDIENTS: Carbaryl, D. Nuprol, N. Methidathion. INERT INGREDIENTS: Sevin is the registered trademark of Union Carbide Corporation for their Carbaryl insecticide.

CAUTION: KEEP OUT OF REACH OF CHILDREN

1 LB. NET WEIGHT

of fleas and ticks on cats and dogs

TO SPRAY: Mix **LAWN, TURF & GARDEN** with water at the rate of 10 tablespoons to 1 gallon of water (5 pounds to 50 gallons of water). Use a good pressure sprayer and apply thoroughly to all parts of the plants as soon as insects appear, repeat at 7 to 10 day intervals or as often as needed.

SHADE TREES, ORNAMENTALS: Japanese beetles, rose chafers, elm leaf beetles, leaf miners, green striped maple worm, bud worms, leaf hoppers, tent caterpillars.

ROSES, ANNUAL AND PERENNIAL FLOWERING PLANTS: Japanese beetles, rose chafers, elm leaf hoppers, flea beetles, tarnished plant bugs, leaf rollers, European corn borer, diamond back worms, exposed to soil.

GARDEN VEGETABLES: Cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, diamond back moth, imported cabbage worm (allow 3 days between last application and harvest). Potatoes, Colorado potato beetle, flea beetles, leaf hoppers. Beans, Mexican bean beetle, flea beetle. Consult your experiment station for best time to apply in your locality.

LAWN & TURF INSECTS: Ants, bluegrass billbug, chinch bugs, cutworms, earwigs, fall armyworm, grasshoppers, millipedes, mosquitoes, and sod webworms (lawn moths). Cut lawn close, remove clippings and water thoroughly before treatment. Apply uniformly and soak in with water spray after treatment. Repeat two to three weeks later if necessary. Apply to soil surface at rate of 88 lbs. per acre or two pounds per 1,000 sq. ft. (or 1 lb. by 20-foot area).

TO CONTROL ANTS: Spot apply liberally to ant hills and wash insecticide into soil.

AREA TREATMENT: Use 2 pounds for each 1,000 square feet of area around homes and other buildings infested with crickets, millipedes, scorpions, and wasps. Treat at least a 15-foot band around structure and extending to the foundation to prevent entry into buildings. Work into soil with coarse spray after treatment. Repeat two to three weeks later if necessary.

CATS AND DOGS: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens or puppies under 4 weeks of age. Do not repeat application within 7 days.

Mfg. by Bonide Chemical Co., Inc., Yorkville, NY 14395

BONIDE

SEVIN® 10%

LAWN, TURF & GARDEN INSECT KILLER

CONTROLS bluegrass billbugs, chinch bugs, cutworms, earwigs, fleas, leafhoppers, fall armyworm, millipedes, mosquitoes and sod webworms (lawn moths), Japanese beetles, rose chafers, leaf miners, elm leaf beetles, exposed thrips, leafrollers, tent caterpillars and certain other insects as listed. For use on cats and dogs.

ACTIVE INGREDIENTS

Chlorpyrifos 10.00%

INERT INGREDIENTS

90.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

1 LB. NET WEIGHT

EPA Reg. No. 478

EPA Reg. No. 478