

Read entire label. Use strictly in accordance with label cautions, warnings and directions. Use a *Paint brush, roller, pressure oil can, Apply to areas to be treated*; making repeat treatment as necessary.

CAUTION: May be absorbed through skin. Harmful if swallowed. Avoid inhalation of spray mist. Avoid skin contact. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets and humans. Store out of reach of children. (This product is toxic to fish and wildlife. Birds feeding on treated area may be harmed.)

Made by
BONIDE CHEMICAL CO., INC., UTICA, N. Y.

Buyer's Guarantee Limited to Label Terms

makes up to 96 gallons of finished spray.



BORER-KILLER is a liquid formulation containing 20% Lindane, with spreader and emulsifier. It dissolves instantly. Put required amount in *PAIL* and add water (self-mixing). Will not plug nozzle.

DIRECTIONS FOR USE

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSAGE	TIME TO APPLY
Elm (English and Scotch) Birch Locust	Leaf Miners	1 teaspoonful/ 1 gal. water	Apply when small miners appear in early and mid-May. Apply when small miners appear in early July.
Pines	Pine Bark Beetles	1 pint/5 gals. water	Spray trunks thoroughly — wet trunks to run-off.
Pine, Spruce	White Pine Weevil	1 Tablespoonful/ gal. water	Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus, Hemlock	Strawberry Root Weevil Taxus Weevil	1 Tablespoonful/ 1 gal. water	Apply in May or early June to the trunks and around base of trees, soaking the bark and wetting the soil surface.
Apples, Cherries, Peaches	Fruit Tree Bark Beetles Flat-Headed Borers Round-Headed Borers	1 Tablespoonful/ 1 gal. water 3 Pints/100 gals. water	Apply to bark of tree trunk and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.
Maple, Aspen, Poplar, Ash, Willows, Elm, Locust	Aphids, Bark Beetles, Wood Boring Beetles	3 pints/50 gals. water 2 Tablespoonfuls/ 1 gal. water	Apply in spring when buds begin to break. Repeat in 10-14 days. Elms — Apply to foliage when Aphids are present during summer, repeat in fall — soak trunks to run-off.
Peaches	Peach Tree Borer	3 pints/100 gals. water 1 Tablespoonful/ gal. water	Spray lower trunk and around base of tree, soaking bark and wetting trunk from crotch to ground. Do not apply to fruit or foliage.
Dogwood, Rhododendron, Lilac	Dogwood Borer, Rhododendron Borer, Lilac Borer	1 Tablespoonful/ 1 gal. water	Make 2 to 3 applications 7 to 10 days apart in May and early June.

292

ACCEPTED
MAR 15 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 127 SUBJECT
TO ATTACHED COMMENTS.

127