

WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

Read entire label. Use strictly in accordance with label cautions, warnings and directions. Use a paint brush, roller, pressure oil can, apply to areas to be treated, making repeat treatment as necessary.

CAUTION: May be absorbed through skin. Harmful if swallowed. Avoid inhalation of spray mist. Avoid skin contact. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets and humans. Store out of reach of children. Food utensils such as teaspoons, tablespoons should not be used for food purposes after use with pesticides.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.


Do not reuse container. Destroy when empty.

Buyer's Guarantee Limited to Label Claims

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N.Y. 13502

makes up to 96 gallons of finished spray.



BONIDE BORER-MINER KILLER

BIRCH LEAF MINER, FLAT AND ROUND HEADED BORERS, PEACH TREE BORERS, PINE BARK BEETLES, STRAWBERRY ROOT WEEVIL, APHIDS AND CERTAIN OTHER BORERS

ACTIVE INGREDIENTS:
Gamma Isomer of Benzene Hexachloride from Lindane 20.00%
INERT INGREDIENTS: 80.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions) 1 PT. NET

BORER KILLER is a liquid formulation containing 20% Lindane, with spreader and emulsifier. It dissolves instantly. Put required amount in pail and add water (self-mixing). Will not plug nozzle.

DIRECTIONS FOR USE

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSAGE	TIME TO APPLY
Elm (English and Scotch) Birch Locust	Leaf Miners	1 teaspoonful/ 1 gal. water	Apply when small miners appear in early and mid-May. Apply when small miners appear in early July.
Pines	Pine Bark Beetles	1 pint/5 gals. water	Spray trunks thoroughly — wet trunks to run-off.
Pine, Spruce	White Pine Weevil	1 Tablespoonful/ gal. water	Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus, Hemlock	Strawberry Root Weevil Taxus Weevil	1 Tablespoonful/ 1 gal. water	Apply in May or early June to the trunks and around base of trees, soaking the bark and wetting the soil surface.
Apples, Cherries, Peaches	Fruit Tree Bark Beetles Flat-Headed Borers Round-Headed Borers	1 Tablespoonful/ 1 gal. water 3 Pints/100 gals. water	Apply to bark of tree trunk and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.
Maple, Aspen, Poplar, Ash, Willows, Elm, Locust	Aphids, Bark Beetles, Wood Boring Beetles	3 pints/50 gals. water 2 Tablespoonfuls/ 1 gal. water	Apply in spring when buds begin to break. Repeat in 10-14 days. Elms — Apply to foliage when Aphids are present during summer, repeat in fall — soak trunks to run-off.
Peaches	Peach Tree Borer	3 pints/100 gals. water 1 Tablespoonful/ gal. water	Spray lower trunk and around base of tree, soaking bark and wetting trunk from crotch to ground. Do not apply to fruit or foliage.
Dogwood Rhododendron, Lilac	Dogwood Borer, Rhododendron Borer, Lilac Borer	1 Tablespoonful/ 1 gal. water	Make 2 to 3 applications 7 to 10 days apart in May and early June.

TESTED

1971

**INSECTICIDE
REGISTERED
EPA**

DE

MINER

**ROUND HEADED
S, PINE BARK
NOT WEEVIL,
Borer Borers**

USDA Reg. No. 4-84
in Lindane 20.00%
. 80.00%

**SAFE FOR CHILDREN
(NS) 1 PT. NET**

BORER KILLER is a liquid formulation containing 20% Lindane, with spreader and emulsifier. It dissolves instantly. Put required amount in pail and add water (self-mixing). Will not plug nozzle.

DIRECTIONS FOR USE

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSAGE	TIME TO APPLY
Elm (English and Scotch) Birch Locust	Leaf Miners	1 teaspoonful/ 1 gal. water	Apply when small miners appear in early and mid-May. Apply when small miners appear in early July.
Pines	Pine Bark Beetles	1 pint/5 gals. water	Spray trunks thoroughly — wet trunks to run-off.
Pine, Spruce	White Pine Weevil	1 Tablespoonful/ gal. water	Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus, Hemlock	Strawberry Root Weevil Taxus Weevil	1 Tablespoonful/ 1 gal. water	Apply in May or early June to the trunks and around base of trees, soaking the bark and wetting the soil surface.
Apples, Cherries, Peaches	Fruit Tree Bark Beetles Flat-Headed Borers Round-Headed Borers	1 Tablespoonful/ 1 gal. water 3 Pints/100 gals. water	Apply to bark of tree trunk and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.
Maple, Aspen, Poplar, Ash, Willows, Elm, Locust	Aphids, Bark Beetles, Wood Boring Beetles	3 pints/50 gals. water 2 Tablespoonfuls/ 1 gal. water	Apply in spring when buds begin to break. Repeat in 10-14 days. Elms — Apply to foliage when Aphids are present during summer, repeat in fall — soak trunks to run-off.
Peaches	Peach Tree Borer	3 pints/100 gals. water 1 Tablespoonful/ gal. water	Spray lower trunk and around base of tree, soaking bark and wetting trunk from crotch to ground. Do not apply to fruit or foliage.
Dogwood Rhododendron, Lilac	Dogwood Borer, Rhododendron Borer, Lilac Borer	1 Tablespoonful/ 1 gal. water	Make 2 to 3 applications 7 to 10 days apart in May and early June.