

Front Panel

S O N I D E

T O M C A T (h)

(Picture of Cat)

Gets Rid of rats and
Mice the easy way

Manufactured under exclusive U.S. Patentrights
No.2,813,058 & A.A.H.F. Pat. No. 2,427,578

VILW
WINDOW

CAUTION: Keep out of reach of children
(See side panel for other cautions)

14 OZ. NET WEIGHT

ACTIVE INGREDIENTS:

3(a-acetylbenzyl)-4-hydroxycoumarin (warfarin).....	0.025%
INERT INGREDIENTS:.....	99.975%
TOTAL	100.000%

U.S.P.A. Reg. No. 4-90

ACCEPTED
 OCT 24, 1967
 UNITED STATES DEPARTMENT OF AGRICULTURE
 FEDERAL BUREAU OF INVESTIGATION
 FOR THE PROTECTION OF THE CONSUMER
 ED WINTERMANN SUBJECT
 TO AFR...

rear Panel #2

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of rat and mouse blood and upon repeated feedings causes fatal hemorrhage.

(Picture of Box)

(Picture of Rat Hole)

(Picture of Wood Pile)

(Picture of Bait Box)

(Picture of
Board Protecting Bait)

Baiting should continue as long as feeding is noticed. Assure an uninterrupted supply of bait for a period of not less than 12-15 days and continue baiting until signs of feeding have stopped. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and refilled as often as necessary.

Note caution on side panel.

MANUFACTURED BY
SONIDE CHEMICAL CO., INC.
UTICA, NEW YORK 13502

Left side panel #3

DIRECTIONS FOR USE

RATS: Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are known to feed. Protect bait from children, pets, poultry, and livestock by placement out of their reach or by use of boards or bait boxes. (see illustration on back of box) Replace stale, damp or mouldy bait with fresh bait and continue baiting until feeding stops.

MICE: Use same general directions for rats except place tablespoon amounts of bait in shallow containers where mice or signs of mice have been seen. Mice being more difficult to control need more bait placements at intervals of 8 to 12 feet and may take a longer period of baiting. Keep baits fresh and continue until feeding stops.

right side panel #4

CAUTION: Protect children, pets, and domestic animals from bait. If swallowed by humans or animals this material may reduce clotting ability of blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions would be indicated as in the case of hemorrhage caused by overdoses of 3-hydroxycoumarin.

Top panel #1

SUMMARIZED TO READ YOUR HOME ON BACK OF RATS AND MICE, IF USED AS DIRECTED, ON YOUR MONEY BACK

Bottom Panel #6

Same as Panel #1