

# TURF KARE GRUB AND INSECT KILLER

... most effective insecticide for use on lawns around prime areas, parks, ... In many instances prevents subsurface insect damage to lawns for up to ... nuisance insects for weeks after application. Can be applied any time spring, ... when soil is not frozen.

## DIRECTIONS

Mow the lawn to one-half inch remove clippings. Measure the area to be treated and apply contents of this bag to area stated on front of bag. Insect areas may be spot treated by sifting lightly direct from bag. Spreader settings for wide openings will work on most spreaders but due to wear and different walking rates, adjustment of spreader is necessary to avoid gross over or under dosage. Some starting point settings follow:

<u>Make &amp; Model</u>	<u>Setting</u>	<u>Make &amp; Model</u>	<u>Setting</u>
Montgomery Ward 89 1031	8 1/2	Jackson 18-26 18"	7
Sears Roebuck 19"	7	Salem Micro-Gut 22"	7
Scott	8	Cyclone B 5' Swath (Still air only)	6

For subsurface insects (ie. grubs) etc. Water in copiously after application to move insecticide into root zone. Avoid wash away.

For surface insects such as sod webworms (lawn moths), mole crickets, etc. Mow lawn thoroughly before application to bring insects to surface.

For other surface insects (ie. chiggers, earwigs): Do not water or plan to apply if rain is expected before or immediately after application.

When applied for surface insects by either method, subsurface insects will be controlled when insecticide is moved to root zone by rain or watering.

With heavy insect infestation or heavy black soils increase spreader openings by 1/3 and apply bag contents to 1/2 the area stated on front of bag. This will apply the equivalent of 10 lbs. Chlor-dane per acre.

## CAUTIONS

Keep out of reach of children. Avoid breathing of dust. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash thoroughly. Avoid contamination of food and foodstuffs, water supplies and feeding dishes of pets.

Do not allow pets or children on treated areas until insecticide has been thoroughly soaked into soil and treated areas have dried completely.

This product is toxic to fish and wildlife. Keep it out of lakes, ponds, streams. Birds feeding on treated areas may be harmed.

WARNING: RESTRICTED TO THE TRADE ONLY

© 1974, 1975

REGISTERED TRADEMARK OF THE CHEMICAL COMPANY