

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS — WARNING

May be fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARD

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty containers. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of smoke.

Buyer's Guarantee Limited to Label Claims and Refund of Purchase Price.

MFG. BY BONIDE CHEMICAL CO., INC., YORKVILLE, N.Y. 13495

BONIDE

CHLORDANE 74%E

FOR TERMITES AND JAPANESE BEETLE GRUB CONTROL

ACTIVE INGREDIENTS

Technical Chlordane
Petroleum distillate

INERT INGREDIENTS

Equivalent to 44.4% Octachloro 4,7-methanotetrahydrodane
and 29.6% inert compounds.
This material contains 8 lbs Technical Chlordane per gallon.

E.P.A. Est. 4-NY-1 E.P.A. Reg. No. 4-96

74%
20%
6%

WARNING: KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Precautionary Statements

In case of spillage on skin wash promptly with soap and water. Internal — Emetic
of 1 tablespoonful mustard in tumbler of water. Call a physician.

1 QT. AV. NET WEIGHT

2 GAL. AV. NET WEIGHT

DIRECTIONS

FOR TERMITES — Mix 1 quart BONIDE CHLORDANE 74%E in 10 gallons water. Buildings Having Basements or Crawl Spaces — Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

JAPANESE BEETLE LARVAE (Federal/State Quarantine Programs — For use on soil and ornamental turf only.) 2.5 - 5 quarts BONIDE CHLORDANE 74%E per acre (9 - 18 ounces per 5,000 square feet). Use higher rate on muck or highly organic soil. Use sufficient water for full coverage. Apply just before a rain or use irrigation or sprinklers to remove material from turf. Keep children and pets off treated turf until material has been washed into the soil by watering or rain, and grass is dry.

4-96
131176