



**WEED & FEED**

**CRANILLES**

**WEED & FEED  
GRANULES**

JUL 21 1967

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 11111 SUBJECT  
TO ATTACHED COMMENTS.

# WEED & FEED GRANULES

LAWN WEED & FEED GRANULES being a combination product, not only kills weeds but also feeds good grasses. This helps good grasses crowd out, by strong healthy growth, undesirable weed pests that may not have been killed by the weed killer present.

WEED & FEED GRANULES may be applied during the spring (after weather has become warm), summer or fall (before freezing temperatures arrive). Fall applications may not show results until following spring and results will be slow when applied during cool periods.



## DIRECTIONS

If rain is expected within 24 hours do not apply. On newly seeded lawns wait until grass has been regularly mowed two or three times. Do not apply during periods of extremely high temperatures 90°F.+ Apply 4 or 5 days after regular mowing when grass is moist from rain, dew, or light watering. Choose calm period to avoid wind drift. Use lawn fertilizer spreader. Measure the area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for 3/8 inch wide openings will work on most spreaders but due to wear and different walking speeds, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. In case of gross overdose watering the area will reduce or prevent damage. Avoid overlapping and double coverage in backing or turning. Close spreader at those times and before filling.

Do not sprinkle or water treated areas for at least 24 hours after application. Do not mow treated areas for 3 days after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.

Dead or dying weeds may be raked out and bare spots reseeded two weeks after application. Soak lawn (heavy rain or watering) 2 or 3 days before reseeding.

## CAUTION

Keep out of reach of children. Avoid contact with skin, eyes or clothing. In case of contact and after use wash thoroughly. Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Tomatoes, peas, grapes, roses, ornamentals, etc., can be injured, avoid application and drift to valuable plantings. Do not use on ground covers such as Pachasandra, Dichondra, etc. Storage in cool dry area but not near fertilizers, seeds, insecticides or fungicides prolongs effectiveness of ingredients. Clean spreader after use.

Spreader Make & Model	Setting	Spreader Make & Model	Setting
Jackson 20-95	10 1/4	Scott 35-8, 75-8	12
Lawn Beauty	10 1/4	Garden Mark	12

WIND DRIFT. Use lawn fertilizer spreader. Measure the distance between contents of this bag to area listed on front of bag. Spreader settings for  $\frac{3}{8}$  inch wide openings will work on most spreaders but due to wear and different walking speeds, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. In case of gross overdose watering the area will reduce or prevent damage. Avoid overlapping and double coverage in backing or turning. Close spreader at those times and before filling.

Do not sprinkle or water treated areas for at least 24 hours after application. Do not mow treated areas for 3 days after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.

Dead or dying weeds may be raked out and bare spots reseeded two weeks after application. Soak lawn (heavy rain or watering) 2 or 3 days before reseeding.

## CAUTION

Keep out of reach of children. Avoid contact with skin, eyes or clothing. In case of contact and after use wash thoroughly. Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Tomatoes, peas, grapes, roses, ornamentals, etc., can be injured, avoid application and drift to valuable plantings. Do not use on ground covers such as Pachasandra, Dichondra, etc. Storage in cool dry area but not near fertilizers, seeds, insecticides or fungicides prolongs effectiveness of ingredients. Clean spreader after use.

<u>Spreader Make &amp; Model</u>	<u>Setting</u>	<u>Spreader Make &amp; Model</u>	<u>Setting</u>
Jackson 20-95	10 $\frac{1}{4}$	Scott 35-8, 75-8	12
Lawn Beauty	10 $\frac{1}{4}$	Garden Mark	12
Sears Cross County 20-26	10	Modern Line 350-360 6-1	7
Cyclone B 5' Swath (Still air only)	6 $\frac{3}{4}$	Precision Products 20F	12 $\frac{1}{2}$
Salem Micro Set	7 $\frac{1}{2}$	Montgomery Ward 89-1081 20"	8 $\frac{1}{2}$

Buyer accepts free of warranty

Manufactured by Bonide Chemical Co., Inc., Utica, N. Y. 13502

Reg. No. 4-114

497-4

FREIGHT SHIPPING BAG  
Meeting requirements of  
APPLICABLE FREIGHT CLASSIFICATION  
Guaranteed by Arkell and Smiths, Hudson Falls, N. Y.

ACCEPTED  
JUL 21 1967  
UNITED STATES DEPARTMENT OF AGRICULTURE  
FEDERAL AGRICULTURAL MECHANIZATION SERVICE  
FOR THE NATIONAL FERTILIZER INSTITUTE  
1000 17th St., N.W., Washington, D.C. 20036  
TO AGRICULTURAL CHEMISTS

## WEED & FEED GRANULES

LAWN WEED & FEED GRANULES being a combination product, not only kills weeds but also feeds good grasses. This helps good grasses crowd out, by strong healthy growth, undesirable weed pests that may not have been killed by the weed killer present.

WEED & FEED GRANULES may be applied during the spring (after weather has become warm), summer or fall (before freezing temperatures arrive). Fall applications may not show results until following spring and results will be slow when applied during cool periods.

## DIRECTIONS

If rain is expected within 24 hours do not apply. On newly seeded lawns wait until grass has been regularly mowed two or three times. Do not apply during periods of extremely high temperatures 90°F.+ Apply 4 or 5 days after regular mowing when grass is moist from rain, dew, or light watering. Choose calm period to avoid wind drift. Use lawn fertilizer spreader. Measure the area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for  $\frac{3}{8}$  inch wide openings will work on most spreaders but due to wear and different walking speeds, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. In case of gross overdose watering the area will reduce or prevent damage. Avoid overlapping and double coverage in backing or turning. Close spreader at those times and before filling.

Do not sprinkle or water treated areas for at least 24 hours after application. Do not mow treated areas for 3 days after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.

Dead or dying weeds may be raked out and bare spots reseeded two weeks after application. Soak lawn (heavy rain or watering) 2 or 3 days before reseeding.



## CAUTION

Keep out of reach of children. Avoid contact with skin, eyes or clothing. In case of contact and after use wash thoroughly. Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Tomatoes, peas, grapes, roses, ornamentals, etc., can be injured, avoid application and drift to valuable plantings. Do not use on ground covers such as Pachasandra, Dichondra, etc. Storage in cool dry area but not near fertilizers, seeds, insecticides or fungicides prolongs effectiveness of ingredients. Clean spreader after use.

<u>Spreader Make &amp; Model</u>	<u>Setting</u>	<u>Spreader Make &amp; Model</u>	<u>Setting</u>
Jackson 20-95	10 $\frac{1}{4}$	Scott 35-8, 75-8	12
Lawn Beauty	10 $\frac{1}{4}$	Garden Mark	12
Sears Cross County 20-26	10	Modern Line 350-360 6-1	7
Cyclone B 5' Swath (Still air only)	6 $\frac{3}{4}$	Precision Products 20F	12 $\frac{1}{2}$
Salem Micro Set	7 $\frac{1}{2}$	Montgomery Ward 89-1081 20"	8 $\frac{1}{2}$

Buyer accepts free of warranty

Manufactured by Bonide Chemical Co., Inc., Utica, N. Y. 13502

. . . . . Reg. No. 4-114

497-4

FREIGHT SHIPPING BAG  
Meeting requirements of  
APPLICABLE FREIGHT CLASSIFICATION  
Guaranteed by Arkell and Smiths, Hudson Falls, N. Y.

BONIDE