


PM-23 4-131 p. 1/2

BEST AVAILABLE COPY



by SLO Cal.  
4-131

*[Handwritten signature]*  
4-131

EPA Est. No. 4-NY-1      EPA Reg. No. 4-131

ACTIVE INGREDIENT:  
 \*Captan ..... 49.0%  
 Related Derivatives ..... 1.0%  
 INERT INGREDIENTS: ..... 50.0%  
 U.S. Pat. Nos. 2,553,770; 2,553,771  
 \*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

**BEST AVAILABLE COPY**

**DANGER**

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Wear chemical resistant gloves when using indoors or outdoors. When using, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product just be laundered separately from other clothing before reusing.

(Cont'd on page 2)

P. 2/3

BONIDE CAPTAN - 50% WP, EPA Reg. No. 4-131

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place protected from excessive heat. Store in an area inaccessible to children or pets.

Container Disposal: Do not reuse empty container. Wrap and put in trash.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

BONIDE CAPTAN-50% WP can be combined with most commonly used insecticides and fungicides, except oil and alkaline materials, that is lime-sulphur, bordeaux mixture or calcium arsenate. Spray injury may result if used with, just before, or closely following an oil spray. At high rates, or in drenching sprays, CAPTAN-50% WP may cause necrotic spotting of tender, immature leaves on certain varieties of apples, peaches, plums and cherries. This is most likely from early cover sprays during long periods of warm, cloudy, humid weather. To avoid this, at such times, use at lowest recommended rates and avoid drenching trees.

Pour recommended amount on water surface of nearly filled spray tank or pre-mix in bucket half filled with water and pour through screen into nearly filled spray tank. Keep agitator running during filling and spraying. Do not allow mixture to stand. Do not use in low volume gear type spray equipment. Do not use with emulsifiable liquids unless previous use proves compatibility. In order that official food and drug residue tolerances not be exceeded use only at recommended rates.

Use a U.S. standard tablespoon to measure lower amounts of BONIDE CAPTAN 50% WP

ORNAMENTALS	DISEASES	WHEN TO APPLY	AMOUNT
AZALEAS	damping off of cuttings	Dip cuttings before bedding.	2-1/5 lbs /gal. or 4 lbs /100 gals. water.
	petal blight	Apply to soil area around plants and spray flowers beginning just before bloom. Repeat at 7 to 14 day intervals through bloom.	1 lbs /gal. or 2 lbs /100 gals. water.
BEGONIAS (Tuberous)	damping off, tuber rot	Dip tubers 30 minutes, drain and plant.	2-1/5 lbs /gal. or 4 lbs /100 gals. water.
	powdery mildew	Begin at first sign of disease. Repeat at weekly intervals as needed.	1/2 to 1 lb /gal. or 1/2 to 1 lb /100 gals.
CARNATIONS	alternating leaf spot, rust	Begin at first signs of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews.	1 lbs /gal. or 2 lbs /100 gals. water.
	damping off of cuttings	Dip cuttings before bedding.	2-1/5 lbs /gal. or 4 lbs /100 gals. water.
CHRYSANTHEMUM	botrytis flower blight, septoria leaf spot	Apply at first sign of disease. Repeat at 7 to 14 day intervals.	1 lb /gal. or 2 lbs /100 gals. water.
	damping off of cuttings	Dip cuttings before bedding.	2-1/5 lbs /gal. or 4 lbs /100 gals. water.
GLADIOLUS (corms)	corm rot and decay, damping off	Dip corms 15 to 30 minutes. Drain and plant.	1/2 to 1 lb /gal. or 1 lb /100 gals.
ROSES	black spot, styx blossom blight	Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals; and more frequently during frequent rains and heavy dews.	1 lbs /gal. or 2 lbs /100 gals. water.
VEGETABLES	DISEASES	WHEN TO APPLY	AMOUNT
CANTALOUPE, CUCUMBERS	angular leaf spot, anthracnose, downy mildew	Begin at first sign of disease. Repeat at 5 to 7 day intervals.	1 1/2 lbs /gal. or 3 lbs /100 gals. water.
TOMATOES	anthracnose, early and late blight, gray leaf spot (Stemphylium), septoria leaf spot	Begin at first sign of disease. Repeat at 5 to 7 day intervals.	2-1/5 lbs /gal. or 4 lbs /100 gals. water.

**BEST AVAILABLE COPY**

BONIDE CAPTAN - 50% WP, EPA Reg. No. 4-131

FRUIT	DISEASES	WHEN TO APPLY	AMOUNT
APPLES	bitter rot, black rot, fraxage leaf spot, black nec. botryosphera (white rot)	Apply in late cover and preharvest periods	1 lbs / gal. or 2 lbs / 100 gals water
	botrytis rot, bullseye rot	Apply in first two cover sprays and in 1 or 2 preharvest sprays or as needed at 5-7 day intervals from prebloom through early cover periods.	
	brooks fruit spot, Nysbeck	Apply in cover periods	1/2 to 2 1/2 lbs / gal. or 1/2 to 2 lbs / 100 gals water
	sooty blotch	Apply in cover periods.	1 lbs / gal. or 2 lbs / 100 gals water
CHERRIES	botrytis rot	Apply at petal fall, shuck shed, and in cover sprays	1 to 2-1/2 lbs / gal. or 2 to 4 lbs / 100 gals water
	blowm blight (brown rot)	Apply during bloom at 3 to 4 day intervals.	1 lbs / gal. or 2 lbs / 100 gals water
	fruit rot (brown rot)	Apply in pink, full bloom, petal fall, shuck shed, and cover period.	
	leaf spot	Apply in pink, bloom, petal fall, and first cover periods, and in 3 or 4 postharvest foliage sprays	
PEACHES	brown rot, rhizopus, scab	Apply in pink, full bloom, shuck shed, cover and preharvest periods.	1 lbs / gal. or 2 lbs / 100 gals water
	brown rot (blowm blight)	Apply at 3 to 4 day intervals during bloom.	
	corymum blight	Apply in pink bud, full bloom, petal fall, and cover periods as necessary, and as a fall foliage spray	
STRAWBERRIES	botrytis rot (gray mold), leaf spots	Begin at new growth. Repeat at 7 day intervals through pick and 2nd after harvest	1 1/2 lbs / gal. or 3 lbs / 100 gals water

Buyer's guarantee limited to label claims.

BONIDE CHEMICAL CO., Ypsilville, K.Y. 13495

**BEST AVAILABLE COPY**