

BONIDE

SEVIN

5% DUST

Insecticide

CONTROLS

MEXICAN BEAN BEETLES, JAPANESE BEETLES, FLEA BEETLES, CABBAGE WORMS, TOMATO HORNWORMS, ARMY WORMS AND CERTAIN OTHER INSECT PESTS OF VEGETABLES, FRUIT, ORNAMENTALS. CHICKENS, PHEASANTS, PIGEONS, PARTRIDGE: FLEAS, MITES, LICE AND TICKS: DOGS AND CATS: FLEAS AND TICKS.

ACTIVE INGREDIENT:

Carbaryl-1 Naphthyl N-Methylcarbamate	5%
INERT INGREDIENTS	95%

Total 100%

CAUTION: "KEEP OUT OF REACH OF CHILDREN"

(NOTE BACK PANEL OTHER CAUTIONS)

©UNION CARBIDE CORP., REG. T.M.

DIRECTIONS FOR USE

Read the Cautions! This is a squeeze duster can, squeeze sides with fingers and thumb to puff out powder as a dust. Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

FOR CONTROL OF

	Time to Stop Use Before Harvest
Beans: Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus and stinkbugs	No Limit
Carrots: Leafhoppers, flea beetles and spittlebugs	No Limit
Corn: Corn earworm, Fall army worm, grasshoppers. (See notes 1 & 2)	Day of Harvest
Cabbage and Cauliflower: Flea beetles, cabbage worms, harlequin bugs, corn earworm	3 days
Brussels Sprouts and Broccoli: Flea beetles, cabbage worms, harlequin bugs, corn earworm	3 days
Cucumbers, Melons and Squash: Pickleworms, squash bugs, flea beetles, cucumber beetles. (See Note 3)	Day of Harvest
Lettuce: Leafhoppers, army worms, stinkbugs	3 days
Okra: Stinkbugs, corn earworm	No Limit
Potatoes: Colorado potato beetle, flea beetle, leafhopper	No Limit
Peppers: Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug	Day of Harvest
Eggplant: Lace bugs, tomato hornworm	Day of Harvest
Tomatoes: Tomato fruitworm, flea beetle, tomato hornworm, stinkbugs	Day of Harvest
Strawberries: Meadow spittlebug, leaf roller and strawberry weevil	1 Day

FOR CONTROL OF

Time to Stop Use Before Harvest

Tobacco: Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed No Limit

Note 1. Do not feed forage to dairy animals or animals being finished for slaughter.

Note 2. Apply to corn silks every 3 days until dry for earworm control.

Note 3. Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80° F. or when vines are wet.

Chickens: Fleas, Lice, Mites and Ticks: One pound of 5% Sevin Dust is enough for one treatment of 100 chickens. It can also be used on pheasants, pigeons and partridge. Dust the chicken house, and roasts for mites and ticks with one pound of 5% Dust per 40 square feet. Do not dust birds within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

Cats and Dogs: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens under 2 months old.

CAUTIONS

HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. Skin Contact With Any Insecticide May Be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note For Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

BUYER ACCEPTS FREE OF WARRANTY

1 LB. NET

USDA Reg. No. 14-X

Made By

BONIDE CHEMICAL CO., INC., UTICA, N. Y.

ACCEPTED
Nov. 13, 1964

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 4-143. SUBJECT TO ALL OTHER COMMENTS.

4-143

11/13/64

1/1