

DIRECTIONS FOR USE

Read the Cautions! This is a squeeze duster can. squeeze sides with fingers and thumb to puff out powder as a dust. Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7 day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

FOR CONTROL OF

Time to Stop Use Before Harvest

Beans: Mexican bean beetle, leafhoppers, flea beetles, corn, earworm, lygus and stinkbugs	No Limit
Carrots: Leafhoppers, flea beetles and spittlebugs	No Limit
Corn: Corn earworm, Fall army worm, grasshoppers, (See notes 1 & 2)	Day of Harvest
Cabbage and Cauliflower: Flea beetles, cabbage worms, harlequin bugs, corn earworm	3 days
Brussels Sprouts and Broccoli: Flea beetles, cabbage worms, harlequin bugs, corn earworm	3 days
Cucumbers, Melons and Squash: Flea beetles, pickleworms, cucumber beetles, squashbugs (See Note 3)	Day of Harvest
Eggplant: Lace bugs, tomato hornworm	Day of Harvest
Head Lettuce: Leafhoppers, stinkbugs army worms	3 days
Okra: Stinkbugs, corn earworms	No Limit
Peppers: Tomato fruitworm, Fall army worm, flea beetles, leafhoppers, lygus bug	Day of Harvest
Potatoes: Colorado potato beetle, flea beetle, leafhopper	No Limit
Strawberries: Meadow spittlebug, leaf roller and strawberry weevil	1 day
Tobacco: Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed	No Limit
Tomatoes: Tomato fruitworm, tomato hornworm, flea beetle, stinkbugs	Day of Harvest

FOR CONTROL OF

Time to Stop Use Before Harvest

- Note 1.** Do not feed forage to dairy animals or animals being finished for slaughter.
- Note 2.** Apply to corn silks every 3 days until dry for earworm control.
- Note 3.** Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80°F. or when vines are wet.

Chickens: Fleas, lice, mites and ticks: One pound of 5% Sevin Dust is enough for one treatment of 100 chickens. It can also be used on pheasants, pigeons and partridge. Dust the chicken house and roosts for mites and ticks with one pound of 5% dust per 40 square feet. Do not dust or house birds in treated houses within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

Cats and Dogs: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens or puppies under 4 weeks of age. Do not repeat application within 7 days.

Ornamentals: Flowers & herbaceous plants (Japanese, blister, flea, June beetles; boxelder, mealy, lace & plant bugs; leafhoppers, leaf roller, psyllid, rose aphid & exposed thrips) dust thoroughly covering both tops & undersides of leaves with light film. Repeat application as needed. Do not apply on Boston Ivy.

Shrubs, Trees & Woody Plants: (gypsy moth, scale insects, bag worm, birch, boxwood, and oak leaf miners, elm leaf aphid, Japanese, June, elm leaf & willow leaf beetles, mimosa webworm, tent & puss caterpillars, rose aphid & slug, periodical cicada, leafhopper, leaf roller, boxelder, lace, mealy, plant and thorn bugs, apple aphid, orange tortrix and exposed thrips) dust thoroughly covering both tops & undersides of leaves with light film. Repeat application as needed.

CAUTIONS

HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. **Skin Contact May Be Harmful.** Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work. **Wear Regular Long Sleeved Work Clothing.** Change to Clean Clothing Daily. Note For Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

BUYER'S GUARANTEE LIMITED TO LABEL CLAIMS

MANUFACTURED BY

BONIDE CHEMICAL CO., INC., UTICA, N.Y. 13502

BONIDE

SEVIN

5% DUST

Insecticide for use on

**VEGETABLES, FRUIT, ORNAMENTALS,
PHEASANTS, PIGEONS, PARTRIDGE,
DOGS AND CATS.**

ACTIVE INGREDIENT: Carbaryl (1-naphthyl N-methylcarbamate)
INERT INGREDIENTS

USDA REC

Union Carbide Corp. Reg. T.M.

Tot

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions)

ACCEPTED WITH COMMENTS

Time to Start Use Before Harvest

forage to dairy animals or unfinished for slaughter.

in silks every 3 days until dry for control.

on watermelon in Florida. Do not use on melons or cucumbers when temperature is below 80°F. or when vines are wet.

lice, mites and ticks: One pound enough for one treatment of 100 birds. Do not be used on pheasants, pigeons in chicken house and roosts for more than one pound of 5% dust per 40 pounds of birds in treated area. Avoid dusting of birds before slaughter. Avoid dusting of birds before use on pet birds.

Fleas and ticks on cats and dogs. Do not use on cats' eyes. Dust quarters weekly. Do not use on puppies under 4 weeks of age. Repeat application within 7 days.

Flowers & herbaceous plants (Japanese beetles; boxelder, mealy, lace bugs, leaf roller, psyllid, rose hoppers, leaf miner, etc.) dust thoroughly covering both sides of leaves with light film. Repeat application as needed. Do not apply on Boston Ivy.

Woody Plants: (gypsy moth, scale, birch, boxwood, and oak leaf miner, Japanese, June, elm leaf miner, mimosa webworm, tent & puss moth, slug, periodical cicada, leaf miner, etc.) dust thoroughly covering both tops & undersides of leaves with light film. Repeat application as needed.

CAUTIONS

DO NOT INHALE OR SWALLOWED. Do not get on face. Do Not Take Internally. Avoid Repeated Breathing of Dust. Contact May Be Harmful. Avoid Contact with Eyes. Wash Hands Before Eating. Take Shower After Work. Wear Regular Work Clothing. Change to Clean Clothing After Work. Note For Physician: Carbaryl is reversible, cholinesterase inhibitor. Antidote: Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

REGISTERED TRADEMARKS

Union Carbide Corp., N.Y. 13502

BONIDE

SEVIN

5% DUST

Insecticide for use on

**VEGETABLES, FRUIT, ORNAMENTALS, CHICKENS,
PHEASANTS, PIGEONS, PARTRIDGE,
DOGS AND CATS.**

ACTIVE INGREDIENT:	USDA REG. NO. 4-143
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
	100%
Union Carbide Corp. Reg. T.M.	Total

CAUTION: KEEP OUT OF REACH OF CHILDREN.

(See back panel cautions)

4 LB. NET