

pm 04 4-143 2/12/98

Page 175

**ACCEPTED**  
FEB 12 1998  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-143

FRONT PANEL

BONIDE  
SEVIN<sup>®</sup> 5% DUST

Brand Carbaryl Insecticide

Ready to Use

(PICTURE)

Insecticide for use on  
vegetables, fruit, ornamentals, dogs and cats

EPA Est. No. 4-NY-1

EPA Reg. No. 4-143

ACTIVE INGREDIENT:


Carbaryl (CAS# 63-25-2).....	5.0%
OTHER INGREDIENTS:.....	<u>95.0%</u>
TOTAL.....	100.00%

Sevin<sup>®</sup> is a registered trademark of Rhone-Poulenc Nederland B.V. for carbaryl insecticides.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Net Contents: 1 lb



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS--CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling. Wash contaminated clothing before reuse. NOTE TO PHYSICIAN; Carbaryl is a moderate, reversible cholinesterase inhibitor. ATROPINE is antidotal. NOTE TO VETERINARIAN: Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to bees and aquatic organisms. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply while bees are actively visiting the area. Additional protective information may be obtained from your Cooperative Extension Service. Subsequent planting of food and feed crops in treated soils must be made with crops on this or other carbaryl labels.

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.

Change clothing as soon as possible after use.

Wash the outside of gloves before removing.

As with any pesticide product, wash hands thoroughly, immediately after handling and before eating, smoking or using toilet.

Avoid contamination of food and feed.

Do not allow children or pets to contact treated area until dust has settled.

395

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun puffer or bulbous duster, rotary duster rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above-chest height, consider using a liquid carbaryl product. Do not repeat applications more than once a week.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area, inaccessible to children.  
DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

This product has been colored green to blend with foliage and avoid unsightly white dust deposits.

Not for use on plants being grown for sale, or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic, or climatic modification and being grown in interior plantscapes, ornamental gardens, or parks, or on golf courses, or lawns and ground.

READ THE CAUTIONS! Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7-day intervals, more often in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects. NOTE 1: Do not feed forage to dairy animals or animals being finished for slaughter. NOTE 2: Apply to corn silks every 7 days until dry for corn earworm control NOTE 3: Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80° F or when vines are wet.

FOR CONTROL OF:		TIME TO STOP USE BEFORE HARVEST
BEANS:	Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus bugs, stink bugs. . .	No Limit
CARROTS:	Leafhoppers, flea beetles, spittlebugs . . . . .	No Limit
CORN:	Corn earworm, Fall armyworm, grasshoppers (See Notes 1&2). . . . .	Day of Harvest
BRUSSELS SPROUTS:	Flea beetles, cabbage worms, harlequin bugs, corn earworm . . . . .	3 days
BROCCOLI:	Flea beetles, cabbage worms, harlequin bugs, corn earworm . . . . .	3 days
CUCUMBERS:	Flea beetles, pickelworms, cucumber beetles, squash bugs (See Note 3). . . . .	Day of Harvest
MELONS:	Flea beetles, pickelworms, cucumber beetles, squash bugs (See Note 3). . . . .	Day of Harvest
SQUASH:	Flea beetles, pickelworms, cucumber beetles, squash bugs (See Note 3). . . . .	Day of Harvest
EGGPLANT:	Lace bugs, tomato hornworm . . . . .	Day of Harvest
HEAD LETTUCE:	Leafhoppers, stink bugs, army worms. . . . .	3 days
OKRA:	Stink bugs, corn earworm . . . . .	No Limit
PEPPERS:	Tomato fruitworm, Fall armyworm, flea beetles, leafhoppers, lygus bugs . . . . .	Day of Harvest
POTATOES:	Colorado potato beetle, flea beetles, leafhoppers. . . . .	No Limit
STRAWBERRIES:	Meadow spittlebug, leaf rollers, strawberry weevil. . . . .	1 day
TOMATO:	Tomato fruitworm, tomato hornworm, flea beetles, stink bugs . . . . .	Day of Harvest

**READ ENTIRE LABEL BEFORE EACH USE  
ONLY ON DOGS AND CATS**

DOGS and CATS: Fleas and ticks on dogs and cats. Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Do not use on pregnant dogs or cats. Do not use on kittens or puppies under 12 weeks of age. Dust bedding and doghouse using a shaker can or other dust applicator. Do not repeat application more than once a week. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. In case of Emergency, call 1-800-548-2423 (Animal Poison Control Center). Consult a veterinarian before using this product on debilitated, aged, nursing or medicated animals.

595

ORNAMENTALS: Flowers and herbaceous plants (Japanese beetle, blister beetle, flea beetles, June beetles, boxelder bug, mealybugs, lacebugs, plant bugs, leafhoppers, leaf rollers, psyllids, rose aphid and exposed thrips) dust thoroughly covering both tops and undersides of leaves with light film. Repeat application as needed no more than once a week. Do not apply on Boston Ivy.

SHRUBS, TREES and WOODY PLANTS: (Gypsy moth, scale insects, bagworm, birch leafminer, boxwood leafminer, oak leafminers, elm leaf aphid, Japanese beetle, June beetles, elm leaf beetle, willow leaf beetles, mimosa webworm, tent caterpillars, puss caterpillar, rose aphid, rose slug, periodical cicada, leafhoppers, leafrollers, boxelder bug, lace bugs, mealybugs, plant bugs, thorn bug, apple aphid, orange tortrix and exposed thrips), dust thoroughly covering both tops and underside of leaves with light film. Repeat application as needed but not more than once a week.

BEE, WASP and Hornet: Using the 10oz duster with snorkel tube attachment, locate the brood nest or center of the colony, if possible by listening with your ear to the wall (inside or out). Where the buzzing and clicking sound is the loudest is generally the area of the brood comb and bees.

Drill a 1/4" hole into that section of the wall from either the inside or outside and push the bit on through to clear a hole into the brood area.

Insert the snorkel tube of the Sevin®5% duster into the 1/4" hole and squeeze the duster, thus pumping the dust into the brood area. Continue squeezing until most of the insecticide dust is pumped in the wall cavity. If dust packs in the snorkel tube, remove the snorkel tube and the nozzle and shake to loosen dust. Allow 24 hours for Sevin®5% dust to completely destroy the colony after which the 1/4" hole and the outside bee entrance can be permanently sealed with wood putty or some other similar impervious material.

If it is impossible to drill a hole into the brood nest, apply Sevin®5% dust heavily around the outside entrance to the colony. Also, pump a large quantity of Sevin®5% dust into the entrance. This method will take longer to destroy the colony and may require two or three applications in order to do so.

Sevin®5% dust will also destroy wasps and hornets but heavy protective clothing, including veil and gloves, should be worn in applying Sevin®5% to the nest, as the hornets will attack savagely before the Sevin®5% dust can be applied effectively.

Stand upwind while pumping Sevin®5% dust into the entrance of the colony. Do not eat any honey taken out of the wall after bees have been destroyed with Sevin®5% dust.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call National Pesticide Telecommunications Network at 1-800-858-7378.

BUYER,S GUARANTEE LIMITED TO LABEL CLAIMS

MANUFACTURED BY:

BONIDE PRODUCTS, INC. YORKVILLE, NY 13495