HOW INSECTS ARE KILLED:

Protects plants, such as roses, chrysanthemums, azaleas, pyracantha, spirea, birch, and other flowers and shrubs from damage by aphids, spider mites, thrips, leaf miners, whiteflies, lace bugs, leafhoppers, SYSTEMIC GRANULES enter a plant through its roots (systemic action) and then moves to all parts of the plant through natural growth and sap flow. Working from inside of the plant, it kills sucking insects that feed on the plant.

HOW TO USE:

Apply the ready-to-use granules uniformly (see Recommended Applications) to the soil around the base of the plant. Mix the granules into the top soil, then water. The chemical, released from the granules, will leach into the root zone and be absorbed into the plant. The granules can also be scattered over planting beds and mixed into the top soil beneeded to maintain insect control after the plants start to grow.

WARNING:

Poisoning may occur as a result of exposure. Wear rubber gloves. May be fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through Rapidly absorbed through In case of contact, remove contaminated clothing and wash skin immediately with soap and water. Do not get in eyes. Do not contaminate feed or food. Keep children and pets away from treated area until product is worked and watered into soil and treated area is completely dry. Antidote: Atropine is antidotal.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Birds and other wildlife in treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Dispose of by burying or burning.


Do not apply broadcast applications to lawns.

RESTRICTION:

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Buyer's Guarantee Limited to Label Claims

Packed and Distributed by BONIDE CHEMICAL CO., INC., Yorkville, N.Y. 13495



RECOMMENDED APPLICATIONS

FOR FLOWERS

TO CONTROL Aphids (plant lice) Lace bug

Carnation Leafhoppers Chrysanthemum Spider mites Delphinium Thrips Whiteflies Petunia

AMOUNT TO USE

BEFORE PLANTING: Apply 16 ounces per 100 square feet (5 x 20 ft.) and work into soil. Sow seed or set plants in normal manner and water thoroughly.

GROWING PLANTS: For Individual Plants: Apply 1 level teaspoonful around base of plant, work into soil and water thoroughly.

FOR FLOWERS IN A ROW: Apply 2 ounces to each 6 feet of row, in a narrow band near base of plants. Work into top soil and water thoroughly, Repeat application at 4-to-6-week intervals if insects reappear.

POTTED PLANTS: Apply one level teaspoonful per 6-inch pot. Mix thoroughly into top soil and drench with water.

FOR SHRUBS

Birch Camellia Euonymus Moraine locust Pine Rhododendron Roses

TO CONTROL Aphids (plant lice) Birch leaf miner Lace Bug Leafhoppers Mimosa webworm Spider mites Pine tip moth Thrips Whiteflies

AMOUNT TO USE

Shrubs 1-3 ft. tall, 1 oz. per foot of height. Shrubs 3-6 ft. tall, 2 ozs. per foot of height. Shrubs 6-12 ft. tail, 3 ozs. per foot of height.

Apply granules uniformly over the area under the shrub canopy; work Into soil and water thoroughly. Repeat application at 4-to-6-week intervals, if required.

ACCEPTED

OCT 11 1974

UNDER THE FEDERAL MASCOTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NO. TO ATTACHED COMMENTS.