

BONIDE

WEED SEEDLING KILLER

**A New Easy Way To Control Weeds
& Grasses As They Germinate In
Flower & Ornamental Plantings**

ACTIVE INGREDIENTS: USDA Reg. No. 4-186
*S-ethyl dipropylthiocarbamate 2.5%
INERT INGREDIENTS: 97.7%

U.S. Pat. No. 2,913,927 TOTAL 100.0%
*Eptam is the registered trademark of Stauffer Chemical Co.
for above named chemical.

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions)

NET WEIGHT 4 LBS. *Treats up to 100 sq. ft.*

INSTRUCTIONS FOR USE

WEED SEEDLING KILLER

WEED SEEDLING
KILLER

Bonide Weed Seedling Killer is a new Chemical "hoe" designed to save hours of tedious hand weeding, plus the additional benefits of neater, cleaner flower beds and borders.

One application will keep a treated area virtually weed free for an average growing season (in semi-tropical areas, more than one application may be needed). **Weed Seedling Killer** controls over 20 species of weeds including annual bluegrass, barnyard grass, crabgrass, common chickweed, foxtails, florida pusley, goosegrass, henbit (deadnettle), lambsquarters, lovegrass, purslane, pigweed, sand bur, nutgrass, bermudagrass, and quackgrass.

Weed Seedling Killer is suitable for use around:

Herbaceous Plants: (ageratum, allysum, asters, amaranthus, balsam, begonia, mums, dahlia, daylillies, dianthus, marigold, nasturtium, pansy, petunia, zinnia).

Ground Covers: (ajuga, nazania, hypericum, ice plant, ivy, pachysandra, periwinkle, sedum, ornamental strawberry)

Evergreens and Deciduous Trees & Shrubs: (berberis, boxwood, camellia, chamaecyparis, citrus, dogwood, euonymus, fir, hemlock, American and Japanese holly, juniper, leucothoe, lilac, linden, magnolia, maple, oak, pieris, pine, podocarpus, rhododendron, rose, spruce, viburnum, taxus yew.)

DIRECTIONS FOR USE

Bonide Weed Seedling Killer can be used anytime during the growing season to prevent weeds from growing. Any weeds that are growing in the area to be treated should be removed before application as it will NOT kill weeds already growing.

FOR HERBACEOUS PLANTS — Apply after growth is 3-5 inches high or two weeks after transplanting.

GROUND COVERS — Apply after growth starts in spring or two weeks after transplanting.

FOR TREES AND SHRUBS — Apply after growth starts in spring or two weeks after transplanting. For control of nutgrass, bermudagrass and quackgrass around established trees and shrubs, existing stands of these perennial grasses must be pulled or hoed out before application. (1 lb. to 160 sq. ft.)

The soil to be treated should be loose and free of clods. Apply by gently shaking the container over the area to be treated. Hold the container about 3 feet above the soil. (1 lb. to 200 sq. ft.) Immediately (within minutes) mix granules thoroughly into the soil using hand rake or cultivator to a depth of 2-3 inches. May also be mixed into the soil by watering for 3-5 minutes or until the soil is wet to a depth of 2 inches.

Additional cultivation to a depth of 1-1½ inches to promote water penetration will not destroy effectiveness.

NOTE: **Weed Seedling Killer** should be used only for recommended purposes and at recommended rates (do not overdose). May cause injury to ornamentals under certain soil and climatic conditions, or if directions are not followed.

CAUTION: Keep out of reach of children. Harmful if swallowed. Wash thoroughly after use. Avoid contact with skin and clothing. Avoid contamination of food, feedstuffs, drinking water, fish ponds, streams and lakes.

Buyers Guarantee Limited to Label Claims

Made by

BONIDE CHEMICAL CO., INC.

UTICA, NEW YORK 13502

ACCEPTED

FOR THE PURPOSES OF THE FEDERAL
FOOD, DRUG AND COSMETIC ACT
AS AMENDED BY THE
FEDERAL ACQUISITION REGULATORY
ADMINISTRATIVE ACT
FEDERAL REGISTER NO. 4