

pm 19

4-157

page 174

BONIDE

OLD PRO MULTI-PURPOSE SPRAY CONCENTRATE FOR NAMED

VEGETABLES & FRUIT TREES

ORNAMENTALS AND FLOWERS

SPRAY

**CONTROLS 100 INSECTS WHICH
ATTACK NAMED FRUITS,
VEGETABLES, ORNAMENTALS &
FLOWERS—SEE SIDE PANEL**

EPA Est. No. 4-NY-1

EPA Reg. No. 4-157

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	13.5%
Malathion: 0,0-diethyl phosphorodithioate of diethyl mercaptosuccinate	13.5%
INERT INGREDIENTS:	73.0%

This product contains 1 lb. Carbaryl and
1 lb. Malathion per gallon at 70° f.

ACCEPTED
FEB 10 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 4-157

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back panel for additional
precautionary statements.

REC'D ENVIRONMENTAL/PPD1
96 NOV -5 AM 1:19

NET CONTENTS:

BONIDE VEGETABLES & FRUIT TREES SPRAY, EPA Reg. No. 4-157

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION

Harmful if inhaled or swallowed. Avoid breathing spray or mist. Do not take internally. Avoid contact with skin, eyes or clothing. In case of contact and before eating and smoking, wash thoroughly with warm water and soap. Note to Physician: Malathion and Carbaryl are moderate, reversible Cholinesterase inhibitors. Atropine is antidotal.

add Page 4
(a.)
(b.)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water.

Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage: Keep container tightly closed to prevent evaporation. Do not store near or contaminate feed or foodstuffs. Do not use, pour, spill or store near heat or open flame. Store in a cool, dry area inaccessible to children and pets.
Disposal: Do not reuse empty container. Wrap container and put in trash collection.

NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES. FOR USE ON PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATION AND BEING GROWN IN INTERIOR PLANTSCAPES, ORNAMENTAL GARDENS OR PARKS, OR ON GOLF COURSES OR LAWNS AND GROUNDS.

add Page 4 (c.) X

BONIDE VEGETABLES & FRUIT TREES SPRAY, EPA Reg. No. 4-157

DIRECTIONS:

Mix at rate of one quart OLD PRO to 25 gallons of water, or 2½ tablespoons per gallon. Choose a cool, calm period preferably early morning or evening when plants will not be subjected to hot direct rays of the sun. Start applications as soon as insects appear and when scale crawlers have settled on the foliage. Make repeat applications as necessary. Apply spray to tops and undersides of leaves, until lightly dripping. Stop applications to fruits and vegetables 3 days before harvest.

(no more than once a week)

BONIDE OLD PRO can be used on 1 apples, 2 pears, 3 peaches, 6 cherries, 7 strawberries, 8 grapes, 9 oranges, 10 lemons, 11 grapefruit, 12 tangerine, 13 beans, 14 cabbage, 15 corn, 16 cucumbers, 17 potatoes, 18 tomatoes, 19 egg plant, 20 peppers, 21 ornamentals and 22 flowers, to control:

codling moth on 1-8 above	clover mites 1-3, 7	Florida red and purple scale	blister beetle 22
green apple aphid on 1-3, 7	European red mites 1-3, 7	9-12	boxelder bugs 21, 22
pear psylla on 1, 2	two spotted mites 1-3, 7	green citrus aphid 9-12	boxwood leaf miner 21, 22
orange tortrix on 1-6,	Willamette mite 1-3	Mexican bean beetle 13	elm leaf aphid 21, 22
9-12, 21, 22	wooly apple aphid 1, 2, 7	aphids 13, 14, 17-20, 21, 22	elm leaf beetle 21, 22
eye spotted bud moth 2, X-6	oriental fruit moth 3	lygus and stink bugs 13, 14, 17	lace bugs 17-20
California pear slug 1, 2	peach twig borer 3-6	-20	oak leaf miner 21, 22
lentiform leaf miner 1, 2	lesser peach tree borer 3-6	bean leaf beetle 13	plant bug 21, 22
San Jose scale 1-3	cat facing bug 3	lima bean pod borer 13	rose aphid 21, 22
Forbes scale 1, 2, X-6	Western spotted cucumber	corn earworm 13, 14	rose slug 22
oystershell scale 1, 2, X-6	beetle 3, 16	army worm 13, 14	willow leaf beetle 21
Lecanium scales 1-7	olive scale 3	velvet bean caterpillar 13	thrips (exposed) 9-12, 15,
white apple leafhopper 1-3, 8	cherry fruit fly X-6	corn earworm 13, 15	21, 22
apple mealy bug 1, 2, 8	black cherry aphid 4-6	European corn borer 15, 17-	gypsy moth 21, 22
apple maggot 1, 2	Western tussock moth 3-6,	20	euonymus scale 21, 22
plum curculio 1-8	9-12	fall armyworm 15, 17-20	white fly 21, 22
rosy apple aphid 1, 2, 7, 21, 22	meadow spittle bug 7	sap beetles 15	tarnished plant bug 13, 14,
red banded leaf roller	strawberry leaf roller 7	flea beetle 13-20	17-20
1-6, 8	grape leaf folder 8	pickleworm 16	rose leaf hopper 22
Eastern tent caterpillar 1, 2,	leaf hopper 8, 13, 14, 16,	melon worm 16	Azalea scale 21, 22
X-6, 21, 22	17-20, 21, 22	imported cabbage worm 14	pine leaf scale 21, 22
fruit tree leaf roller 1-6, 9-12	grape berry moth 8	cabbage looper 14	Magnolia scale 21, 22
periodical cicada 1-3, 21, 22	spider mite 7, 8, 13	Colorado potato beetle 17-20	oak kermes 21, 22
European apple sawfly 1, 2	citrus cutworm 9-12	tomato fruit worm 17-20	Fletcher scale 21, 22
Japanese beetle 1-6, 8, 13,	California red, yellow, purple,	tomato horn worm 17-20	Juniper scale 21, 22
15, 21, 22	black, soft, Citricola scales	tomato russet mite 18	Monterey pine scale 21, 22
bag worm 1, 2, 21, 22	9-12	birch leaf miner 21, 22	
mealy bug 1, 2, 8, 21, 22			

Do not apply to peaches, oranges, lemons, grapefruit, tangerines and cabbage within 7 days of harvest, or to corn within 5 days of harvest. Corn (corn earworm) apply at 2-3 day intervals as silks appear and until silks begin to dry. 3 or more applications may be needed as to duration of silking and severity of infestation.

Buyer's guarantee limited to label claims

Mfg. by Bonide Chemical Co., Inc. • Yorkville, NY 13495
Products

3
4

BONIDE VEGETABLES & FRUIT TREES SPRAY, EPA Reg. No. 4-157

STATEMENT OF PRACTICAL TREATMENT

- (a.) If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.
- If on skin: Wash with plenty of water. Get medical attention if irritation persists.
- If swallowed: Call a doctor or get medical attention. Do not induce vomiting. Do not give anything by mouth to an unconscious person. Avoid alcohol.

USER SAFETY REQUIREMENTS

- (b.) Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

- (c.) Apply when air is calm to avoid drift and contact with eyes and skin.
- Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- For trees taller than 10 feet, consider hiring a licensed professional.
- Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- Do not repeat applications more than once a week.