

Side Panel:

Home Environmental Area Usage: See front panel listing for use with foggers, misters, aerosols and space sprayers such as Aero-Dyne "500", Dynafog, Tifa, Swing Fog, Microsol, Vibro-Fog and others.

To CONTROL: flies, flying insects, moths, gnats, hornets, mosquitoes and wasps.

Indoors: Occupants should vacate the premises before treatment and should not re-occupy until thoroughly ventilated after treatment (opening several doors and windows to create air currents which dissipate the vapors). Use only space or contact sprays indoors; **DO NOT** use fog. Do not apply to skin or animals. Remove birds and pets; cover fish bowls and aquaria before application. Human and pet food and food handling equipment should be removed or covered with wrapping paper. Apply upwards into four corners of the room, 10-15 seconds per 1,000 cu. ft. of room space (average room 15-20 seconds). Close doors and windows for about 15 minutes after treatment. Repeat as necessary. To control hornets and wasps on nests spray nests directly at night.

Outdoors: Best results are obtained in still air. Strong wind dissipates fog too rapidly for good results. Apply 1/4 gallon per acre of lawn and shrubbery -- a thorough application around outdoor areas to repel and kill annoying flying insects. Use dense fog (3-7 second application) around dark, moist shrubbery and insect hiding places for flies, mosquitoes, and wasps. Repeat as necessary. Avoid close direct application to shrubs as spotting or injury may result. Do not apply in areas where food crops may be contaminated.

To Control ants, centipedes, cockroaches, millipedes, silverfish, spiders, and ticks. Apply to cracks of woodwork, wall, floors, around doors, window frames, baseboards, behind shelving of storage areas and other places where these insects may congregate, walk or hide. Apply to ant trails around doors and window sills. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, containers or cooking utensils. Do not treat animals.

To Control fleas & ticks. Apply to animal sleeping quarters freely; force insecticide into cracks in floors and about the room. Apply to animal bedding. Repeat as necessary. Observe "Home Environmental area" cautions above.

To Control bedbugs. Take bed apart and spray entire surface of bedstead, slats, and springs. Spray thoroughly into cracks, and joints in bedstead and on tufts and seams of mattress. Do not spray general surface of mattress. Spray on and under baseboards, moldings, cracks in floors and walls, underneath loose wallpaper and other places where insects may be found. Air mattress until completely dry (at least 4 hours before re-use).

[393]

1-1-60

Use Panel, Cont'd.

Use Control: fly maggots in breeding areas (manure piles and garbage dumps). Apply 1-2 quarts per 100 sq. ft., 10' X 10' area. Do not treat areas where food or animal feed, or crops are growing. Birds feeding on treated areas **may** be killed.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin or eyes. If swallowed, induce vomiting immediately and get **medical attention**. Do not breathe vapor, mist or spray. Do not get in eyes, on skin or clothing; rapidly absorbed through skin or eyes. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get **medical attention**. Wash contaminated clothing with soap and hot water before use. Do not re-use contaminated shoes or boots until free of all contamination. Keep out of reach of children. Do not contaminate animal feed, water or foodstuffs. Wash thoroughly with soap and water after handling and before eating or smoking. Atropine is antidotal.

Toxic to fish and wildlife. Keep out of any body of water. Avoid contamination of backward fish ponds.

Apply this product only as specified on the label. Do not re-use empty container. Destroy it by perforating and crushing. Bury or discard in safe place.

WARRANTY LIMITED TO LABEL CLAIMS.

RONIDE CHEMICAL CO., INC.
Yonkers, New York 10502

[2 of 3]

1-16-69

Side Panel:

Home Environmental Area Usage: See front panel listing for use with foggers, misters, aerosols and space sprayers such as Aero-Dyne "500", Dynafog, Tiff, Swing Fog, Microsol, Vibro-Fog and others.

To CONTROL: flies, flying insects, moths, gnats, hornets, mosquitoes and wasps.

Indoors: Occupants should vacate the premises before treatment and should not re-occupy until thoroughly ventilated after treatment (opening several doors and windows to create air currents which dissipate the vapors). Use only space or contact sprays indoors; DO NOT use fog. Do not apply to skin or animals. Remove birds and pets; cover fish bowls and aquaria before application. Human and pet food and food handling equipment should be removed or covered with wrapping paper. Apply upwards into four corners of the room, 10-15 seconds per 1,000 cu. ft. of room space (average room 15-20 seconds). Close doors and windows for about 15 minutes after treatment. Repeat as necessary. To control hornets and wasps on nests spray nests directly at night.

Outdoors: Best results are obtained in still air. Strong wind dissipates fog too rapidly for good results. Apply 1/4 gallon per acre of lawn and shrubbery -- a thorough application around outdoor areas to repel and kill annoying flying insects. Use dense fog (3-7 second application) around dark, moist shrubbery and insect hiding places for flies, mosquitoes, and wasps. Repeat as necessary. Avoid close direct application to shrubs as spotting or injury may result. Do not apply in areas where food areas may be contaminated.

To Control ants, centipedes, cockroaches, millipedes, silverfish, spiders, and ticks. Apply to cracks of woodwork, wall, floors, around doors, window frames, baseboards, behind shelving of storage areas and other places where these insects may congregate, walk or hide. Apply to ant trails around doors and window sills. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, containers or cooking utensils. Do not treat animals.

To Control fleas & ticks. Apply to animal sleeping quarters freely; force insecticide into cracks in floors and about the room. Apply to animal bedding. Repeat as necessary. Observe "Home Environmental Area" cautions above.

To Control bedbugs. Take bed apart and spray entire surface of bedstead, slats, and springs. Spray thoroughly into cracks, and joints in bedstead and in tufts and seams of mattress. Do not spray general surface of mattress. Spray on and under baseboards, walls, and ceiling. Spray on walls, around base of wall, and in cracks. Do not spray on furniture, drapery, or clothing.

[392]

1-16-69

Side Panel, Cont'd.

To Control: fly maggots in breeding areas (manure piles and garbage dumps). Apply 1-2 quarts per 100 sq. ft., 10' X 10' area. Do not treat areas where food or animal feed, or crops are growing. Birds feeding on treated areas may be killed.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin or eyes. If swallowed, induce vomiting immediately and get medical attention. Do not breathe vapor, mist or spray. Do not get in eyes, on skin or clothing; rapidly absorbed through skin or eyes. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Wash contaminated clothing with soap and hot water before use. Do not re-use contaminated shoes or boots until free of all contamination. Keep out of reach of children. Do not contaminate animal feed, water or foodstuffs. Wash thoroughly with soap and water after handling and before eating or smoking. Atropine is antidotal.

Toxic to fish and wildlife. Keep out of any body of water. Avoid contamination of backyard fish pools.

Apply this product only as specified on the label. Do not re-use empty container. Destroy it by perforating and crushing. Bury or discard in safe place.

BUYERS GUARANTEE LIMITED TO LABEL CLAIMS.

BONIDE CHEMICAL CO., INC.
Utica, New York 13502



Kills and Repels Stable Flies, House Flies, Face Flies, Deer Flies, Horse Flies, Gnats and Mosquitoes

Active Ingredients	
Butoxypolypropylene glycol	50.0%
Technical Methoxychlor*	5.0%
Technical Piperonyl Butoxide**	1.6%
Pyrethrins	0.2%
Petroleum Distillates	33.2%
Inert Ingredients	10.0%
	<hr/>
	100.0%

*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% related reaction products.

**Equivalent to 1.28% (butylcarbityl) (6-propylpiperonyl) ether and 0.32% related compounds.

CAUTION Keep out of reach of children.
See other cautions on back panel.

Net Contents 1 Pint (U.S. Measure)

EPA Reg. No. 59-138



59-138

Insect Spray For Horses

DIRECTIONS FOR USE

Insect Spray for Horses can be effectively used on horses of all breeds and ages. This product must be diluted with water as directed before application.

Dilute 1 1/2 fluid ounces (3 tablespoons) in a pint of water. Thoroughly mix. Spray or sponge on animal to dampen the coat. Give particular attention to areas where flies most often attack such as legs and under belly. Repeat application as required.

For larger quantities of spray follow dilution table.

ACCEPTED
MAY 26, 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 59-138 SUBJECT TO ATTACHED COMMENTS.

Dilution Table	
Insect Spray for Horses	Water
3 fl. oz. (6 tablespoons)	1 quart
1/2 pint	3 quarters
1 pint	1 gallon + 1 quart
1 quart	2 1/2 gallons
1 gallon	10 gallons

Horses may be ridden or worked immediately upon drying following application.

CAUTION
Harmful if swallowed. Remove or cover exposed foods before spraying.
Do not apply to dairy animals.
Do not reuse container. Destroy when empty.

COOPER U.S.A., INC.
Chicago, Illinois 60614 USA
A Wellcome Company

FIEI-NM-WWH


SEVIN DUST

EPA REG. NO. 14782-2

NET WGT. 50 LBS.

DIRECTIONS FOR USE

- ① **APPLES:** Carolina Moth, Green Apple Acker, Red-banded Leaf Roller, Pear Psylla, Apple Maggot, Periodical Cicada — To avoid possible fruit thinning do not apply until 30 days after full bloom. Repeat at 10 day intervals if needed. Do not apply within 1 day of harvest. Use 70-120 pounds per acre.
- ② **CORN:** Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetles — Use 40 pounds per acre. For larvae in whorl and foliage feeders, apply to entire plant. Apply to silks as they appear. Make 3 - 4 applications at 2 - 3 day intervals as infestation warrants. Re-apply immediately after rain. For European corn borer—consult your state agricultural experiment station for proper timing of treatments.
Application to the tassel region of sweet corn during the pollen shed period will seriously reduce bee population.
- ③ **COTTON:** Boll Weevil, Bollworm, Cotton Fleahopper, Lygus Bugs, Thrips, Pink Bollworm — Use 30 to 50 pounds of dust per acre. Apply when insects first threaten damage to crop, and repeat at 5 to 7 day intervals as required to maintain control.



Manufactured by
CARUSO FARM SUPPLY CO., INC.
120 N. WHITE HORSE PINE
HAMMONTON, NEW JERSEY 08037

ACTIVE INGREDIENT:

*Carbaryl (1-Naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS:	95.0%
	<hr/>
	100.0%

*Sevin, U. S. Pat. Nos. 2,903,478; 3,009,855

CAUTION Keep Out of Reach of Children

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes. Do not burn. Do not reuse container.

DIRECTIONS FOR USE (CONT.)

④ **SORGHUM (GRAIN AND FORAGE):** Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers, Sorghum Midge - Use 30 to 40 pounds per acre. For optimum control on grain sorghum direct dust into forming heads. For sorghum midge control, make applications 3 to 4 days after heads have emerged from boot. Do not apply within 21 days of harvest of grain. May be applied to day of harvest for sorghum forage.

⑤ **TOBACCO:** Hornworms, Budworms, Tobacco Flea Beetles, Grasshoppers, Japanese Beetles - Use at the rate of 30 to 50 pounds of dust per acre in broadcast treatment. Full coverage is essential. Begin treatments when worms are small. Repeat as needed. Use the highest dosage on large plants. If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Do not apply within 3 days of priming or cutting.

*Trade-mark of Union Carbide Corporation for the active ingredient carbaryl.