

ACCEPTED

MAY 0 2 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under

EPA Reg. No. 4-165

BLUEBERRY	Cherry fruitworm, cranberry fruitworm, plum conculion, leathoppers	14 days	Mix 1 1/2 lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply when insects first appear. Repeat application at 7-10 day intervals as in- festation warrants.
GRAPE	Flea beetles, Japanese beetles, grape leaf skeleton- izers, leathoppers, rose chalcids, Grape berry moths	14 days	Mix 1 1/2 lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply as a post blossom cover spray and repeat 10 days later; make third and fourth application at 10 day intervals if necessary.
RASPBERRY	Japanese beetles, rose chalcids	3 days	Mix 1 1/2 lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply when insects first appear. Repeat application at 7-10 day intervals as in- festation warrants.
STRAWBERRY	Flea beetles, scirtid bugs, strawberry leaf beetles, strawberry weevils	3 days	Mix 1 1/2 lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply as a thorough cover spray to upper and lower surfaces of foliage and vegetable. Begin treatment at first sign of insect infestation and repeat 7-10 day intervals as infestation warrants.
BEANS	Bean leaf beetles, corn earworms, fall armyworms, Japanese beetles, leathoppers, Mexican bean beetles	3 days	
CABBAGE	Imported cabbage worms, flea beetles	3 days	
CALCUTROWER	Imported cabbage worms, flea beetles	7 days	
EGGPLANT	Flea beetles, leathoppers	1 day	
PEAS	Pea weevils	7 days	
PEPPER	Flea beetles	None	
POTATO	Blister beetles, flea beetles, Colorado potato beetles, leathoppers	None	
TOMATO	Blister beetles, fall armyworms, flea beetles, tomato fruitworms	1 day	

AREA MOSQUITO SPRAY: To aid in the control of mosquitoes, gnats around home grounds, patios and picnic areas. Mix 4 teaspoons per gallon water (2 oz. in 3 gallons or 2 quarts in 100 gallons water per 5 acres). Do not apply more than 13 fluid ounces of Methoxychlor 25E or 20 gallons of spray solution per acre (0.2 lbs. actual Methoxychlor) per 5 acres. Apply as a coarse spray of resting surfaces, such as grass, shrubs and tree trunks and branches to a height of 8 to 10 feet. Thoroughly wet plant material so not apply to point of run off on the soil. Do not spray patio surfaces. Keep children and pets off treated areas until spray dries. May also be applied to the same areas with a fog or mist applicator, using 1 part concentration to 4 parts of water. Best results are obtained by timing application to coincide with greatest mosquito activity in early morning or night. Treat as large an area as possible. Repeat as necessary.

Buyer's guarantee limited to label claims

Mfg. by BONIDE CHEMICAL CO., INC., Yorkville, N.Y. 13495

BONIDE®**METHOXYCHLOR
25% E
INSECTICIDE**For control of mosquitoes, insect pests of shade
trees, flowers, ornamentals and vegetables.

ACTIVE INGREDIENTS: EPA Est. 4-RY-1 EPA Reg. No. 4-165

Methoxychlor, technical	25%
Xylene	63%
Heavy Aromatic Naphtha	7%
INERT INGREDIENTS:	5%

*equivalent to 22% of 2,2-bis(4-methoxyphenyl)-1,1,1-trichloroethane and 3% of other isomers and related compounds. Contains 2 lbs. methoxychlor technical per gallon.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS -- CAUTION**

Keep out of reach of children. Harmful if swallowed or inhaled. Avoid breathing of spray mist. Avoid contact with skin, eyes. In case of contact, wash immediately with soap and water -- may be absorbed through skin. Wash thoroughly after using. Do not use on pets or humans. Keep children and pets off treated areas until washed into soil and the grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed do not induce vomiting. Call a physician immediately.
If Inhaled remove to fresh air.
If on Skin wash immediately with soap and water.
If in Eyes wash with plenty of water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to bees and aquatic organisms. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply while bees are actively visiting the area. Additional protective information may be obtained from your Cooperative Extension Service.

PHYSICAL & CHEMICAL HAZARD: S

Flammable. Keep away from fire sparks and heated surfaces.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Store tightly sealed in a cool, dry place, and in an area inaccessible to children or pets. Do not reuse container. Wrap in paper and discard in trash. Avoid contamination of food and feedstuffs.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel cautions.

 ONE QUART ONE GALLON

DIRECTIONS

IMPORTANT: Users should observe the usual precautions concerning application of oil containing products to crops during high temperatures, since many crops are then more susceptible to oil injury. Do not use this product with spray materials incompatible with oil. Do not apply within 14 days of sulphur product application.

BONIDE METHOXYCHLOR 25E INSECTICIDE contains 2 pounds Methoxychlor per gallon and is to be used in a mixture with water for spray control of insects shown on label. It is suitable for high or low volume application. Use this product only according to directions given on the label.

SHADE TREES, ORNAMENTALS: To control Cankerworms, gypsy moth, tussock moth, hemlock looper, imported willow leaf beetle — Mix 1½ to 2 tablespoonfuls per gallon of water (4-6 oz. per 6 gallons or 2-3 quarts in 100 gallons per acre). Apply to thoroughly cover all plant surfaces when small larvae begin feeding.

FLOWERS: To control rose chafer, leafhopper, spittlebug, Japanese beetle, lace bug, flea beetle — Mix 1½ to 2 tablespoonfuls per gallon of water (4-6 oz. per 6 gallons or 2-3 quarts in 100 gallons per acre). Apply to thoroughly cover all plant surfaces when small larvae begin feeding.

CROP	PEST	INTERVAL (Last spray to Harvest)	DIRECTIONS
APPLE	Apple maggots	7 days	Mix 1½ lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Spray all surfaces of foliage and fruit when adults first appear. Repeat at 7-10 day intervals as long as adults are present.
	Codling moths, Japanese beetles, leafhoppers, Plum curculio	7 days	Mix 1½ lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply first spray at petal fall and make two or more additional applications at 7-10 day intervals if infestation warrants.
APRICOT	Plum curculio	21 days	
CHERRY	Plum curculio	7 days	
	Cherry fruit flies, Cherry fruit worms	7 days	Mix 1½ lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply as a thorough cover spray when insects first appear. Repeat at 7-10 day intervals as long as adults are present.
PEACH	Plum curculio	21 days	
PEAR	Codling moths, pear slug	7 days	Mix 1½ lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply as a thorough cover spray at petal fall and make two or more additional applications at 7-10 day intervals if infestation warrants.
PLUM & PRUNE	Plum curculio	7 days	
BLACKBERRY	Flea beetles, Japanese beetles, rose chafers, strawberry weevils	3 days	Mix 1½ lbs. per gal. water (4 oz. per 6 gals. or 2 qts. in 100 gals. per acre). Apply when insects first appear. Repeat application at 7-10 day intervals as infestation warrants.