



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 24 1998

Anne Montalone
Bonide Products Inc.
2 Wurz Ave.
Yorkville, N.Y. 13495

Subject: Revised Labeling in Response to EPA Letter of 12/29/97
Bonide Methoxychlor 25% E Insecticide
EPA Registration No. 4-165
Received at EPA January 13, 1998

Dear Ms. Montalone,

This letter acknowledges receipt of the subject labels, sent in response to the EPA letter of December 29, 1997. The labels are acceptable provided that the following correction is made:

Under page 3, Flowers, the directions should read "To control rose chafer, leafhopper, spittlebug, Japanese beetle, lace bug, flea beetle--Mix 1 ½ to 2 tablespoonfuls per gallon of water. Apply to thoroughly cover plant surfaces when small larvae begin feeding."

A stamped copy of the approved label is enclosed for your files.

Sincerely,

A handwritten signature in cursive script that reads "Elizabeth T. Haerberer".

Elizabeth T. Haerberer
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

BONIDE

METHOXCHLOR 25%E INSECTICIDE

For Control of insect pest on shade trees, flowers, ornamentals
and vegetables

EPA Reg. No. 4-165

EPA EST. No. 4-NY-1

ACTIVE INGREDIENTS:

Methoxychlor: cas # 72-43-5.....	25%
INERT INGREDIENTS:	75%
TOTAL.....	100%

Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

CAUTION

see back panel for additional precautionary statements


NET CONTENTS: 1 pint

ACCEPTED
with COMMENTS
In EPA Letter Dated:

2/24/98 FEB 24 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

4-165



Crop	Pest	Interval (Last spray to Harvest)	Directions
APPLE	Apple Harvest	7 Days	Mix 1½ tbs. per gal. water. Spray all surfaces of foliage and fruit when adults first appear. Repeat at 7-10 day intervals as long as adults are present.
	Codling moths, Japanese beetles, leafhoppers, plum curculios	7 days	Mix 1½ tbs. per gal. water. Apply first spray at petal fall and make two or more additional applications at 7-10 day intervals if infestation warrants.
APRICOT	Plum curculios	21 days	
CHERRY	Plum curculios	7 days	
	Cherry fruit flies, Cherry fruit worms	7 days	Mix 1½ tbs. per gal. water. Apply as a thorough cover spray when insects first appear. Repeat at 7-10 day intervals as long as adults are present.
PEACH	Plum curculios	21 days	Mix 1½ tbs. per gal. water. Apply as a thorough cover spray at petal fall and make two or more additional applications at 7-10 day intervals if infestation warrants.
PEAR	Codling moths, pear slugs	7 days	
PLUM & PRUNE	Plum curculios,	7 days	Mix 1½ tbs. per gal. water. Apply as a thorough cover spray at petal fall and make two or more additional applications at 7-10 day intervals if infestation warrants.
BLACKBERRY	Japanese Beetles, rose chafers, strawberry weevils, flea beetles	3 days	Mix 1½ tbs. per gal. water. Apply when insects first appear. Repeat application at 7-10 day intervals if infestation warrants.
BLUEBERRY	Cherry fruitworms, cranberry fruitworms, Plum curculios, leafhoppers	14 days	Mix 1½ tbs. per gal. water. Apply when insects first appear. Repeat application at 7-10 day intervals if infestation warrants.
GRAPE	Flea beetles, Japanese beetles, Grape leaf skeletonizers, leafhoppers, rose chafers	14 days	Mix 1½ tbs. per gal. water. Apply when insects first appear. Repeat application at 7-10 day intervals if infestation warrants.
	Grape berry moths	14 days	Mix 1½ tbs. per gal. water. Apply as a post-blossom cover spray and repeat 10 days later: make third and fourth applications at 10 day intervals if necessary.

RASPBERRY	Japanese beetles, rose chafers	3 days	Mix 1½ tbs. per gal. water. Apply when insects first appear. Repeat application at 7-10 day intervals if infestation warrants.
STRAWBERRY	Flea beetles, spittlebugs, strawberry leaf beetles, strawberry weevils	3 days	
BEANS	Bean leaf beetles, corn earworms, fall armyworms, Japanese beetles, leafhoppers, Mexican bean beetles	3 days	Mix 1½ tbs. to 2 tbs. per gallon water. Apply as a thorough cover spray to upper and lower surfaces of foliage and vegetable. Begin treatment at first sign of insect infestation and repeat at 7-10 day intervals as infestation warrants.
CABBAGE	Imported cabbage worms, flea beetles	3 days	
CAULIFLOWER	Imported cabbage worms, flea beetles	7 days	
EGGPLANT	Flea beetles, leafhoppers	1 day	
PEAS	Pea weevils	7 days	
PEPPER	Flea beetles	7 days	
POTATO	Blister beetles, flea beetles, Colorado potato beetles, Leafhoppers	None	
TOMATO	Blister beetles, fall armyworms, flea beetles, tomato fruitworms	1 day	

Buyer's Guarantee Limited to Label Claims

Manufactured by:
Bonide Products, Inc.
Yorkville, NY 13495

