MIXING TABLE FOR CONTROL OF SCALE INSECT AND MITE EGGS

AMOUNTS OF OIL & LIME SULPHUR ADDED TO AMOUNT OF WATER

5 tablespoons	(2½ oz.)	ONE GALLON
8 ounces	(½ pint)	THREE GALLONS
16 ounces	(1 pint)	SIX GALLONS
2 GALLONS	(—)	ONE HUNDRED GALLONS

TO CONTROL OVERWINTERING PLANT DISEASES such as black spot, rust blights, leaf spots on dormant deciduous roses, ornamentals, shade trees, apples, plums, prunes and quinces mix and apply at 4½ times above table rates. For currants, gooseberries, grapes and pears at 7½ times above table rates.

Do not store mixtures, make fresh for each use. Keep container tightly closed when not in use. Keep this product in this labelled container until used. When container is empty, wash and destroy; never reuse. Do not use food or drink containers as measuring or transfer units.

CAUTION: Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist or contact with skin, eyes and clothing. In case of contact, and after use, wash thoroughly. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Buyer's Guarantee Limited to Label Claims

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N. Y., 13502

ACCEPTED
MAY 7 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT

BONIDE

OIL & LIME SULPHUR SPRAY

Kills certain overwintering insects and diseases on dormant shrubs, roses, trees and fruit trees.

ACTIVE INGREDIENTS:
Calcium Polysulfides
Petroleum Oils
INERT INGREDIENTS
Petroleum oils minimum unsulfonated residue 92%
USDA Reg. No. 4-100-
5-0[illegible]
80-0[illegible]
15-0[illegible]

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See side panel, other cautions)
NET CONTENTS 1 PINT LIQUID

Oil & Lime Sulphur Spray is a combination of insecticidal oil and fungicidal sulphur for use in controlling rose scale, San Jose scale, black and brown scales, Italian pear scale, spider mite eggs, mealy bugs, and overwintering fungus spores or plant diseases such as black spot, rust blights and leaf spots on dormant deciduous, roses, ornamentals, shade trees, almond, apple, cherry, nectarine, peach, plum, walnut trees and raspberry bushes.

DIRECTIONS

Apply in fall, winter or spring only when plants are completely dormant. Prune and destroy all old leaves and branches of roses, ornamentals and shade trees. If applied in spring after buds show new growth, injury may result. Do not apply when freezing temperatures are expected and do not apply to evergreens. Protect painted surfaces from spray, as staining or bleaching will result if sprayed.

Shake thoroughly before using as separation may occur in bottle. Remixing reconstitutes the product effectively. May be used in hose-end, tank or hand sprayers. Add small amount of water to sprayer, add correct amount of Oil & Lime Sulphur Spray, mix and add balance of water to the sprayer. Keep mixture agitated while spraying. Wet pests and all parts of plant thoroughly with penetrating spray.