

CAUTION:

Avoid prolonged or repeated breathing of spray. Do not take internally. Skin contact may be harmful, avoid contact. Wash before eating or smoking and after use. Wear regular long sleeved work clothing. Do not reuse clothing without cleaning.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor; atropine is antidotal.

DO NOT CONTAMINATE FOOD, FEED FEEDING TROUGHS AND WATER RECEPTACLES.

PRODUCT FREEZING TEMPERATURE 32° F; satisfactory utility is restored after thawing and mixing of product stored at temperatures as low as 0° F. Repeated 0-10° freeze thawing may impair utility.

Replace cover on container to prevent drying.

PRODUCT INFORMATION

Sevin 4 Flowable is an extremely fine dispersion of Sevin in water. A dilute spray of Sevin 4 Flowable is tenacious and quick-setting, giving an adhesive residue to plants on partial drying. Sevin is compatible with commonly used insecticides such as malathion and other organic phosphates and botanicals; also with such fungicides as fixed coppers, dithiocarbamates, sulphur, Captan, Cyrex, Glyodin, and Phaltan. Sevin does not control spider mites, but is compatible with all common miticides. Sevin is unstable under highly alkaline conditions. Do not use with lime, Bordeaux, lime-sulphur, casein-lime spreaders or other alkaline materials.

OBSERVE THE FOLLOWING USE PRECAUTIONS

For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. This product is highly toxic to bees exposed to direct treatment or residues on crops.

48 teaspoons = 16 tablespoons = 8 fl. ounces = 1/2 pint = 1/4 quart

Buyer's guarantee limited strictly to label claims.

Manufactured for

BCHIDE CHEMICAL COMPANY, INC., UTICA, NEW YORK 13502


**SEVIN 4 LB.
FLOWABLE INSECTICIDE**

**AN AQUEOUS SUSPENSION OF AGRICULTURAL
INSECTICIDE CONTAINING 4 POUNDS
SEVIN PER GALLON**

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate) 41.5%
INERT INGREDIENTS 58.2%
U.S. Pat. No. 2,903,478 Sevin is a trademark of Union Carbide Corp.

USDA Reg. No. 4-167

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS.
HARMFUL IF INHALED OR SWALLOWED.** 1 QT. NET
(See side panel cautions)



DIRECTIONS FOR USE

BEFORE USING SHAKE OR STIR WELL: Pour recommended amount of this material into nearly filled spray tank, or pre-mix in a bucket half-filled with water and pour mixture into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids unless previous experience has proven them physically compatible and safe to use.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. (Do not use on Boston Ivy)—Apply 1 qt. per 100 gal. water (2 level teaspoonfuls per gal.) when insects or their damage appear. Repeat weekly or as needed to control: aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, roseslug, scale crawlers, tent caterpillars, thrips (exposed), willow leaf beetles.

ADULT MOSQUITO CONTROL: For use on pasture and rangelands, non-agricultural lands, such as yards, parks, recreation areas, logging camps, military posts and adjacent forested lands or wastelands: Use 1/2 to 1 pt. per acre in suitable low pressure ground equipment. For all methods apply in sufficient water to effectively cover area to be treated. By airplane use in 3-5 gallons of water per acre. For optimum control apply in early morning or evening when air is calm and adult mosquitoes are active.

Use 1 quart per 100 gallons in hydraulic sprayers or 2 teaspoons per gallon in hand sprayers for treating backyards or other limited areas. Thoroughly cover lower shade tree foliage, shrubbery, flower beds, grass and shaded areas around buildings where adult mosquitoes congregate. Do not apply to water. Repeat at 7 to 10-day intervals or as necessary. Sevin injures Boston Ivy. Do not use on this ornamental.

LAWN INSECT CONTROL: Use one quart in 150 to 200 gallons of water for each 5,000 square feet of established lawn area for control of Chinch bugs, leafhoppers, sod webworms (lawn moths), fall armyworm, cutworms, cent-wigs, ants, fleas, mosquitoes and millipedes. In Florida use 3 pints for chinch bug control.

Use garden hose sprayers or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2-3 weeks later if necessary.