

DIRECTIONS

Apply when weeds are actively growing (temperatures above 50°F). Select period of little or no wind when rain is not expected for 8-12 hours after application. Rain or watering within 6 hours after application will interfere with results necessitating additional application. Shake can before and occasionally during application. Aim arrow on button at weed to be treated and away from you. Not necessary to stoop over weed but avoid spattering or accidental spraying of nearby desirable plants. In case of accidental application to plants dilute foam immediately with hose or sprinkling can. A once-over, light covering of weeds is sufficient. If certain resistant varieties are not dead in 3 weeks, repeat application. Do not mow lawn weeds for 2 days after application.

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

BONIDE

THE FEDERAL BUREAU OF INVESTIGATION HAS CONDUCTED TESTS AND HAS DETERMINED THAT THIS PRODUCT IS NOT A TOXIC POISON SUBSTANCE.

WEED KILLER

FOAMING JET ACTION

WITH 2,4-D & SILVEX

POWERFUL SPOT KILLER

FOR LAWN WEEDS

NO STOOPING

ACTIVE INGREDIENTS

2,4-D (2,4-DICHLOROPHENOXYACETIC ACID) 1.0%

SILVEX (2,4,5-TRICHLOROPHENOXYACETIC ACID) 0.5%

INERT INGREDIENTS 98.5%

CONTENTS UNDER PRESSURE

CAUTION: KEEP OUT OF REACH OF CHILDREN

1.1 LBS. NET WT.

LB. 6 07 NET

Broadleaf Weed Killer provides quick, simple spot kill of many lawn weeds. Foaming action slows where application has been made and reduces drift of spray which can damage nearby desirable plants. Broadleaf Weed Killer with 2,4-D and Silvex gets common chickweed, narrow leaf chickweed, cinquefoil, clovers, dandelion, dock, field bindweed, ground ivy, henbit, knotweed, annual morning glory, oxalis, pennywort, broadleaf plantain, buckhorn plantain, poison ivy, poison oak, purslane, spotted purple, thistles.

CAUTION: Always use with care. Do not use on lawn if rain is expected within 24 hours after use. Wash thoroughly. Store out of reach of children. Do not use when temperature is near 32°F or near fertilizers, seeds, insecticides or fungicides. Do not puncture the can or fill with empty cans and expose to fire. Avoid freezing. May be more susceptible plants and reduce toxicity of application if any, to creeping grasses, Dichondra and clover lawns. Do not apply to newly seeded lawns. Do not re-apply to lawn more than twice.

DIRECTIONS

Apply when weeds are actively growing (temperatures above 60°F). Select period of little or no wind when rain is not expected for 8-12 hours after application. Rain or watering within 6 hours after application will interfere with results necessitating additional application. Shake can before, and occasionally during application. Aim arrow on button at weed to be treated and away from you. Not necessary to stoop over weed but avoid spattering or accidental spraying of nearby desirable plants. In case of accidental application to plants, dilute foam immediately with hose or sprinkling can. A once over, light covering of weeds is sufficient. If certain resistant varieties are not dead in 3 weeks, repeat application. Do not mow lawn weeds for 2 days after application.

Buyer's Guarantee Limited to Label Claims

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

BONIDE

ACCURATE

John

• MEET THE
FUNGICIDE A
FOR ECONOM

FOAMING

WEED

WITH 2

POWERFUL

FOR LAWN

NO SPOTS

ACTIVE INGREDIENTS:

* Triethanolamine Salt of 2,4-D

* Triethanolamine Salt of Silvex

(propionic acid)

INERT INGREDIENTS:

Equivalent to 0.75% (0.41% ch
Equivalent to 0.75% (Silvex
propionic acid)

CONTENTS

CAUTION: KEEP OUT OF REACH OF CHILDREN

See back

1 LB