

CERTIFICATION OF COMPLIANCE

Amendment to the Registration Standard for
2,4-dichlorophenoxyacetic acid (2,4-D) and
its inorganic salts, amines and esters

February 7, 1990

I, being an authorized representative of Bonide Products, Inc. understand the labeling requirements specified in the Registration Standard for 2,4-Dichlorophenoxyacetic Acid as amended and included therein by reference. Further, I certify that all containers of Bonide Lawn Weed Killer w/Banvel D, EPA Reg. No. 4-169, released for shipment after March 1, 1991, will bear revised labeling in accordance with those requirements.

Annabel Olson
Annabel Olson

Registration Specialist
Title

March 14, 1990
Date

455

YEL0 295
BONIDE®

TREATS 10,000 SQUARE FEET

A 100 Foot by 100 Foot Lawn Area—Net Weight 14 lbs.

Just Pour in Spreader and Walk • Clean, Easy to Use, Dry Granules

YEL0
**LAWN
WEED KILLER**

W/BANVEL® D & 2,4-D

LAWN WEED KILLER

WBANVEL D & 2,4-D

kills...

Dandelion, plantain, buckhorn, thistle, sow thistle, poison ivy, poison oak, ground ivy, ragweed, burdock, nettle, pennywort, pigweed, healall, sheep sorrell, yarrow, wild onion, purslane, chicory, gypsum weed, black medic and white clover in established lawns.

E.P.A. Est. 4-NY-1

E.P.A. Reg. No. 4-169

ACTIVE INGREDIENTS:

*Dimethylamine salt or 2,4-dichlorophenoxyacetic acid	3.0%
**Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.6%
***Dimethylamine salts of related acids	0.2%
INERT INGREDIENTS:	96.3%

100.00%

*Equivalent to 2,4-Dichlorophenoxyacetic acid 2.5%.

**equivalent to 3,6-dichloro-o-anisic acid (dicamba) 0.5%

***Banvel is a registered trademark of the Velsicol Chemical Co., Chicago

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

ENVIRONMENTAL HAZARD

~~This product is toxic to aquatic invertebrates.~~ Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters. Avoid contamination of feed or foodstuffs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Securely wrap original container in several layers of newspaper and discard in trash.
Store in a cool dry area but not near seeds, fertilizers, insecticides or fungicides.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds. On Bermuda, St. Augustine, bent, Dichondra and carpet grass lawns spot treat only as injury may occur. Avoid application on or near desirable trees and other plants, or on areas over their root systems or there movement and washing may cause contact with roots.

~~This product can reach groundwater from improper handling. To minimize groundwater contamination from spills during loading and cleaning of equipment, take the following steps:~~

~~Handling: When handling this product, wear chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label.~~

~~Cleaning of Equipment: When cleaning equipment, do not pour the washwater on the ground, spray or drain over a large area away from wells and other water sources.~~

DIRECTIONS

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Weeds may be spot treated by lightly sifting direct from bag. Spreader settings for 1/4 inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. Push shut-off open when re-starting. Keep machine handle low or container of machine vertical to ground level. In case of accidental heavy overdose, watering the over dosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded three weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

LAWN WEED KILLER

Lawn Weed Killer can be applied any time spring, summer or fall for general weed control. Application during warm periods 60-85° when weeds are actively growing gives best results. During cool or cold periods action is slow and retreatment may be necessary. Do not apply to a new lawn until after second regular mowing.

Starting Point suggested spreader settings:

Make & Model	Setting	Make & Model	Setting
Scott	8	Montgomery Ward 89-1081	8 1/4
Jackson 18-26 18"	7	Salem Micro Set 22"	7
Scars Products Come Country 18"	7	Curtina R R' Swath	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Securely wrap original container in several layers of newspaper and discard in trash.
Store in a cool dry area but not near seeds, fertilizers, insecticides or fungicides.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds. On Bermuda, St. Augustine, bent, Dichondra and carpet grass lawns spot treat only as injury may occur. Avoid application on or near desirable trees and other plants, or on areas over their root systems or their movement and washing may cause contact with roots.

~~This product can reach groundwater from improper handling. To minimize groundwater contamination from spills during loading and cleaning of equipment, take the following steps:~~

~~Handling: When handling this product, wear chemical resistant gloves. Wash non-disposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label.~~

~~Cleaning of Equipment: When cleaning equipment, do not pour the wash water on the ground; spray or drain over a large area away from wells and other water sources.~~

DIRECTIONS

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Weeds may be spot treated by lightly sifting direct from bag. Spreader settings for 1/4 inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. Push shut-off open when re-starting. Keep machine handle low or container of machine vertical to ground level. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded three weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

LAWN WEED KILLER

Lawn Weed Killer can be applied any time spring, summer or fall for general weed control. Application during warm periods 60-85° when weeds are actively growing gives best results. During cool or cold periods action is slow and retreatment may be necessary. Do not apply to a new lawn until after second regular mowing.

Starting Point suggested spreader settings:

Make & Model	Setting	Make & Model	Setting
Scott.....	8	Montgomery Ward 89-1081.....	8 1/4
Jackson 18-26 18".....	7	Salem Micro Set 22".....	7
Sears-Roebuck Cross Country 19".....	7	Cyclone B 5' Swath (Still Air Only).....	6
Lawn Beauty.....	7		

Buyer's Guarantee Limited to Label Claims

Manufactured by

BONIDE CHEMICAL CO., INC. • YORKVILLE, NY 13495

