

**CHERRIES (RED TART):** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals making last application to 14 days after harvest.

**CITRUS (ORANGES, GRAPEFRUIT, LEMONS, LIMES, TANGERS, TANGERINES):** Make first application at the first appearance of symptoms. Repeat at 7 to 10 day intervals. Do not apply within 3 days of harvest.

**CRANBERRIES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals. Do not apply within 3 days of harvest.

**CUCUMBERS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals. Do not apply within 3 days of harvest.

**LETTUCE:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**ONIONS, GARLIC, LEEKS, SHALLOTS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**STRAWBERRIES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**TOMATOES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**ROSES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**CHRYSANTHEMUMS, IRIS, DEPTEROCARPUS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**CARNATIONS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**ZINNIAS, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**POINSETTIA (GREENHOUSE):** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**AZALEA CUTTINGS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

Packed by  
**BONIDE CHEMICAL COMPANY, INC., UTICA, N. Y. 13502**

# BONIDE

## FOLPET 50% WP

### A FUNGICIDE FOR PLANT DISEASE CONTROL

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 Avoid inhalation of dust or spray. Avoid contact with skin. Avoid contamination of feed or foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Burn empty bag and keep out of smoke. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**1 LB. NET WEIGHT**

#### COMPATIBILITY

FOLPET 50 WP may be used in conjunction with other fungicides and insecticides. Do not use in conjunction with other materials which are known to be incompatible.

#### DIRECTIONS FOR USE

Four required amount of this material is sufficient for use in a spray tank. Add to the spray tank in the order listed. Mix thoroughly through the tank. Add to the spray tank in the order listed. Mix thoroughly through the tank. Add to the spray tank in the order listed. Mix thoroughly through the tank.

#### RECOMMENDATIONS

Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**APPLES, CRABAPPLES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**AVOCADOS:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**BLUEBERRIES, HUCKLEBERRIES:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**CANTALOUPE, MUSKMELONS, HONEYDEW MELONS, WATERMELONS, SUMMER SQUASH, PUMPKINS, WINTER SQUASH:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**CELERY:** Apply 1/2 lb. per acre. Apply at the first appearance of symptoms. Repeat at 7 to 10 day intervals until harvest.

**CHERRIES (RED FART):** Fruit Rot - 4 lbs. per 100 gals. water plus a spreader. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Apply to foliage post harvest. Downy Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Apply to foliage post harvest. Sweet cherries as for above. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Apply to foliage post harvest.

**CITRUS (ORANGES, GRAPEFRUIT, LEMONS, LIMES, TANGELUS, TANGERINES):** Mealybugs - 4 lbs. per 100 gals. water plus a spreader. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Apply to foliage post harvest. The full flush in August. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Apply to foliage post harvest.

**CRANBERRIES:** Storage Rot - 4 lbs. per 100 gals. water plus a spreader. Apply during bloom period. Repeat at 7 to 10 day intervals until harvest. Do not apply within 7 days of harvest.

**CUCUMBERS:** Fruit Rot - Rhizoctonia and other - 4 to 8 lbs. per 100 gals. water plus a spreader. Apply in a minimum of 100 gals. water per acre. Directly spray at the base of the row. Apply when plants are beginning to fruit. Downy Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply in a minimum of 100 gals. water per acre. Directly spray at the base of the row. Apply when plants are beginning to fruit. Powdery Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply in a minimum of 100 gals. water per acre. Directly spray at the base of the row. Apply when plants are beginning to fruit. Anthracnose - 4 lbs. per 100 gals. water plus a spreader. Apply in a minimum of 100 gals. water per acre. Directly spray at the base of the row. Apply when plants are beginning to fruit. Start application as soon as plants are above the ground. Make repeat applications at 7 to 10 day intervals as necessary.

**GRAPES:** Downy Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply 250 to 300 gals. per acre. Least 400 gals. per acre. Make first application when new growth is 1 to 2 inches long. Repeat when growth is 1 to 2 inches long. Black Rot - 4 lbs. per 100 gals. water plus a spreader. Apply 250 to 300 gals. per acre. Least 400 gals. per acre. Make first application just before bloom. Repeat immediately after bloom. Depending upon the susceptibility of the grape variety, from 1 to 3 more sprays may be needed at 7 to 10 day intervals. Powdery Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply 250 to 300 gals. per acre. Least 400 gals. per acre. Make first application 7 to 10 weeks after bloom and repeat at 2 week intervals as needed. Note: During extended periods of hot dry weather, foliage injury to several varieties of grapes may occur.

**GOOSEBERRIES, CURRANTS:** Leaf Spot - 4 lbs. per 100 gals. water plus a spreader. Apply 250 gals. per acre. Make first application just before bloom period. Repeat at 7 to 10 day intervals in the post bloom period.

**LETTUCE:** Downy Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply up to 200 gals. per acre. Make first application when plants are established in field. Repeat at 7 day intervals until harvest.

**ONIONS, GARLIC, LEEKS, SHALLOTS:** Purple Blotch and Downy Mildew - 4 lbs. per 100 gals. water plus a spreader. Apply 200 gals. per acre. Make first application before first infection. Repeat at weekly intervals until tops begin to dry before harvest.

**STRAWBERRIES:** Leaf Spot, Fruit Rot - 2 lbs. per 100 gals. water plus a spreader. Apply up to 200 gals. per acre. Make first application before bloom and repeat at 7 day intervals until after harvest.

**TOMATOES:** Anthracnose - 4 lbs. per 100 gals. water plus a spreader. Use up to 200 gals. per acre. Make first application during first bloom and repeat at 7 to 10 day intervals until fruit is harvested.

**ROSES:** Black Spot, Powdery Mildew - 2 lbs. per 100 gals. water plus a spreader. 2 tablespoonfuls per 1 gallon of water. Make first application when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Make thorough application. Do not use wetting agents on roses as injury may result.

**CHRYSANTHEMUMS, IRIS:** Septoria Leafspot, Didymella Leafspot - 2 lbs. per 100 gals. water plus a spreader. 2 tablespoonfuls per 1 gal. of water. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as needed throughout the season.

**CARNATIONS:** Rust, Alternaria Leafspot - 2 lbs. per 100 gals. water plus a spreader. 2 tablespoonfuls per 1 gal. of water. Begin applications 2 weeks after planting and continue bi weekly applications for 3 to 4 applications.

**ZINNIAS, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS:** Alternaria Leafspot, Powdery Mildew, Anthracnose - 2 lbs. per 100 gals. water plus a spreader. 2 tablespoonfuls per 1 gal. of water. Treat weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity.

**POINSETTIA (GREENHOUSE):** Pythium Root Rot - 2 lbs. per 100 gals. water plus a spreader and use 1 pint per 1000 cu. ft. of space. Make first application at time of planting. Repeat in 10 days.

**AZALEA CUTTINGS:** Dipping off - 2 lbs. per 100 gals. water plus a spreader. Allow to soak for 10 to 15 minutes before placing in rooting media.

Buyer's Guarantee Limited to Label Claim

Packed by

BONIDE CHEMICAL COMPANY, INC., UTICA, N. Y. 13502

# BONIDE

## FOLPET 50% W

### A FUNGICIDE FOR PLANT DISEASE CONTROL

ACTIVE INGREDIENT  
Folpet, 50% (Folpet 50% W)  
NET WT. 1 LB. (454 G)  
EPA REG. NO. 101-108-01

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
Avoid inhalation of dust or spray. Avoid contact with skin. Avoid contamination of foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Burn or dispose of in accordance with local, state, and federal regulations. Do not contaminate water, soil, or air. Clean equipment, or disposal of wastes.

1 LB. NET WEIGHT

# BONIDE

## FOLPET 50% WP

### A FUNGICIDE FOR PLANT DISEASE CONTROL

ACTIVE INGREDIENT  
Folpet, N-(1-hydroxypropyl)-N'-methyl-2-pyrrolidone 50%  
NET WEIGHT 1 LB. 5 OZ.  
Pat. No. 2,811,111

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
Avoid inhalation of dust or spray. Avoid contact with skin. Avoid contamination of feed or foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Burn empty bag and keep out of smoke. Do not contaminate water by cleaning of equipment, or disposal of wastes.

1 LB. NET WEIGHT

#### COMPATIBILITY

FOLPET 50-WP may be combined with most commonly used insecticides and plant and fungicides. Fungicidal activity of FOLPET is enhanced if combined with strongly alkaline water such as hydrated lime. This material should not be used in combination with the following or sprays: Do not use with DDT when a concentrate material is to be applied.

#### DIRECTIONS FOR USE

Pour required amount of this material on surface of water in nearly filled spray tank or pre-mix in a bucket half filled with water and pour mixture through screen into nearly filled spray tank. Add balance of water to the tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible. Consult State Agricultural Experiment Station or State Extension Service Specialist for specific information as the timing, number and rate of application needed will vary with local conditions.

The Federal Food and Drug Administration has set official tolerances for the use of pesticides. Enter that residue on food and for use of low-moisture products. State to obtain use only at recommended rates.

#### RECOMMENDATIONS

Unless otherwise stated, tolerances are given in terms of pounds of Folpet 50-WP per 100 gallons of water.  
**APPLES, CRABAPPLES:** Bitter Rot, Botryphaeria, Alternaria Spot, Blight, Fruit Rot, Sooty Blotch, Bark Rot, Fly Speck and other diseases. Apply 1 to 2 lbs. per acre. Make first application at delayed dormancy and repeat at 7-14 day intervals until harvest. (A TION, East.) Use during the entire growing period but 10-15 days after petal fall may cause russeting on sensitive varieties such as Golden Delicious, Red Delicious and Jonathan. Waxap.  
**AVOCADOS:** Scab. 1 to 2 lbs. Make a 100-gallon at 2 to 3 week intervals as needed.  
**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES:** Anthracnose, Mildew, Blight, Botrytis, and other diseases. Make first application when berries have set and repeat at 7-14 day intervals until harvest.  
**BLUEBERRIES, HUCKLEBERRIES:** Botrytis, Mildew, Powdery Mildew, Leaf Spot, and other diseases. Apply 1 to 2 lbs. per 100 gallons of water. Make first application at 7-14 day intervals until harvest.  
**CANTALOUPE, MUSKMELONS, HONEYDEW MELONS, WATERMELONS, SUMMER SQUASH, PUMPKINS, WINTER SQUASH:** Mildew, Powdery Mildew, Blight, and other diseases. Apply 1 to 2 lbs. per 100 gallons of water. Make first application at 7-14 day intervals until harvest.  
**CELERY:** Early Blight, Late Blight, Leaf Spot, and other diseases. Apply 1 to 2 lbs. per 100 gallons of water. Make first application at 7-14 day intervals until harvest.