

300 176497 182
17 1 4-181

SEP 30 1986

Bonide Chemical Company, Inc.
2 Wurz Avenue
Yorkville, NY 13495

Gentlemen:

Subject: Bonide Folpet 50% WP
EPA Registration No. 4-181
Your Submission Dated January 20, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comment listed below. Five copies of the finished labeling must be submitted.

- Add the appropriate Statements of Practical Treatment for your product.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

88773: Jacoby: J-3; KENCO: 9/18/86; 9/30/86; bjp: vo

CONCURRENCES

SYMBOL ▶							
SURNAME ▶							
DATE ▶							

201-4

BONIDE®

FOLPET 50% WP

A FUNGICIDE FOR PLANT DISEASE CONTROL

ACTIVE INGREDIENT:
Folpet (1-(4-chlorophenyl)thio phthalimide) 50%
INERT INGREDIENTS: 50%
Pat. No. 2,553,770, 2,553,771. Also known as Phialen.
EPA Est. 4-NV-1 EPA Reg. No. 4-181

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See label panel for additional precautionary statements.

NET CONTENTS: 1 LB.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION
Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Avoid breathing spray or mist. Wash thoroughly after handling. Wash contaminated clothing before reuse.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Storage: Keep pesticide in original container. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area, inaccessible to children and pets.
Container Disposal: Securely wrap original container in several layers of paper and discard in trash. Do not reuse container.

DIRECTIONS FOR USE
Pour required amount of this material on surface of water in nearly filled spray tank, or pre-mix in a bucket half-filled with water and pour mixture through screen into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.
Consult State Agriculture Experiment Station or State Extension Service Specialist for specific information as the timing, number and rate of application needed will vary with local conditions.
The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

RECOMMENDATIONS
Unless otherwise stated, dosages are given in terms of pounds of Folpet 50-WP per 100 gallons of water.
ROSES: Black Spot, Powdery Mildew — 2 lbs. per 100 gals. (2 tablespoonfuls per 1 gal. of water). Make first application when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Make thorough application. Do not use wetting agents on roses as injury may result.
CHRYSANTHEMUMS, IRIS: Septoria Leafspot, Didymellina Leafspot — 2 lbs. per 100 gals. (2 tablespoonfuls per 1 gal. of water). Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as needed throughout the season.
CARNATIONS: Rust, Alternaria Leafspot — 2 lbs. per 100 gals. (2 tablespoonfuls per 1 gal. of water). Begin applications 2 weeks after planting and continue bi-weekly application for 3 to 4 applications.
ZINNIAS, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX, SNAPDRAGONS: Alternaria Leafspot, Powdery Mildew, Anthracnose — 2 lbs. per 100 gals. (2 tablespoonfuls per 1 gal. of water). Treat weekly during periods of rapid new growth and severe infection, and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity.
POINSETTIA (GREENHOUSE): Pythium, Root Rot — 2 lbs. per 100 gals. water plus a spreader and use 1 pint per 8 inch pan. Make first application at time of panning. Repeat in 10 days.
AZALEA CUTTINGS: Damp-off — 1½ to 2 lbs. per 100 gals. Allow to soak for 15 to 30 minutes before placing in rooting media.

COMPATIBILITY
FOLPET 50-WP may be combined with most commonly used insecticides, adjuvants and fungicides. Fungicidal activity of FOLPET is not reduced when used with strongly alkaline materials such as hydrated lime. This material should not be used in combination with or closely following oil sprays. Do not use with TEPP when a concentrate mixture is being applied.

**ACCEPTED
with COMMENTS**

Buyer's Guarantee Limited to Label Claims
Packed by BONIDE CHEMICAL CO., INC., YORKVILLE, N.Y. 13495

SEP 30 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 4-181