

CAUTION: Keep out of reach of children. Avoid breathing spray and contact with eyes or skin. May be absorbed through the skin. In case of contact and after use, wash with soap and water. Avoid contamination of feed and foodstuffs.

Cover fish bowls or aquariums near use areas before use. Do not spray Maidenhair Fern, Lantana, Cyclamen or apply to household pets or humans.

Buyer's guarantee limited to label claims.

ACTIVE INGREDIENTS
Carbaryl (1-naphthyl N-methylcarbamate)
Folpet (N-trichloromethylthio-1,1-dithio-2-urea)
Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate)
1,1-bis-(p-chlorophenyl)-2,2,2-trichloroethanol

EPA Reg. No. 4-124
7500
5000
5000
0750
98-1750

INERT INGREDIENTS

Sevin is registered trademark of Union Carbide Corp.
Phaltan is a registered trademark for Phaltimide fungicide. U.S. Patent Nos. 2,553,770; 2,553,771; 2,553,776.
Kelthane is registered trademark of Rohm & Haas Co.

Mfg. by BONIDE CHEMICAL CO., INC., UTICA, N.Y. 13502

ACCEPTED

U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF PLANT INDUSTRY
WASHINGTON, D.C. 20250
SUBJECT: PESTICIDES
DATE: 10/15/58

BONIDE

Rose & Floral SPRAY BOMB

1 LB. NET

Flowers (Roses) — Ornamentals

House Plants (African Violets)

Ready to Use Indoors-Outdoors

CAUTION: KEEP OUT OF REACH OF CHILDREN

(See side panel cautions)

WARNING: Contents under pressure. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

FLORA-BOMB IS A MULTI-PURPOSE SPRAY for house plants, small gardens, and may be used in large gardens for spot treatment to prevent spreading of localized infestations to larger areas. The formula is designed for use on African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Delphiniums, Dogwood, English Ivy, Euonymus, Fuchsia, Geraniums, Crassula, Grape vine, Kentia palm, Laurel, Marigolds, Rhododendrons, Roses, Rubber plants, Snapdragons, Stocks, Wandering Jew and Zinnias. Insecticide for aphids, spider and clover mites, exposed thrips, lace bugs, flea beetle, bag worms, tent caterpillars, leaf hoppers, rose chafer, Japanese beetles, pavement ants, and army worms. Fungicide for black spot and powdery mildew on roses, alternaria leaf spot and rust on carnation, and powdery mildew and septoria leaf spot on Chrysanthemums.

DIRECTIONS

Shake can before use, remove cap, hold in upright position with spray opening (at end of arrow on button) pointed toward object to be sprayed. Keep spray opening and can 14 (fourteen) inches away from foliage. Press button downward to release mist. Apply mist to all parts of plant, top and undersides of foliage. Stop application before droplets form on leaf surfaces or liquid starts to drip off foliage. Apply as preventative before diseases (black spot, powdery mildew, alternaria leaf spot, rust, septoria leaf spot) occur and every 5-7 days during warm, moist periods when diseases are prevalent. Properly treated foliage will be protected for about one week.